

Beam Single Spin Asymmetries in Electron-Proton Scattering

Pratik Sachdeva

Washington University in St. Louis

Mentors:

Wally Melnitchouk

Jefferson Laboratory Theory Center

Peter Blunden

University of Manitoba

July 23, 2014



U.S. DEPARTMENT OF
ENERGY

Office of
Science

The logo for Jefferson Lab, featuring a stylized red and yellow swoosh above the text "Jefferson Lab".
Jefferson Lab
Thomas Jefferson National Accelerator Facility

Outline

- 1 Introduction
 - Beam Single Spin Asymmetries (BSSA)
 - Transverse and Normal Polarizations
 - Motivation: Q-weak Collaboration
- 2 Elastic Scattering
 - Beam Normal SSA
 - Beam Transverse Single Spin Asymmetry
 - Combination of Asymmetries
- 3 Inelastic Hadronic Intermediate State
 - $\mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma}$ Interference
 - Leptonic and Hadronic Tensors
 - Normal BSSA
- 4 Conclusion

Introduction

Beam Single Spin Asymmetries (BSSA)

- Polarized electron beam (longitudinal, transverse, normal), but unpolarized target.

Introduction

Beam Single Spin Asymmetries (BSSA)

- Polarized electron beam (longitudinal, transverse, normal), but unpolarized target.
- In general the beam asymmetry, B , is

$$B = \frac{\sigma_{\uparrow} - \sigma_{\downarrow}}{\sigma_{\uparrow} + \sigma_{\downarrow}}.$$

Introduction

Beam Single Spin Asymmetries (BSSA)

- Polarized electron beam (longitudinal, transverse, normal), but unpolarized target.
- In general the beam asymmetry, B , is

$$B = \frac{\sigma_{\uparrow} - \sigma_{\downarrow}}{\sigma_{\uparrow} + \sigma_{\downarrow}}.$$

- BSSA disappear for one-photon exchange.

What exchanges can cause BSSA?

What exchanges can cause BSSA?

Consider the scattering amplitude:

$$\begin{aligned} |\mathcal{M}|^2 &= |\mathcal{M}_\gamma + \mathcal{M}_Z + \mathcal{M}_{\gamma\gamma} + \dots|^2 \\ &= |\mathcal{M}_\gamma|^2 + \mathcal{M}_\gamma^* \mathcal{M}_Z + \mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma} + \dots \end{aligned}$$

What exchanges can cause BSSA?

Consider the scattering amplitude:

$$\begin{aligned} |\mathcal{M}|^2 &= |\mathcal{M}_\gamma + \mathcal{M}_Z + \mathcal{M}_{\gamma\gamma} + \dots|^2 \\ &= |\mathcal{M}_\gamma|^2 + \mathcal{M}_\gamma^* \mathcal{M}_Z + \mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma} + \dots \end{aligned}$$

- At the Born level, the spin orientation has no effect on $|\mathcal{M}_\gamma|^2$, so the asymmetry vanishes.

What exchanges can cause BSSA?

Consider the scattering amplitude:

$$\begin{aligned} |\mathcal{M}|^2 &= |\mathcal{M}_\gamma + \mathcal{M}_Z + \mathcal{M}_{\gamma\gamma} + \dots|^2 \\ &= |\mathcal{M}_\gamma|^2 + \mathcal{M}_\gamma^* \mathcal{M}_Z + \mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma} + \dots \end{aligned}$$

- At the Born level, the spin orientation has no effect on $|\mathcal{M}_\gamma|^2$, so the asymmetry vanishes.
- Higher order effects result in asymmetries dependent on whether the spin is transversely, normally, or longitudinally polarized.

Transverse and Normal Polarization

- Transverse asymmetry B_t behaves like $\cos \phi_s$ and normal asymmetry B_n behaves like $\sin \phi_s$.





Figure 2. Electron-Proton Scattering

Motivation: Q-weak Collaboration

- Precise measurement of the weak charge of the proton, Q_W^P , using asymmetry produced by longitudinally polarized electrons.


Motivation: Q-weak Collaboration

- Precise measurement of the weak charge of the proton, Q_W^p , using asymmetry produced by longitudinally polarized electrons.
- Transversely polarized electrons: BSSA has azimuthal (ϕ) dependence (Buddhini Waidyawansa, CIPANP 2012 Conference).



Motivation: Q-weak Collaboration

- Precise measurement of the weak charge of the proton, Q_W^p , using asymmetry produced by longitudinally polarized electrons.
- Transversely polarized electrons: BSSA has azimuthal (ϕ) dependence (Buddhini Waidyawansa, CIPANP 2012 Conference).



- Concerned about possible phase offsets due to unconsidered BSSA effects.

Approach

- Elastic Scattering Contribution

Approach

- Elastic Scattering Contribution
 - Interference of one- and two-photon exchange.
 - Interference of one-photon and Z exchange.

Approach

- Elastic Scattering Contribution
 - Interference of one- and two-photon exchange.
 - Interference of one-photon and Z exchange.
- Hadronic Inelastic Intermediate State


Approach

- Elastic Scattering Contribution
 - Interference of one- and two-photon exchange.
 - Interference of one-photon and Z exchange.
- Hadronic Inelastic Intermediate State
 - Pasquini and Vanderhaeghen (**PRC 70**, 2004) showed that inelastic hadronic intermediate states have a larger contribution to the asymmetry than the elastic case.
 - Consider the case of near forward limit.

Elastic Scattering

Elastic scattering for electron-proton scattering,

$$e(p_1) + N(p_2) \rightarrow e(p_3) + N(p_4).$$



Kinematic Variables

Elastic scattering process:

$$e(p_1) + N(p_2) \rightarrow e(p_3) + N(p_4)$$

Using the following notation for kinematic variables:

Kinematic Variables

Elastic scattering process:

$$e(p_1) + N(p_2) \rightarrow e(p_3) + N(p_4)$$

Using the following notation for kinematic variables:

$$P = \frac{p_2 + p_4}{2}, \quad K = \frac{p_1 + p_3}{2}, \quad q = p_1 - p_3, \quad Q^2 = -q^2;$$

$$s = (p_1 + p_2)^2, \quad \tau = \frac{Q^2}{4M^2}, \quad \nu = P \cdot K, \quad \epsilon = \frac{\nu^2 - M^4 \tau(1 + \tau)}{\nu^2 + M^4 \tau(1 + \tau)};$$

$$W^2 = (p_2 + q_1)^2, \quad Q_1^2 = -q_1^2, \quad Q_2^2 = -q_2^2,$$

for electron mass m_e and hadron mass M .

Beam Normal Single Spin Asymmetry: Setup

Using six invariant amplitudes given by Goldberger et al. (Ann. Phys. 2, 1957), we can construct a general elastic lepton-nucleon scattering amplitude:

$$T = T^{\text{non-flip}} + T^{\text{flip}},$$

Beam Normal Single Spin Asymmetry: Setup

Using six invariant amplitudes given by Goldberger et al. (Ann. Phys. 2, 1957), we can construct a general elastic lepton-nucleon scattering amplitude:

$$T = T^{\text{non-flip}} + T^{\text{flip}},$$

where

$$T^{\text{non-flip}} = \frac{e^2}{Q^2} \bar{u}_e(p_3) \gamma_\mu u_e(p_1) \cdot \bar{u}_N(p_4) \left(\tilde{G}_M \gamma^\mu - \tilde{F}_2 \frac{P^\mu}{M} + \tilde{F}_3 \frac{\not{K} P^\mu}{M^2} \right) u_N(p_2)$$

and

Beam Normal Single Spin Asymmetry: Setup

Using six invariant amplitudes given by Goldberger et al. (Ann. Phys. 2, 1957), we can construct a general elastic lepton-nucleon scattering amplitude:

$$T = T^{\text{non-flip}} + T^{\text{flip}},$$

where

$$T^{\text{non-flip}} = \frac{e^2}{Q^2} \bar{u}_e(p_3) \gamma_\mu u_e(p_1) \cdot \bar{u}_N(p_4) \left(\tilde{G}_M \gamma^\mu - \tilde{F}_2 \frac{P^\mu}{M} + \tilde{F}_3 \frac{\not{K} P^\mu}{M^2} \right) u_N(p_2)$$

and

$$T^{\text{flip}} = \frac{e^2}{Q^2} \frac{m_e}{M} \left[\bar{u}_e(p_3) u_e(p_1) \cdot \bar{u}_N(p_4) \left(\tilde{F}_4 + \tilde{F}_5 \frac{\not{K}}{M} \right) u_N(p_2) + \tilde{F}_6 \bar{u}_e(p_3) \gamma_5 u_e(p_1) \cdot \bar{u}_N(p_4) \gamma_5 u_N(p_2) \right].$$

Beam Normal Single Spin Asymmetry: Setup

In the two previous equations,

$$\tilde{G}_M, \tilde{F}_2, \tilde{F}_3, \tilde{F}_4, \tilde{F}_5, \tilde{F}_6$$

are complex functions of ν and Q^2 . In the Born approximation, these reduce to

Beam Normal Single Spin Asymmetry: Setup

In the two previous equations,

$$\tilde{G}_M, \tilde{F}_2, \tilde{F}_3, \tilde{F}_4, \tilde{F}_5, \tilde{F}_6$$

are complex functions of ν and Q^2 . In the Born approximation, these reduce to

$$\tilde{G}_M^{\text{Born}}(\nu, Q^2) = G_M(Q^2)$$

$$\tilde{F}_2^{\text{Born}}(\nu, Q^2) = F_2(Q^2)$$

$$\tilde{F}_{3,4,5,6}^{\text{Born}}(\nu, Q^2) = 0.$$

Beam Normal Single Spin Asymmetry: Setup

In the two previous equations,

$$\tilde{G}_M, \tilde{F}_2, \tilde{F}_3, \tilde{F}_4, \tilde{F}_5, \tilde{F}_6$$

are complex functions of ν and Q^2 . In the Born approximation, these reduce to

$$\tilde{G}_M^{\text{Born}}(\nu, Q^2) = G_M(Q^2)$$

$$\tilde{F}_2^{\text{Born}}(\nu, Q^2) = F_2(Q^2)$$

$$\tilde{F}_{3,4,5,6}^{\text{Born}}(\nu, Q^2) = 0.$$

We can also write

$$\tilde{G}_E = \tilde{G}_M - (1 + \tau)\tilde{F}_2.$$

Beam Normal SSA

Gorchtein et al. found that for a spin parallel (anti-parallel) to the normal polarization vector

$$S^\mu = (0, \vec{S}_n), \quad \vec{S}_n = (\vec{p}_1 \times \vec{p}_3) / |\vec{p}_1 \times \vec{p}_3|$$

there is a beam normal SSA B_n due to $\mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma}$. It is equal to

Beam Normal SSA

Gorchtein et al. found that for a spin parallel (anti-parallel) to the normal polarization vector

$$S^\mu = (0, \vec{S}_n), \quad \vec{S}_n = (\vec{p}_1 \times \vec{p}_3) / |\vec{p}_1 \times \vec{p}_3|$$

there is a beam normal SSA B_n due to $\mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma}$. It is equal to

$$B_n = \frac{2m_e}{Q} \sqrt{2\epsilon(1-\epsilon)} \sqrt{1 + \frac{1}{\tau} (G_M^2 + G_E^2)}^{-1} \\ \times \left\{ -\tau G_M \operatorname{Im} \left(\tilde{F}_3 + \frac{1}{1+\tau} \frac{\nu}{M^2} \tilde{F}_5 \right) - G_E \operatorname{Im} \left(\tilde{F}_4 + \frac{1}{1+\tau} \frac{\nu}{M^2} \tilde{F}_5 \right) \right\}.$$

Beam Normal SSA




Figure 3. Transverse Spin Polarization

Beam Normal SSA




Figure 3. Transverse Spin Polarization

If we consider a general transverse spin

$$\vec{s} = \cos \phi_s \hat{x} + \sin \phi_s \hat{y},$$

the general beam normal SSA can be written

Beam Normal SSA




Figure 3. Transverse Spin Polarization

If we consider a general transverse spin

$$\vec{s} = \cos \phi_s \hat{x} + \sin \phi_s \hat{y},$$

the general beam normal SSA can be written

$$B_{n, \text{gen}} = B_n \sin \phi_s.$$

Beam Transverse SSA

We found that a beam transverse SSA exists at the Born level due to $\mathcal{M}_\gamma^* \mathcal{M}_Z$. It is equal to

Beam Transverse SSA

We found that a beam transverse SSA exists at the Born level due to $\mathcal{M}_\gamma^* \mathcal{M}_Z$. It is equal to

$$B_t = \frac{G_F m_e}{2\pi\alpha} \frac{Q^2}{(s - M^2)} \sqrt{\frac{\epsilon(1 - \epsilon)}{\tau^2(\tau + 1)}} \left(G_M^2 + \frac{\epsilon}{\tau} G_E^2 \right)^{-1} \\ \times \left\{ g_V^e G_M G_M^Z \tau(\tau + 1) + g_A^e \left(G_M G_A^Z \tau(1 + \tau - \nu) + G_E G_E^Z (1 - \tau - \nu) \right) \right\},$$

where G_M^Z and G_E^Z are the weak form factors and G_F is Fermi's coupling constant.

Beam Transverse SSA

We found that a beam transverse SSA exists at the Born level due to $\mathcal{M}_\gamma^* \mathcal{M}_Z$. It is equal to

$$B_t = \frac{G_F m_e}{2\pi\alpha} \frac{Q^2}{(s - M^2)} \sqrt{\frac{\epsilon(1 - \epsilon)}{\tau^2(\tau + 1)}} \left(G_M^2 + \frac{\epsilon}{\tau} G_E^2 \right)^{-1} \\ \times \left\{ g_V^e G_M G_M^Z \tau(\tau + 1) + g_A^e \left(G_M G_A^Z \tau(1 + \tau - \nu) + G_E G_E^Z (1 - \tau - \nu) \right) \right\},$$

where G_M^Z and G_E^Z are the weak form factors and G_F is Fermi's coupling constant.

For a general transverse spin,

$$B_{t, \text{ gen}} = B_t \cos \phi_S.$$

Combination of Asymmetries

Combination of Asymmetries

- The interference between one- and two-photon exchange only produces a normal BSSA.
- The interference between one-photon and Z exchange only produces a transverse BSSA.

Combination of Asymmetries

- The interference between one- and two-photon exchange only produces a normal BSSA.
- The interference between one-photon and Z exchange only produces a transverse BSSA.

The combination of these two asymmetries gives

$$\begin{aligned} B &= B_t \cos \phi_s + B_n \sin \phi_s \\ &= \sqrt{B_t^2 + B_n^2} \sin(\phi_s + \delta) \end{aligned}$$

where $\delta = \tan^{-1} \left(\frac{B_t}{B_n} \right)$.

Combination of Asymmetries: Q-weak Kinematics

In Q-weak, B_n was measured as

$$B_n \approx -5.350 \text{ ppm.}$$

Combination of Asymmetries: Q-weak Kinematics

In Q-weak, B_n was measured as

$$B_n \approx -5.350 \text{ ppm.}$$

Using Q-weak kinematics, $Q^2 = 0.025 \text{ GeV}$, $E_1 = 1.125 \text{ GeV}$ (electron beam energy), we find

Combination of Asymmetries: Q-weak Kinematics

In Q-weak, B_n was measured as

$$B_n \approx -5.350 \text{ ppm.}$$

Using Q-weak kinematics, $Q^2 = 0.025 \text{ GeV}$, $E_1 = 1.125 \text{ GeV}$ (electron beam energy), we find

$$|\delta| = \left| \tan^{-1} \left(\frac{B_t}{B_n} \right) \right| \approx \left| \frac{B_t}{B_n} \right| \approx \frac{1.116 \times 10^{-11}}{5.350 \times 10^{-6}} = 2.086 \times 10^{-6}.$$

Combination of Asymmetries: Q-weak Kinematics

In Q-weak, B_n was measured as

$$B_n \approx -5.350 \text{ ppm.}$$

Using Q-weak kinematics, $Q^2 = 0.025 \text{ GeV}$, $E_1 = 1.125 \text{ GeV}$ (electron beam energy), we find

$$|\delta| = \left| \tan^{-1} \left(\frac{B_t}{B_n} \right) \right| \approx \left| \frac{B_t}{B_n} \right| \approx \frac{1.116 \times 10^{-11}}{5.350 \times 10^{-6}} = 2.086 \times 10^{-6}.$$

→ This is too small to affect the Q-weak measurements.

Order of Magnitude Estimates

- Beam Normal SSA:

$$B_n \sim \frac{\alpha m_e}{M} \sim 5 \times 10^{-6} \rightarrow 5 \text{ ppm}$$

Order of Magnitude Estimates

- Beam Normal SSA:

$$B_n \sim \frac{\alpha m_e}{M} \sim 5 \times 10^{-6} \rightarrow 5 \text{ ppm}$$

- A_{PV} (from longitudinal polarization):

$$A_{PV} \sim \frac{Q^2}{M_Z^2} \sim Q^2 \times 10^{-4}$$

Order of Magnitude Estimates

- Beam Normal SSA:

$$B_n \sim \frac{\alpha m_e}{M} \sim 5 \times 10^{-6} \rightarrow 5 \text{ ppm}$$

- A_{PV} (from longitudinal polarization):

$$A_{PV} \sim \frac{Q^2}{M_Z^2} \sim Q^2 \times 10^{-4}$$

- Beam Transverse SSA:

$$B_t \sim \frac{Q^2}{M_Z^2} \frac{m_e}{M} \sim Q^2 \times (5 \times 10^{-8})$$

Order of Magnitude Estimates

- Beam Normal SSA:

$$B_n \sim \frac{\alpha m_e}{M} \sim 5 \times 10^{-6} \rightarrow 5 \text{ ppm}$$

- A_{PV} (from longitudinal polarization):

$$A_{PV} \sim \frac{Q^2}{M_Z^2} \sim Q^2 \times 10^{-4}$$

- Beam Transverse SSA:

$$B_t \sim \frac{Q^2}{M_Z^2} \frac{m_e}{M} \sim Q^2 \times (5 \times 10^{-8})$$


- What factors contribute to B_n ?

Inelastic Hadronic Intermediate State

Scattering process:

$$e(p_1) + N(p_2) \rightarrow e(p_3) + N(p_4)$$

with intermediate leptonic momentum k and intermediate hadronic momentum $W = p_2 + q_1$.



One- and Two- Photon Interference

- We consider the contribution to the normal BSSA from $\mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma}$.

One- and Two- Photon Interference

- We consider the contribution to the normal BSSA from $\mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma}$.
- For polarized beam, spinor relation becomes

$$\bar{u}_e(k) u_e(k) = (1 + \gamma_5 \not{\epsilon})(\not{k} + m_e).$$

One- and Two- Photon Interference

- We consider the contribution to the normal BSSA from $\mathcal{M}_\gamma^* \mathcal{M}_{\gamma\gamma}$.
- For polarized beam, spinor relation becomes

$$\bar{u}_e(k) u_e(k) = (1 + \gamma_5 \not{\epsilon})(\not{k} + m_e).$$

- The BSSA comes from the absorptive part of the two-photon exchange amplitude (Pasquini & Vanderhaeghen PRC 70, 2004):

$$B_n = \frac{2 \operatorname{Im} \left(\sum_{\text{spins}} \mathcal{M}_\gamma^* \cdot \text{Abs } \mathcal{M}_{\gamma\gamma} \right)}{\sum_{\text{spins}} |\mathcal{M}_\gamma|^2}.$$

Leptonic and Hadronic Tensors

We can write this as

$$B_n = \frac{e^6}{(2\pi)^3 Q^2} \left(\sum_{\text{spins}} |\mathcal{M}_\gamma|^2 \right)^{-1} \int \frac{d^3 \vec{k}}{E_k} \frac{1}{Q_1^2 Q_2^2} \text{Im} \{ L_{\alpha\mu\nu} H^{\alpha\mu\nu} \},$$

Leptonic and Hadronic Tensors

We can write this as

$$B_n = \frac{e^6}{(2\pi)^3 Q^2} \left(\sum_{\text{spins}} |\mathcal{M}_\gamma|^2 \right)^{-1} \int \frac{d^3 \vec{k}}{E_k} \frac{1}{Q_1^2 Q_2^2} \text{Im} \{ L_{\alpha\mu\nu} H^{\alpha\mu\nu} \},$$

where

$$L_{\alpha\mu\nu} = \text{Tr} \left[\frac{1}{2} (1 + \gamma_5 \not{\epsilon}) (\not{p}_1 + m_e) \gamma_\alpha (\not{p}_1 - \not{q} + m_e) \gamma_\mu (\not{p}_1 - \not{q}_1 + m_e) \gamma_\nu \right].$$

Leptonic and Hadronic Tensors

We can write this as

$$B_n = \frac{e^6}{(2\pi)^3 Q^2} \left(\sum_{\text{spins}} |\mathcal{M}_\gamma|^2 \right)^{-1} \int \frac{d^3 \vec{k}}{E_k} \frac{1}{Q_1^2 Q_2^2} \text{Im} \{ L_{\alpha\mu\nu} H^{\alpha\mu\nu} \},$$

where

$$L_{\alpha\mu\nu} = \text{Tr} \left[\frac{1}{2} (1 + \gamma_5 \not{\epsilon}) (\not{p}_1 + m_e) \gamma_\alpha (\not{p}_1 - \not{q} + m_e) \gamma_\mu (\not{p}_1 - \not{q}_1 + m_e) \gamma_\nu \right].$$

and we take the forward limit on the hadronic tensor:

$$H^{\alpha\mu\nu} = 2p_2^\alpha W^{\mu\nu}$$

Leptonic and Hadronic Tensors

We can write this as

$$B_n = \frac{e^6}{(2\pi)^3 Q^2} \left(\sum_{\text{spins}} |\mathcal{M}_\gamma|^2 \right)^{-1} \int \frac{d^3 \vec{k}}{E_k} \frac{1}{Q_1^2 Q_2^2} \text{Im} \{ L_{\alpha\mu\nu} H^{\alpha\mu\nu} \},$$

where

$$L_{\alpha\mu\nu} = \text{Tr} \left[\frac{1}{2} (1 + \gamma_5 \not{\epsilon}) (\not{p}_1 + m_e) \gamma_\alpha (\not{p}_1 - \not{q} + m_e) \gamma_\mu (\not{p}_1 - \not{q}_1 + m_e) \gamma_\nu \right].$$

and we take the forward limit on the hadronic tensor:

$$H^{\alpha\mu\nu} = 2p_2^\alpha W^{\mu\nu}$$

where (de Rujula et al. Nucl. Phys. B35, 1971)

$$W^{\mu\nu} = \left(-g^{\mu\nu} + \frac{q_1^\mu q_1^\nu}{q_1^2} \right) W_1 + \frac{1}{M^2} \left(p_2^\mu - \frac{p_2 \cdot q_1}{q_1^2} q_1^\mu \right) \left(p_2^\nu - \frac{p_2 \cdot q_1}{q_1^2} q_1^\nu \right) W_2.$$

Finally, we can write the asymmetry as

Finally, we can write the asymmetry as

$$B_n = -\frac{1}{(2\pi)^3} \frac{e^2 Q^2}{D(s, Q^2)} \frac{1}{8E_1 E_3 M} \int_0^{2\pi} d\phi_k \int_{M^2}^s dW^2 \int_0^{Q_{1,\max}^2} dQ_1^2$$

$$\times \frac{\text{Im} \{L_{\alpha\mu\nu} H^{\alpha\mu\nu}\}}{Q_1^2 |\vec{k}| (1 - \cos \theta \cos \theta_k - \sin \theta \sin \theta_k \cos \phi_k)}$$

Finally, we can write the asymmetry as

$$B_n = -\frac{1}{(2\pi)^3} \frac{e^2 Q^2}{D(s, Q^2)} \frac{1}{8E_1 E_3 M} \int_0^{2\pi} d\phi_k \int_{M^2}^s dW^2 \int_0^{Q_{1,\max}^2} dQ_1^2 \\ \times \frac{\text{Im} \{L_{\alpha\mu\nu} H^{\alpha\mu\nu}\}}{Q_1^2 |\vec{k}| (1 - \cos \theta \cos \theta_k - \sin \theta \sin \theta_k \cos \phi_k)}$$

where E_1 and E_3 are the energies of the incoming and outgoing leptons, θ_k is the angle between \vec{p}_1 and \vec{k} , ϕ_k is the azimuthal angle, and

$$D(s, Q^2) = \frac{8Q^4}{1 - \epsilon} \left\{ G_M^2 + \frac{\epsilon}{\tau} G_E^2 \right\},$$

Once again,

$$B_n = -\frac{1}{(2\pi)^3} \frac{e^2 Q^2}{D(s, Q^2)} \frac{1}{8E_1 E_3 M} \int_0^{2\pi} d\phi_k \int_{M^2}^s dW^2 \int_0^{Q_{1,\max}^2} dQ_1^2$$

$$\times \frac{\text{Im} \{L_{\alpha\mu\nu} H^{\alpha\mu\nu}\}}{Q_1^2 |\vec{k}| (1 - \cos \theta \cos \theta_k - \sin \theta \sin \theta_k \cos \phi_k)}$$

Once again,

$$B_n = -\frac{1}{(2\pi)^3} \frac{e^2 Q^2}{D(s, Q^2)} \frac{1}{8E_1 E_3 M} \int_0^{2\pi} d\phi_k \int_{M^2}^s dW^2 \int_0^{Q_{1,\max}^2} dQ_1^2 \\ \times \frac{\text{Im} \{L_{\alpha\mu\nu} H^{\alpha\mu\nu}\}}{Q_1^2 |\vec{k}| (1 - \cos \theta \cos \theta_k - \sin \theta \sin \theta_k \cos \phi_k)}$$

The tensor contraction is

$$\text{Im} L_{\alpha\mu\nu} H^{\alpha\mu\nu} = \frac{8m_e}{M^2 Q_1^2} \left[2W_1 M^2 Q_1^2 (\epsilon^{p_1 p_2 q s} + \epsilon^{p_2 q q_1 s}) \right. \\ \left. + W_2 \epsilon^{p_2 q q_1 s} \left((M^2 - Q_1^2 - W^2)(p_1 \cdot p_2) - M^2 Q_1^2 \right) \right].$$

Once again,

$$B_n = -\frac{1}{(2\pi)^3} \frac{e^2 Q^2}{D(s, Q^2)} \frac{1}{8E_1 E_3 M} \int_0^{2\pi} d\phi_k \int_{M^2}^s dW^2 \int_0^{Q_{1,\max}^2} dQ_1^2$$

$$\times \frac{\text{Im} \{L_{\alpha\mu\nu} H^{\alpha\mu\nu}\}}{Q_1^2 |\vec{k}| (1 - \cos \theta \cos \theta_k - \sin \theta \sin \theta_k \cos \phi_k)}$$

The tensor contraction is

$$\text{Im} L_{\alpha\mu\nu} H^{\alpha\mu\nu} = \frac{8m_e}{M^2 Q_1^2} \left[2W_1 M^2 Q_1^2 (\epsilon^{p_1 p_2 q s} + \epsilon^{p_2 q q_1 s}) \right.$$

$$\left. + W_2 \epsilon^{p_2 q q_1 s} \left((M^2 - Q_1^2 - W^2)(p_1 \cdot p_2) - M^2 Q_1^2 \right) \right].$$

→ Currently evaluating for Q-weak kinematics.

Conclusion

- 1 Beam single spin asymmetries result from the difference in cross sections produced by the beam polarization in electron-proton scattering experiments.

Conclusion

- 1 Beam single spin asymmetries result from the difference in cross sections produced by the beam polarization in electron-proton scattering experiments.
- 2 The interference between one- and two-photon exchange amplitudes produces a beam normal single spin asymmetry.

Conclusion

- ① Beam single spin asymmetries result from the difference in cross sections produced by the beam polarization in electron-proton scattering experiments.
- ② The interference between one- and two-photon exchange amplitudes produces a beam normal single spin asymmetry.
- ③ The interference between one-photon and Z exchange amplitudes produces a beam transverse single spin asymmetry.

Conclusion

- 1 Beam single spin asymmetries result from the difference in cross sections produced by the beam polarization in electron-proton scattering experiments.
- 2 The interference between one- and two-photon exchange amplitudes produces a beam normal single spin asymmetry.
- 3 The interference between one-photon and Z exchange amplitudes produces a beam transverse single spin asymmetry.
- 4 The combination of these two asymmetries produces a small phase shift that may be detectable in the future.

Conclusion

- 1 Beam single spin asymmetries result from the difference in cross sections produced by the beam polarization in electron-proton scattering experiments.
- 2 The interference between one- and two-photon exchange amplitudes produces a beam normal single spin asymmetry.
- 3 The interference between one-photon and Z exchange amplitudes produces a beam transverse single spin asymmetry.
- 4 The combination of these two asymmetries produces a small phase shift that may be detectable in the future.
- 5 The inelastic hadronic intermediate state produces an important contribution to the BSSA (soon to be quantified).

Conclusion

- 1 Beam single spin asymmetries result from the difference in cross sections produced by the beam polarization in electron-proton scattering experiments.
- 2 The interference between one- and two-photon exchange amplitudes produces a beam normal single spin asymmetry.
- 3 The interference between one-photon and Z exchange amplitudes produces a beam transverse single spin asymmetry.
- 4 The combination of these two asymmetries produces a small phase shift that may be detectable in the future.
- 5 The inelastic hadronic intermediate state produces an important contribution to the BSSA (soon to be quantified).
- 6 The framework developed here could potentially be used to compute other beam or target SSA at near-forward angles.

References

- ① Waidyawansa, B. P. (2013). *A 3% Measurement of the Beam Normal Single Spin Asymmetry in Forward Angle Elastic Electron-Proton Scattering using the Q_{weak} Setup*. Ohio University: U.S.A.
- ② B. Pasquini and M. Vanderhaeghen, Physical Review C **70**, 045206 (2004).
- ③ M. Gorchtein et al., Nucl. Phys. **A741**, 234 (2004).
- ④ M.L. Goldberger, Y. Nambu, R. Oehme, Ann. Phys. **2** (1957) 226.
- ⑤ A. De Rujula, J. M. Kaplan, and E. de Rafael, Nucl. Phys. **B35**, 365 (1971).

Acknowledgements

I would like to thank my mentor, Dr. Wally Melnitchouk, for his guidance, instruction, and patience. I would also like to thank Dr. Peter Blunden for his aid in completing the calculations for this project.

Questions?