

Version xx as of November 11, 2019

Primary authors: Carlos Yero, Werner Boeglin, Mark Jones

To be submitted to PRL

Comment to cyero002@fiu.edu by xxx, yyy

DØ INTERNAL DOCUMENT – NOT FOR PUBLIC DISTRIBUTION

First Measurements of the $D(e,e'p)n$ Cross Section at Very High Recoil Momenta and Large Q^2

C. Yero and W.U. Boeglin

Florida International University, University Park, Florida 33199, USA

M.K. Jones

Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

(for the Hall C Collaboration)

(Dated: November 11, 2019)

First results of cross section measurements of the $^2H(e,e'p)n$ reaction at 4-momentum transfers $4 \leq Q^2 \leq 5$ (GeV/c)² and neutron recoil momenta up to 1.18 GeV/c are presented. At the selected kinematics, Meson Exchange Currents (MEC) and Isobar Configurations (IC) are suppressed. Final State Interactions (FSI) have also been suppressed by choosing a kinematic region where the neutron recoil angle (θ_{nq}) is between 35 and 45 degrees with respect to the 3-momentum transfer, \vec{q} . In this region, the Plane Wave Impulse Approximation (PWIA) dominates and comparison to recent theoretical calculations show data to be sensitive to momentum distributions up to ~ 700 MeV/c recoil momenta.

Being the most simple np bound state, the deuteron serves as a starting point to study the strong nuclear force at the subfermi level which is currently not well understood. At such small internucleon distances the NN (nucleon-nucleon) potential is expected to exhibit a repulsive core in which the interacting nucleon pair begins to overlap. The overlap is directly related to two-nucleon short range correlations (SRC) observed in $A > 2$ nuclei [1–4]. Short-distance studies of the deuteron are also important in determining whether or to what extent the description of nuclei in terms of nucleon/meson degrees of freedom must be supplemented by the inclusion of explicit quark effects, which is an issue of fundamental importance in nuclear physics[5]. To date, there are only a few nuclear experiments for which a transition between nucleonic to quark degrees of freedom can be observed [6–8]. This Letter presents first results of $^2H(e,e'p)n$ in which kinematics were taken to the limit where a transition to non-nucleonic degrees of freedom is expected.

The most direct way to study the short range structure of the deuteron wavefunction (or equivalently, its high momentum components) is via the exclusive deuteron electro-disintegration reaction at very high neutron recoil (or missing) momenta and within the PWIA kinematics. In this approximation, the virtual photon couples to the proton which is ejected from the nucleus without further interaction with the recoiling neutron, which carries a momentum equal in magnitude but opposite in direction to the initial state proton, $\vec{p}_r = -\vec{p}_{i,p}$. This gives direct access to the deuteron momentum distributions since the scattered neutron momentum remains unchanged.

In reality, the ejected particles undergo subsequent interactions resulting in re-scattering of the proton and neutron (FSIs). Another possibility is that the photon may couple to the virtual meson being exchanged between the nucleons (MECs), or the photon may excite either nucleon in the deuteron into a resonance state (ICs) which decays back into the ground state nucleon causing further re-scattering between the proton and neutron. The above-mentioned long-range processes alter the final neutron momentum making the deuteron momentum distributions difficult to access.

Previous deuteron electro-disintegration experiments performed at Jefferson Lab (JLab) have helped disentangle and quantify the contributions from FSI, MEC and IC on the $^2H(e,e'p)n$ cross-section and determine the kinematics at which they are either suppressed (MECs and ICs) or under control (FSIs). The first of these was performed in Hall A [9] at a relatively low momentum transfer of $Q^2 = 0.665$ (GeV/c)² and neutron recoil momenta up to $p_r = 550$ MeV/c where it was shown that for $p_r > 300$ MeV/c, the inclusion of FSI, MEC and IC was necessary in Arenhovel's calculations for a satisfactory agreement between the theory and data.

The next experiment was performed in Hall B [10] using the CEBAF Large Acceptance Spectrometer (CLAS) which took advantage of its large detector acceptance to simultaneously measure a wide variety of kinematic settings giving an overview of the $^2H(e,e'p)n$ reaction kinematics. This was the first experiment to probe the deuteron at high momentum transfers ($1.75 \leq Q^2 \leq 5.5$ (GeV/c)²) and presented angular distributions of

cross-sections that confirmed the onset of the General Eikonal Approximation (GEA), predicting a strong angular dependence of FSI with neutron recoil angles with FSI peaking at $\theta_{nq} \sim 70^\circ$. The cross-sections versus neutron recoil momenta up to 2 GeV/c were also presented with integrated neutron recoil angles in the range $20^\circ < \theta_{nq} < 160^\circ$ to gain better statistical precision. As a result, it was not possible to choose kinematical regions binned in θ_{nq} in which FSI were minimal to extract the momentum distributions.

Finally, a third ${}^2H(e, e'p)n$ experiment was performed in Hall A [11] at $Q^2 = 0.8, 2.5, 3.5$ (GeV/c) 2 and recoil momenta up to 550 MeV/c at kinematics which allowed the extraction of angular and momentum distributions for significantly smaller kinematical bins than in Hall B/CLAS. The angular distributions were presented as the cross-section ratio, $R = \sigma_{exp}/\sigma_{PWIA}$ versus θ_{nq} , and verified the strong anisotropy of FSI with recoil angle previously observed in Hall B. Most importantly, for recoil neutron momentum bins, $p_r = 0.4 \pm 0.02$ and 0.5 ± 0.02 GeV/c, the ratio $R \sim 1$ for $35^\circ \leq \theta_{nq} \leq 45^\circ$ indicating a reduced sensitivity of the experimental cross-section to FSI, in which, $\sigma_{exp} \sim \sigma_{PWIA}$. This kinematic window in which FSI are small can also be seen in the momentum distributions for $\theta_{nq} = 35 \pm 5^\circ$ and $45 \pm 5^\circ$, where data and theory agree well within the PWIA kinematics. The experiment concluded that the kinematic window found at $35^\circ \leq \theta_{nq} \leq 45^\circ$ gives for the first time a direct access to the high momentum components of the deuteron momentum distribution.

The experiment presented on this Letter takes advantage of the kinematic window found previously in Hall A and extends the ${}^2H(e, e'p)n$ cross section measurements to $Q^2 = 4.5$ (GeV/c) 2 and neutron recoil momenta up to 1.18 GeV/c. In this configuration, MECs and ICs are suppressed and FSIs are under control for neutron recoil angles between 35 and 45 degrees giving access to unprecedented high momentum components of the deuteron wavefunction.

This experiment was part of a group of four experiments that commissioned the new Hall C Super High Momentum Spectrometer (SHMS) as part of the 12 GeV upgrade at JLab. An electron beam was incident on a 10 cm long liquid deuterium target (LD2). The scattered electron and knocked-out proton were detected in coincidence by the SHMS and High Momentum Spectrometer (HMS), respectively. The “missing” (undetected) neutron was reconstructed from momentum conservation laws. The beam currents delivered by the accelerator ranged between 40-60 μ A due to frequent beam trips at higher currents and the beam was rastered over a 2x2 mm 2 area to reduce the effects of localized boiling on the cryogenic targets (hydrogen and deuterium).

Both spectrometers at Hall C have similar standard detector packages, each with 1) four sets of hodoscope planes (scintillator arrays) used for triggering, 2) a pair

of drift chambers used for tracking, 3) a calorimeter used for e^-/π^- discrimination and 4) a gas Čerenkov used also for e^-/π^- and an additional Noble Gas Čerenkov used in the SHMS for $e^-/\pi^+/K^+$ at momenta > 6 GeV/c. Due to the absence of significant background on this experiment and the low coincidence trigger rates ($\sim 1 - 3$ Hz) at the higher missing momentum settings, the use of additional particle identification (PID) was found to have little to no effect on the final cross section.

We measured three missing momentum settings: $p_r = 80, 580$ and 750 MeV/c. In the high missing momentum settings the spectrometer configuration was changed multiple times resulting in either the spectrometer angle or momentum not being exactly the same. As a result, two separate data sets measured for the 580 MeV/c and three data sets for the 750 MeV/c setting. The spectrometer central settings are approximately as follows: the SHMS central angle and momentum settings were kept fixed at (12.194 deg, 8.5342 GeV/c) and the HMS central angle and momentum settings were changed from (38.896 deg, 2.840 GeV/c) at the 80 MeV setting to (54.992 deg, 2.1925 GeV/c) and (58.391 deg, 2.0915 GeV/c) at the 580 and 750 MeV/c settings, respectively. At these kinematics, the 3-momentum transfer is $|\vec{q}| = 2.86$ GeV/c and is on the order of the final proton momentum indicating that most of the energy and momentum are transferred to the proton. As a result, the ejected proton scatters at angles $\theta_{pq} \sim 0$, relative to the \vec{q} . This configuration is known as the “parallel-kinematics” and suppresses the process in which the neutron is struck and the proton is a spectator.

In addition to deuteron kinematics, ${}^1H(e, e'p)$ elastic data was also taken at kinematics close to the deuteron 80 MeV setting for cross-checks with the spectrometer acceptance model as well as for normalization purposes using the Hall C Monte Carlo simulation program, SIMC. Additional ${}^1H(e, e'p)$ data was also taken at three other kinematic settings that covered the entire SHMS momentum acceptance range and were used for spectrometer optics studies and calibration.

The event selection criteria was done exactly the same for the hydrogen and deuterium data. The criteria was determined by making 1) standard cuts on the spectrometer momentum fraction (δ) to select a region in which the reconstruction optics is well known, 2) an HMS collimator cut to restrict the spectrometer solid angle acceptance to events that only passed through the collimator and not by re-scattering from the edges, 3) a missing energy cut (peak ~ 2.2 MeV for the deuteron) to select true ep coincidences and not events from the radiative tail, 4) a coincidence time cut to select true coincidence events and not accidentals, 5) a PID cut on the SHMS calorimeter to select electrons and not other sources of background, mostly pions and 6) a z-vertex difference cut between the HMS and SHMS z reaction vertex difference to select events that truly originated from the same reaction

vertex at the target.

The experimental data yield was normalized by the total charge and corrected for tracking efficiencies, total live time, proton absorption and target boiling factors. For $^1\text{H}(e, e'p)$, the corrected data yield was compared to SIMC using P. Bosted's proton form factor parametrization[12] to check the spectrometer acceptance model. The data to SIMC yield ratio integrated over invariant mass W was determined to be unity, so there was no need to include an overall hydrogen normalization factor. For the $^2\text{H}(e, e'p)n$ data, the measured cross sections were compared to the model cross sections (incorporated as a SIMC subroutine) from calculations by J.M. Laget using the Paris potential. Variations of up to $\sim 20\%$ for recoil momenta up to ~ 250 MeV/c were observed which are typical for this setting using the Paris potential. The 80 MeV data was also checked for reproducibility against the Hall A data. This agreement gives us confidence on the measurements made at higher missing momentum setting for which no data exists.

The systematic uncertainties on the measured cross sections were determined from 1) normalization and 2) kinematic uncertainties. The individual contributions from normalization uncertainties were determined to be: tracking efficiencies (0.40%-HMS, 0.59%-SHMS), target boiling (0.39%), total live time (3.0%) and total charge (2.0%) for an overall normalization uncertainty added in quadrature of 3.7%.

The kinematic uncertainties were determined point-to-point in (θ_{nq}, p_r) bins for each data set independently, and added in quadrature for overlapping p_r bins of different data sets. For $\theta_{nq} = 35, 45$ and 75 deg (presented on this Letter) the overall kinematic uncertainty varied up to 6.5% for $p_r \leq 1.01$ GeV/c. The overall systematic uncertainty in the cross section was determined by the quadrature sum of the normalization and kinematic uncertainties. This results was then added in quadrature to the statistical uncertainty to obtain the final uncertainty in the cross section.

The data was radiatively corrected for each bin in (θ_{nq}, p_r) by multiplying measured cross sections to the ratio of the SIMC yield without and with radiative effects. Bin-centering corrections were also applied by multiplying the radiative corrected cross sections to the ratio of theoretical cross sections (external to SIMC) to the average cross sections calculated from SIMC. The theoretical calculations used in the bin-centering corrections were done by J.M. Laget using the Paris potential.

Both experimental and theoretical reduced cross sections were extracted from the measured (or model) cross sections for each data of set independently and were averaged for overlapping bins in p_r , where the reduced cross sections are defined as follows:

$$\sigma_{red} \equiv \frac{\sigma_{exp(th)}}{K f_{rec} \sigma_{cc1}} \quad (1)$$




FIG. 1. The ratio $R(p_r) = \sigma_{red}/n(p_r)$ is shown in (a)-(c) for $\theta_{nq} = 35^\circ, 45^\circ$ and 75° , respectively, each with a bin width of $\pm 5^\circ$. The dashed reference (magenta) line refers to CD-Bonn momentum distribution ($n(p_r)$) by which the data and all models are divided.

where $\sigma_{exp(th)}$ is the 5-fold experimental (or theoretical) differential cross section, K is a kinematical factor, f_{rec} is a the recoil factor that arises from the integration over missing energy and σ_{cc1} is the de Forest[13] electron-proton offshell cross section calculated from P. Bosted's form factor parametrization [12]. Only within the PWIA, σ_{red} can be interpreted as the proton momentum distribution inside the deuteron where most of the kinematical dependencies that arise from the cross section have been factored out by σ_{cc1} , leaving only a dependency on p_r .

To quantify by how much and to what extent the data agrees with theory, the ratio of the experimental (or theoretical) reduced cross sections (σ_{red}) to the deuteron momentum distributions ($n(p_r)$) using the charge-dependent Bonn (CD-Bonn) potential is shown in Fig. 1. The theoretical calculations for the CD-Bonn and Argonne v_{18} (AV18) potentials were performed by M. Sargsian and those for the Paris potential were

done by J.M. Laget. For $p_r \lesssim 300$ MeV/c, the data is in good agreement with all models for the neutron recoil angles shown in Fig. 1. For $p_r \gtrsim 300$, at recoil angles $\theta_{nq} = 35^\circ \pm 5^\circ$ [Fig. 1(a)], the data is best described by the CD-Bonn curves which exhibit a reduced sensitivity to FSIs and an enhanced sensitivity to the momentum distribution with a ratio $R \sim 1$ for recoil momenta up to ~ 800 MeV/c before being overwhelmed by FSIs. The data, however, is sensitive to the momentum distributions only up to $p_r \sim 750$ MeV/c before transitioning to other theoretical curves with a maximal ratio of $R \sim 3$ at $p_r \sim 940$ MeV/c due to statistical limitations. For recoil angles $\theta_{nq} = 45^\circ \pm 5^\circ$ [Fig. 1(b)], both the data and CD-Bonn curves show sensitivities to momentum distributions up to $p_r \sim 580$ MeV/c. At $p_r > 580$ MeV/c, FSIs become increasingly important for CD-Bonn curves, whereas the data shows a similar behaviour as in Fig. 1(a) with an early rise in the ratio than predicted by the CD-Bonn FSI model. For $\theta_{nq} = 75^\circ \pm 5^\circ$ [Fig. 1(c)], the onset of GEA is observed beyond $p_r \sim 300$ MeV/c, with a strong angular dependence of the reduced cross sections on FSIs predicted by all models and verified by data which was statistically limited at larger recoil angles.

Figure 2 shows the extracted experimental and theoretical reduced cross sections as a function of neutron recoil momentum, p_r for three angular settings at $4 \leq Q^2 \leq 5$ (GeV/c)². The data is compared to the results from the previous Hall A experiment[11] at a $Q^2 = 3.5$ (GeV/c)². The overlay of the Hall A data (cyan) in Fig. 2 provides a continuity to the data from this experiment in the transition region from low (80

MeV/c) to high (580, 750) MeV/c missing momentum settings in which there was no data. There is also an overall good agreement between the two experiments in the regions in which there is an overlap in p_r .




FIG. 2. The reduced cross section $\sigma_{red}(p_r)$ as a function of neutron recoil momentum p_r is shown in (a)-(c) for recoil angles $\theta_{nq} = 35^\circ, 45^\circ$ and 75° , respectively, with a bin width of $\pm 5^\circ$. The data is compared to the previous Hall A experiment results as well as the Paris, AV18 and CD-Bonn theoretical reduced cross sections.

The authors would like to thank Tex, LaTeX and Friends for the answer to this question.

-
- [1] K. S. Egiyan *et al.* (CLAS Collaboration), Observation of nuclear scaling in the $A(e, e')$ reaction at $x_B > 1$, *Phys. Rev. C* **68**, 014313 (2003).
 - [2] K. S. Egiyan *et al.* (CLAS Collaboration), Measurement of two- and three-nucleon short-range correlation probabilities in nuclei, *Phys. Rev. Lett.* **96**, 082501 (2006).
 - [3] R. Shneor *et al.* (Jefferson Lab Hall A Collaboration), Investigation of proton-proton short-range correlations via the $^{12}\text{C}(e, e'pp)$ reaction, *Phys. Rev. Lett.* **99**, 072501 (2007).
 - [4] N. Fomin, D. Higinbotham, M. Sargsian, and P. Solvignon, New results on short-range correlations in nuclei, *Annual Review of Nuclear and Particle Science* **67**, 129159 (2017).
 - [5] P. Ulmer *et al.*, Short-Distance Structure of the Deuteron and Reaction Dynamics in $^2\text{H}(e, e'p)n$, https://www.jlab.org/exp_prog/proposals/01/PR01-020.pdf (2001), *Jefferson Lab Proposal E01-020*.
 - [6] C. Bochna *et al.*, Measurements of deuteron photodisintegration up to 4.0 gev, *Phys. Rev. Lett.* **81**, 4576 (1998).
 - [7] E. C. Schulte *et al.*, Measurement of the high energy two-body deuteron photodisintegration differential cross section, *Phys. Rev. Lett.* **87**, 102302 (2001).
 - [8] E. C. Schulte *et al.*, High energy angular distribution measurements of the exclusive deuteron photodisintegration reaction, *Phys. Rev. C* **66**, 042201 (2002).
 - [9] P. E. Ulmer *et al.*, $^2\text{H}(e, e'p)n$ reaction at high recoil momenta, *Phys. Rev. Lett.* **89**, 062301 (2002).
 - [10] K. S. Egiyan *et al.* (CLAS Collaboration), Experimental study of exclusive $^2\text{H}(e, e'p)n$ reaction mechanisms at high Q^2 , *Phys. Rev. Lett.* **98**, 262502 (2007).
 - [11] W. U. Boeglin *et al.* (For the Hall A Collaboration), Probing the high momentum component of the deuteron at high Q^2 , *Phys. Rev. Lett.* **107**, 262501 (2011).
 - [12] P. E. Bosted, Empirical fit to the nucleon electromagnetic form factors, *Phys. Rev. C* **51**, 409 (1995).
 - [13] T. D. Forest, Off-shell electron-nucleon cross sections: The impulse approximation, *Nuclear Physics A* **392**, 232 (1983).