

Version xx as of October 29, 2019

Primary authors: Carlos Yero, Werner Boeglin, Mark Jones

To be submitted to PRL

Comment to cyero002@fiu.edu by xxx, yyy

DØ INTERNAL DOCUMENT – NOT FOR PUBLIC DISTRIBUTION

First Measurements of the $D(e,e'p)n$ Cross Section at Very High Recoil Momenta and Large Q^2

C. Yero and W.U. Boeglin

Florida International University, University Park, Florida 33199, USA

M.K. Jones

Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

(for the Hall C Collaboration)

(Dated: October 29, 2019)

First results of cross section measurements of the $^2H(e,e'p)n$ reaction at 4-momentum transfers $4 \leq Q^2 \leq 5$ (GeV/c)² and neutron recoil momenta up to 1.18 GeV/c are presented. At the selected kinematics, Meson Exchange Currents (MEC) and Isobar Configurations (IC) are suppressed. Final State Interactions (FSI) have also been suppressed by choosing a kinematic region where the neutron recoil angle (θ_{nq}) is between 35 and 45 degrees with respect to the 3-momentum transfer, \vec{q} . In this region, the Plane Wave Impulse Approximation (PWIA) dominates and comparison to recent theoretical calculations show data to be sensitive to momentum distributions up to ~ 700 MeV/c recoil momenta.

Being the most simple np bound state, the deuteron serves as a starting point to study the strong nuclear force at the subfermi level which is currently not well understood. At such small internucleon distances the NN (nucleon-nucleon) potential is expected to exhibit a repulsive core in which the interacting nucleon pair begins to overlap. The overlap is directly related to two-nucleon short range correlations (SRC) observed in $A > 2$ nuclei [1–4]. Short-distance studies of the deuteron are also important in determining whether or to what extent the description of nuclei in terms of nucleon/meson degrees of freedom must be supplemented by the inclusion of explicit quark effects, which is an issue of fundamental importance in nuclear physics[5]. To date, there are only a few nuclear experiments for which a transition between nucleonic to quark degrees of freedom can be observed [6–8]. This Letter presents first results of $^2H(e,e'p)n$ in which kinematics were taken to the limit where a transition to non-nucleonic degrees of freedom is expected.

The most direct way to study the short range structure of the deuteron wavefunction (or equivalently, its high momentum components) is via the exclusive deuteron electro-disintegration reaction at very high neutron recoil (or missing) momenta and within the PWIA kinematics. In this approximation, the virtual photon couples to the proton which is ejected from the nucleus without further interaction with the recoiling neutron, which carries a momentum equal in magnitude but opposite in direction to the initial state proton, $\vec{p}_r = -\vec{p}_{i,p}$. This gives direct access to the deuteron momentum distributions since

the scattered neutron momentum remains unchanged.

In reality, the ejected particles undergo subsequent interactions resulting in re-scattering of the proton and neutron (FSIs). Another possibility is that the photon may couple to the virtual meson being exchanged between the nucleons (MECs), or the photon may excite either nucleon in the deuteron into a resonance state (ICs) which decays back into the ground state nucleon causing further re-scattering between the proton and neutron. The above-mentioned long-range processes alter the final neutron momentum making the deuteron momentum distributions difficult to access.

Previous deuteron electro-disintegration experiments performed at Jefferson Lab (JLab) have helped disentangle and quantify the contributions from FSI, MEC and IC on the $^2H(e,e'p)n$ cross-section and determine the kinematics at which they are either suppressed (MECs and ICs) or under control (FSIs). The first of these was performed in Hall A [9] at a relatively low momentum transfer of $Q^2 = 0.665$ (GeV/c)² and neutron recoil momenta up to $p_r = 550$ MeV/c where it was shown that for $p_r > 300$ MeV/c, the inclusion of FSI, MEC and IC was necessary in Arenhovel's calculations for a satisfactory agreement between the theory and data.

The next experiment was performed in Hall B [10] using the CEBAF Large Acceptance Spectrometer (CLAS) which took advantage of its large detector acceptance to simultaneously measure a wide variety of kinematic settings giving an overview of the $^2H(e,e'p)n$ reaction kinematics. This was the first experiment to probe the

deuteron at high momentum transfers ($1.75 \leq Q^2 \leq 5.5$ (GeV/c)²) and presented angular distributions of cross-sections that confirmed the onset of the General Eikonal Approximation (GEA), predicting a strong angular dependence of FSI with neutron recoil angles with FSI peaking at $\theta_{nq} \sim 70^\circ$. The cross-sections versus neutron recoil momenta up to 2 GeV/c were also presented with integrated neutron recoil angles in the range $20^\circ < \theta_{nq} < 160^\circ$ to gain better statistical precision. As a result, it was not possible to choose kinematical regions binned in θ_{nq} in which FSI were minimal to extract the momentum distributions.

Finally, a third $^2\text{H}(e, e'p)n$ experiment was performed in Hall A [11] at $Q^2 = 0.8, 2.5, 3.5$ (GeV/c)² and recoil momenta up to 550 MeV/c at kinematics which allowed the extraction of angular and momentum distributions for significantly smaller kinematical bins than in Hall B/CLAS. The angular distributions were presented as the cross-section ratio, $R = \sigma_{exp}/\sigma_{PWIA}$ versus θ_{nq} , and verified the strong anisotropy of FSI with recoil angle previously observed in Hall B. Most importantly, for recoil neutron momentum bins, $p_r = 0.4 \pm 0.02$ and 0.5 ± 0.02 GeV/c, the ratio $R \sim 1$ for $35^\circ \leq \theta_{nq} \leq 45^\circ$ indicating a reduced sensitivity of the experimental cross-section to FSI, in which, $\sigma_{exp} \sim \sigma_{PWIA}$. This kinematic window in which FSI are small can also be seen in the momentum distributions for $\theta_{nq} = 35 \pm 5^\circ$ and $45 \pm 5^\circ$, where data and theory agree well within the PWIA kinematics. The experiment concluded that the kinematic window found at $35^\circ \leq \theta_{nq} \leq 45^\circ$ gives for the first time a direct access to the high momentum components of the deuteron momentum distribution.

This experiment takes advantage of the kinematic window found previously in Hall A and extends the $^2\text{H}(e, e'p)n$ cross section measurements to $Q^2 = 4.5$ (GeV/c)² and neutron recoil momenta up to 1.18 GeV/c. In this configuration, MECs and ICs are suppressed and FSIs are under control for recoil angles between 35 and 45 degrees giving access to unprecedented high momentum components of the deuteron wavefunction.

This experiment was part of a group of four experiments that commissioned the new Hall C Super High Momentum Spectrometer (SHMS) as part of the 12 GeV upgrade at JLab. An electron beam with beam energy of 10.6005 GeV was incident on a 10 cm long liquid deuterium target (LD2). The scattered electron and knocked-out proton were detected in coincidence by the SHMS and High Momentum Spectrometer (HMS), respectively. The “missing” (undetected) neutron was reconstructed from momentum conservation laws. The beam currents delivered by the accelerator ranged between 40-60 μA due to frequent beam trips at higher currents and the beam was rastered over a 2×2 mm² area to reduce the effects of localized boiling on the cryogenic targets (hydrogen and deuterium).

Both spectrometers at Hall C have similar standard

detector packages, each with 1) four sets of hodoscope planes (scintillator arrays) used for particle triggering, 2) a pair of drift chambers used for tracking, 3) a calorimeter used for e^-/π^- discrimination and 4) a gas Čerenkov used for additional e^-/π^- and π^+/K^+ (only Heavy Gas Čerenkov in SHMS) discrimination. Given the low trigger rates and absence of significant background on this experiment, the use of additional particle identification (PID) cuts was found to have little to no effect on the final cross section.

The authors would like to thank Tex, LaTeX and Friends for the answer to this question.

-
- [1] K. S. Egiyan *et al.* (CLAS Collaboration), Observation of nuclear scaling in the $A(e, e')$ reaction at $x_B > 1$, *Phys. Rev. C* **68**, 014313 (2003).
 - [2] K. S. Egiyan *et al.* (CLAS Collaboration), Measurement of two- and three-nucleon short-range correlation probabilities in nuclei, *Phys. Rev. Lett.* **96**, 082501 (2006).
 - [3] R. Shneor *et al.* (Jefferson Lab Hall A Collaboration), Investigation of proton-proton short-range correlations via the $^{12}\text{C}(e, e'pp)$ reaction, *Phys. Rev. Lett.* **99**, 072501 (2007).
 - [4] N. Fomin, D. Higinbotham, M. Sargsian, and P. Solvignon, New results on short-range correlations in nuclei, *Annual Review of Nuclear and Particle Science* **67**, 129159 (2017).
 - [5] P. Ulmer *et al.*, Short-Distance Structure of the Deuteron and Reaction Dynamics in $^2\text{H}(e, e'p)n$, https://www.jlab.org/exp_prog/proposals/01/PR01-020.pdf (2001), *Jefferson Lab Proposal E01-020*.
 - [6] C. Bochna *et al.*, Measurements of deuteron photodisintegration up to 4.0 gev, *Phys. Rev. Lett.* **81**, 4576 (1998).
 - [7] E. C. Schulte *et al.*, Measurement of the high energy two-body deuteron photodisintegration differential cross section, *Phys. Rev. Lett.* **87**, 102302 (2001).
 - [8] E. C. Schulte *et al.*, High energy angular distribution measurements of the exclusive deuteron photodisintegration reaction, *Phys. Rev. C* **66**, 042201 (2002).
 - [9] P. E. Ulmer *et al.*, $^2\text{H}(e, e'p)n$ reaction at high recoil momenta, *Phys. Rev. Lett.* **89**, 062301 (2002).

- [10] K. S. Egiyan *et al.* (CLAS Collaboration), Experimental study of exclusive ${}^2\text{H}(e, e'p)n$ reaction mechanisms at high Q^2 , *Phys. Rev. Lett.* **98**, 262502 (2007).
- [11] W. U. Boeglin *et al.* (For the Hall A Collaboration), Probing the high momentum component of the deuteron at high Q^2 , *Phys. Rev. Lett.* **107**, 262501 (2011).