AdCflox (Vet) Bolus

Ciprofloxacin Hydrochloride BP

For Veterinary Use Only

Composition:

Each Bolus contains Ciprofloxacin Hydrochloride BP 1164.4 mg (as Ciprofloxacin BP 1000 mg).

Indications:

AdCflox (Vet) Bolus is effective against several diseases as Respiratory Tract infections, Gastro Intestinal infections, Enterotoxaemia, Skin Bone & Ear infections, Swine dysentery, Uterine infections, Udder infections, Urinary Tract infection.

Dosage& Administration:

Cattle: 1-2 Bolus/ 70-150 kg body weight or 6.7-13.4 mg/kg body weight twice daily for 3-5 days.

Or, as directed by registered veterinary physician.

Adverse effects:

Central nervous system (CNS) toxicity; gastrointestinal reactions; hematuria; hepatotoxicity; hypersensitivity reactions; Stevens-Johnson Syndrome (blistering, itching, loosening, peeling, or redness of skin; diarrhea).

Drug Interaction:

Ciprofloxacin administered together with Theophylline can lead to elevated blood levels of Theophylline.

Precaution:

Patients receiving Ciprofloxacin should be well hydrated and should be instructed to drink fluids liberally. Because of the risk of crystalluria, it is recommended that the usual dosage of the drug should not be exceeded. Patients with a history of hypersensitivity to Ciprofloxacin or to other quinolones.

Use in Pregnancy and Lactation:

Intravenous doses of ciprofloxacin of up to 20 mg per kg of body weight (mg/kg) in pregnant rats and mice have not shown evidence of maternal toxicity, embryo toxicity, or teratogenic effects. Ciprofloxacin can be distributed into milk, sometimes at a higher concentration than in plasma, but it is not known under what conditions significant amounts might be absorbed by nursing animals.

Withdrawal Period

Meat: 14 days, Milk: 3 days.

Storage:

Store in a cool and dry place (below 30^o C temperatures) & protect from direct sunlight. Keep all medicines out of reach of children.

Supply:

10 X 2's bolus in blister pack.

