

Cone Theorem Readings Notes

Spring 2024

Note 4 — 04, 03, 2025 (draft version 0)

Yi Li

In this note, we will give a proof of Cone theorem for Kähler MMP. The major reference of this note are [HP24], [HP16] and [DH24].

Let us first state the theorem that we want to prove.

Theorem 0.1. Let $(X, B + \beta)$ be a generalized KLT pair. Assume the BDPP conjecture holds, then $K_X + B + \beta$ is metrically nef iff it's numerical nef.

Theorem 0.2. Let $(X,)$

Contents

1	Höring-Peternell's approach to the cone theorem for Kähler 3-folds	1
2	Das-Hacon's approach to cone theorem for DLT Kähler 3-folds	1
3	Kawamata's subadjunction for Kähler variety	1
4	Hacon-Păun's approach to cone theorem in any dimension	1
1	Höring-Peternell's approach to the cone theorem for Kähler 3-folds	
2	Das-Hacon's approach to cone theorem for DLT Kähler 3-folds	
3	Kawamata's subadjunction for Kähler variety	

The major technical tool used in the

4 Hacon-Păun's approach to cone theorem in any dimension

References

- [DH24] O. Das and C.D. Hacon. *On the Minimal Model Program for Kähler 3-folds*. 2024. arXiv: [2306.11708](https://arxiv.org/abs/2306.11708) [math.AG].

- [HP24] C.D. Hacon and M. Păun. *On the Canonical Bundle Formula and Adjunction for Generalized Kaehler Pairs*. 2024. arXiv: [2404.12007](#) [[math.AG](#)].
- [HP16] A. Höring and T. Peternell. “Minimal models for Kähler threefolds”. In: *Invent. Math.* 203.1 (2016), pp. 217–264.