


# Interrogation of human hematopoietic traits at single-cell and single-variant resolution


(Caleb, Erik, Jacob), Will?, Hilary, Joel, (Martin, Jason, Vijay)

Supplemental Information


## Overview of UK Biobank data




Supplemental Figure 1: Phenotypic and genetic correlations across the 16 traits examined.




Supplemental Figure 2: Numbers of pleiotropic variants




Supplemental Figure 3: Heritability estimates from LD Score Regression across 16 hematopoietic traits. The estimates of the narrow-sense SNP heritabilities are plotted in gray with their corresponding standard errors. Heritability estimates for all variants with fine-mapped posterior probability  $\geq 0.001$  are plotted in blue for each trait, and the proportions of total narrow-sense heritability captured by these fine-mapped variants (blue bar / gray bar) are indicated by the numbered labels.




Supplemental Figure 4: Haplotypes frequencies of the CCND3 variants rs2408955 and rs112233623 in the UKBB Population. Haplotype labels are formatted as rs2408955/rs112233623.



Supplemental Figure 5: Moderating effects of CCND3 haplotypes on red blood cell count.




Supplemental Figure 6: Local z-scores for enrichment of hematopoietic nucleosome-depleted regions in the set of fine-mapped variants with posterior probability  $> 0.10$ .




Supplemental Figure 7: Local annotation enrichments for fine-mapped variants, excluding all fine-map variants with  $R^2 > 0.80$  to the sentinel variant of any region.

(A)




Supplemental Figure 8: Side-by-side heatmaps showing overlap of hematopoietic nucleosome-depleted regions by cell type with fine-mapped variants ( $PP > 0.50$ ) by trait. The two heatmaps share a common y-axis of specific variants.







Supplemental Figure 9: Hematopoietic cell type enrichments for Mean Retic Volume and Monocyte count using various methods.




Supplemental Figure 10: All enrichments from g-chromVAR. The horizontal line shows a Bonferroni multiple testing adjusted threshold for statistical significance of enrichment.




Supplemental Figure 11: All enrichments from chromVAR. The horizontal line shows a Bonferonni multiple testing adjusted threshold for statistical significance of enrichment.



Supplemental Figure 12: All enrichments from LD Score. The horizontal line shows a Bonferonni multiple testing adjusted threshold for statistical significance of enrichment.




Supplemental Figure 13: Cell type - trait enrichments for g-chromVAR across different finemap variants posterior probability cutoffs. The horizontal line shows a Bonferonni multiple testing adjusted threshold for statistical significance of enrichment.



Supplemental Figure 14: Two subpopulations of CMP and MEP cells were obtained by k-means clustering on ATAC principal components or g-ChromVAR enrichments, respectively. Rank-order plots showing transcription factor binding sites ranked by difference in chromVAR enrichment between the two clusters of (A) CMP and (B) MEP populations

(B)



Supplemental Figure 15: Sample embedding of figure in document.

(C)