

Multivariable Calculus

MATH 1920 Lecture Notes

Author: Yichen

Course: MATH 1920

Date: September 21, 2026

Contents

1	Vectors in Two and Three Dimensions	1
1.1	Coordinates	1
1.2	Vectors and Position Vectors	1
1.3	Unit Vectors	3
1.4	Vector Operations	3
1.5	Parallel Vectors	4
2	Discussion Week 1: Distance Formulae and Geometric Shapes	5
2.1	Equations of Geometric Shapes	6
2.2	Linear Equations across Dimensions	6
3	Vectors and Directions	8
3.1	Dot Product	10
3.2	Determinants and Cross Products	14
3.2.1	Properties of the Cross Product	17
3.2.2	Algebraic Rules of the Cross Product	17
3.3	Scalar Triple Product	18
4	Discussion Week 2: Quadric Surfaces and Traces	19
5	Planes and Surfaces	21
5.1	Determining a Plane	21
5.2	Mathematical Formulations of a Plane	21
5.3	Properties of Normal Vectors	22
6	Spheres and Planes	24
6.1	Determining a Sphere	24
6.2	Plane-Sphere Intersections	25
6.3	Traces of Surfaces	25
6.4	Quadric Surfaces	26

7	Vector-Valued Functions	28
7.1	Coordinate Systems in 3D Space	28
7.1.1	Polar Coordinates (2D)	28
7.1.2	Cylindrical Coordinates	29
7.1.3	Spherical Coordinates	29
7.2	Coordinate System Transformations	30
7.3	Vector-Valued Functions	31
7.4	Gallery of Parametric Space Curves	34
7.5	Parameterization of Curves and Surfaces	34
7.6	Calculus of Vector-Valued Functions	35
7.7	Properties of Vector Derivatives & Integration	37
7.8	Arc Length of Curves	39
8	Scalar-Valued Functions of Several Variables	41
8.1	Visualizing Multivariable Functions: Traces and Level Sets	41
8.2	Level Sets and Contour Maps	41
8.3	Visualizing Higher Dimensions via Slices	42
9	Limits and Continuity of Functions of Several Variables	44

1 Vectors in Two and Three Dimensions

Question: What information specifies a line?


- 2 points
- A point and a direction vector
- Slope and y -intercept
- Intersection of 2 non-parallel planes

Comparison:


- **Single-variable calculus:** $f(x) \rightarrow$ rate of change of x
- **Multivariable calculus:** $f(x, y) \rightarrow$ functions of multiple variables

1.1 Coordinates

2-dimension (2D): $(2, 3)$



3-dimension (3D): $(1, \pi, 3)$



1.2 Vectors and Position Vectors

Definition 1.1: Equivalent Vectors


Two vectors are **equivalent** if one is a translation (pure shift without rotation) of the other.

Example 1.1. Given points $P = (1, 3)$, $Q = (5, 4)$ and $P' = (1, 1)$, $Q' = (5, 2)$. Let

$R = (4, 1)$ and $O = (0, 0)$. Then:

$$\overrightarrow{PQ} = \overrightarrow{P'Q'} = \overrightarrow{OR}$$

This could be indicated by the diagram below:



Definition 1.2: Position Vector

A vector whose initial point is at the origin $O(0, 0)$ is called a **position vector**. It is written using angle brackets:

$$\overrightarrow{OR} = \langle 4, 1 \rangle$$

Definition 1.3: Magnitude

The **length** or **magnitude** of a vector $\vec{v}$, denoted by $\|\vec{v}\|$, is the length of the line segment between its endpoints.

If $\vec{v} = \langle v_1, v_2 \rangle$, then its magnitude is given by:

$$\|\vec{v}\| = \sqrt{v_1^2 + v_2^2}$$

For example, using $\overrightarrow{PQ} = \langle 4, 1 \rangle$:

$$\|\overrightarrow{PQ}\| = \sqrt{4^2 + 1^2} = \sqrt{17}$$

1.3 Unit Vectors

A vector $\vec{v}$ is called a **unit vector** if its magnitude is 1, i.e., $\|\vec{v}\| = 1$.


Problem 1.1: Describing Unit Vectors

Describe all unit vectors in $\mathbb{R}^2$.

Solution 1.1

A vector $\vec{v} = \langle x, y \rangle$ is a unit vector if and only if $x^2 + y^2 = 1$. Equivalently, using polar parameterization:

$$\vec{v}(t) = \langle \cos t, \sin t \rangle \quad \text{for } 0 \leq t < 2\pi$$



Standard Basis Vectors:

- **2D Basis Vectors:** $\hat{i} = \langle 1, 0 \rangle$, $\hat{j} = \langle 0, 1 \rangle$
- **3D Basis Vectors:** $\hat{i} = \langle 1, 0, 0 \rangle$, $\hat{j} = \langle 0, 1, 0 \rangle$, $\hat{k} = \langle 0, 0, 1 \rangle$

1.4 Vector Operations


Addition and Subtraction:

$$\langle v_1, v_2 \rangle + \langle w_1, w_2 \rangle = \langle v_1 + w_1, v_2 + w_2 \rangle$$

$$\langle v_1, v_2 \rangle - \langle w_1, w_2 \rangle = \langle v_1 - w_1, v_2 - w_2 \rangle \quad (\text{Equivalent to } \vec{v} + (-1)\vec{w})$$

Scalar Multiplication: For any scalar $\lambda \in \mathbb{R}$ and vector $\vec{v} = \langle v_1, v_2 \rangle$:

$$\lambda \vec{v} = \langle \lambda v_1, \lambda v_2 \rangle$$



Important Note

Vectors are **not** designed to be divided! (Vector division is undefined).

1.5 Parallel Vectors

Definition 1.4: Parallel Vectors

Two non-zero vectors $\vec{v}$ and $\vec{w}$ ($\vec{v}, \vec{w} \neq \langle 0, 0 \rangle$) are *parallel* (denoted as $\vec{v} \parallel \vec{w}$) if there exists a non-zero scalar $\lambda \in \mathbb{R}$ such that:

$$\vec{w} = \lambda \vec{v}$$

Equivalent Geometric Definition: $\vec{v} \parallel \vec{w}$ if and only if both vectors define or lie on the *same line passing through the origin*.


2 Discussion Week 1: Distance Formulae and Geometric Shapes

Definition 2.1: Distance Formula in $\mathbb{R}^2$

Let $(x, y) \in \mathbb{R}^2$ and points $P = (x_1, y_1)$, $Q = (x_2, y_2)$. The distance between P and Q is given by:

$$d(P, Q) = \|\overrightarrow{PQ}\| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

This distance formula is a direct application of the Pythagorean theorem, where the distance $d(P, Q)$ forms the hypotenuse of a right-angled triangle with legs $|x_2 - x_1|$ and $|y_2 - y_1|$, as shown in the figure below.



The case in $\mathbb{R}^3$ follows similarly:

Definition 2.2: Distance Formula in $\mathbb{R}^3$

Let $(x, y, z) \in \mathbb{R}^3$ and points $P = (x_1, y_1, z_1)$, $Q = (x_2, y_2, z_2)$. The distance between P and Q is given by:

$$d(P, Q) = \|\overrightarrow{PQ}\| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

2.1 Equations of Geometric Shapes

Consider the equation $x^2 + y^2 = 4$.

In $\mathbb{R}^2$, this equation is equivalent to the set:

$$\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 4\}$$

This represents a circle with radius $r = 2$ centered at the origin. The distance from the origin $(0, 0)$ to any point (x, y) on this circle is constant and equal to 2, which can be verified directly using the distance formula:


$$\sqrt{(x - 0)^2 + (y - 0)^2} = 2$$

Hence, the geometry of the circle is directly implied by the Pythagorean theorem.

Now consider the equation $x^2 + y^2 = 1$ in three-dimensional space, $\mathbb{R}^3$. This set is formally expressed as:

$$\{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 1\}$$

Because z is completely unrestricted, the cross-section at any constant height $z = c$ forms the unit circle $x^2 + y^2 = 1$. Consequently, this equation describes a *right circular cylinder* extending infinitely along the z -axis, as shown.



2.2 Linear Equations across Dimensions

In $\mathbb{R}^2$, the general form of a linear equation is given by:

$$Ax + By + C = 0 \quad (A, B, C \in \mathbb{R}, \text{ with } A \text{ and } B \text{ not both zero})$$

This equation represents a *line* in two-dimensional space.

In $\mathbb{R}^3$, however, the exact same algebraic equation $Ax + By + C = 0$ represents a *plane* parallel to the z -axis (since z is a free variable), not a line.

To uniquely define a line in three-dimensional space, a single linear equation is insufficient; we require a system of *two* independent linear equations (representing the intersection of two non-parallel planes).

3 Vectors and Directions

Problem 3.1: Vector Directions

Let $\vec{v} = \langle 3, -4, 2 \rangle$. Find a unit vector in the same direction as $\vec{v}$.

Solution 3.1

To find a unit vector in the same direction as $\vec{v}$, we first compute its magnitude:

$$\|\vec{v}\| = \sqrt{3^2 + (-4)^2 + 2^2} = \sqrt{9 + 16 + 4} = \sqrt{29}$$

The unit vector in the same direction as $\vec{v}$ is then:

$$\hat{v} = \frac{\vec{v}}{\|\vec{v}\|} = \frac{1}{\sqrt{29}} \langle 3, -4, 2 \rangle$$


In general, for any non-zero vector $\vec{w}$,

$$\begin{aligned}\|\lambda \vec{w}\| &= \sqrt{(\lambda w_1)^2 + (\lambda w_2)^2 + (\lambda w_3)^2} \\ &= \sqrt{\lambda^2(w_1^2 + w_2^2 + w_3^2)} \\ &= |\lambda| \|\vec{w}\|\end{aligned}$$

Hence, if we want a unit vector in the same direction as $\vec{w}$, we can choose $\lambda = \frac{1}{\|\vec{w}\|}$, giving us:

$$\hat{w} = \frac{\vec{w}}{\|\vec{w}\|}$$

Right-Hand Rule The relative position of the x -, y -, and z -axes in $\mathbb{R}^3$ is determined by the *right-hand rule*. If you point your right thumb along the positive x -axis and your index finger along the positive y -axis, then your middle finger (perpendicular to both) will point along the positive z -axis.



Definition 3.1: Parameterizations

A parameterization is a function whose range is a given geometric object.


For example, a parameterization of the unit circle is given by:

$$f(t) = (\cos t, \sin t)$$

Example 3.1. To parameterize the line $y = 2x + 3$:

$$\begin{aligned}\ell(t) &= (t, 2t + 3) \\ &= \langle 0, 3 \rangle + \langle t, 2t \rangle\end{aligned}$$

The plot is then given:



To generalize, to parameterize a line in $\mathbb{R}^2$, we can first take a point on the line. Then, we

take some $\vec{v}$ in the direction of the line. Then, we can write the parameterization as:

$$\ell(t) = (p_1 + v_1t, p_2 + v_2t)$$

In $\mathbb{R}^3$, we can do the same thing.

Example 3.2. Parameterize the line passing through $P(1, 2, 0)$ and $Q(5, -1, 1)$.

$$\begin{aligned}\overrightarrow{PQ} &= \langle 5 - 1, -1 - 2, 1 - 0 \rangle \\ &= \langle 4, -3, 1 \rangle \\ \ell(t) &= (1, 2, 0) + t(4, -3, 1) \\ &= (1 + 4t, 2 - 3t, t)\end{aligned}$$

3.1 Dot Product

Definition 3.2: Dot Product

The dot product of two vectors $\vec{v} = \langle v_1, v_2 \rangle$ and $\vec{w} = \langle w_1, w_2 \rangle$ is defined as:

$$\vec{v} \cdot \vec{w} = v_1w_1 + v_2w_2$$

Example 3.3.

$$\begin{aligned}\vec{v} \cdot \vec{v} &= v_1^2 + v_2^2 = \|\vec{v}\|^2 \\ \vec{i} \cdot \vec{j} &= 1 \cdot 0 + 0 \cdot 1 = 0\end{aligned}$$

Problem 3.2: Dot Product and Angle

Find pairs of vectors with:

- Dot product > 0
- Dot product $= 0$
- Dot product < 0

Solution 3.2

- $\vec{v} \cdot \vec{v} = \|\vec{v}\|^2 > 0$ for any non-zero vector $\vec{v}$.
- $\vec{i} \cdot \vec{j} = 0$ since they are orthogonal.
- $\langle -1, -1 \rangle \cdot \langle 1, 1 \rangle = -1 - 1 = -2 < 0$.

To conclude, the dot products of any two vectors corresponds to the following situations:

- Dot product = 0 if and only if the two vectors are orthogonal (perpendicular).
- Dot product > 0 if the angle between the two vectors is acute (less than 90 degrees).
- Dot product < 0 if the angle between the two vectors is obtuse (greater than 90 degrees).


This is because $\vec{v} \cdot \vec{w} = \|\vec{v}\| \|\vec{w}\| \cos \theta$, where θ is the angle between the vectors. Hence,

$$\theta = \arccos \left(\frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \right)$$

Definition 3.3: Orthogonal Projection

Write $\vec{u} = \vec{u}_{\parallel} + \vec{u}_{\perp}$, where $\vec{u}_{\parallel} \parallel \vec{v}$ and $\vec{u}_{\perp} \perp \vec{u}_{\parallel}$. Then, $\vec{u}_{\parallel}$ is called the **orthogonal projection** of $\vec{u}$ onto $\vec{v}$.

Example 3.4. Below is an illustration of orthogonal projection:



By this, we could see that

$$\vec{u} = \vec{u}_{\parallel} + \vec{u}_{\perp}$$

Then, we solve for $\vec{u}_{\parallel}$ and $\vec{u}_{\perp}$.

Taking the dot product with $\vec{v}$:

$$\begin{aligned}
 \vec{u} \cdot \vec{v} &= (\vec{u}_{\parallel} + \vec{u}_{\perp}) \cdot \vec{v} \\
 &= \vec{u}_{\parallel} \cdot \vec{v} + \vec{u}_{\perp} \cdot \vec{v} \\
 &= \vec{u}_{\parallel} \cdot \vec{v} + 0 \quad (\text{since } \vec{u}_{\perp} \perp \vec{v}) \\
 &= \vec{u}_{\parallel} \cdot \vec{v}
 \end{aligned}$$

Let $\vec{e}_v = \frac{1}{\|\vec{v}\|} \vec{v}$ be the unit vector in the direction of $\vec{v}$, and write $\vec{u}_{\parallel} = \lambda \vec{e}_v$. Then:

$$\begin{aligned}
 \vec{u} \cdot \vec{v} &= (\lambda \vec{e}_v) \cdot (\|\vec{v}\| \vec{e}_v) \\
 &= \lambda \|\vec{v}\| (\vec{e}_v \cdot \vec{e}_v) \\
 &= \lambda \|\vec{v}\| \|\vec{e}_v\|^2 \\
 &= \lambda \|\vec{v}\| \quad (\text{since } \|\vec{e}_v\| = 1)
 \end{aligned}$$

Solving for λ :

$$\lambda = \frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|}$$

Thus, the parallel projection vector $\vec{u}_{\parallel}$ and the orthogonal component $\vec{u}_{\perp}$ are given by:

$$\begin{aligned}
 \vec{u}_{\parallel} &= \left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|} \right) \vec{e}_v \\
 &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \vec{v} \\
 &= \frac{\vec{u} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \vec{v} \\
 \vec{u}_{\perp} &= \vec{u} - \vec{u}_{\parallel} \\
 &= \vec{u} - \frac{\vec{u} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \vec{v}
 \end{aligned}$$

Furthermore, the magnitude of the parallel projection is:

$$\|\vec{u}_{\parallel}\| = \|\vec{u}\| |\cos \theta|$$

Problem 3.3: Vector Projection


Find the projection of $\vec{u} = \langle 3, 2, 2 \rangle$ onto $\vec{v} = \langle 1, -2, 0 \rangle$

Solution 3.3

$$\begin{aligned}\vec{u}_{\parallel} &= \frac{3 \cdot 1 + 2 \cdot (-1) + 2 \cdot 0}{1^2 + (-1)^2 + 0^2} \langle 1, -1, 0 \rangle \\ &= \frac{1}{2} \langle 1, -1, 0 \rangle \\ &= \left\langle \frac{1}{2}, -\frac{1}{2}, 0 \right\rangle\end{aligned}$$

Problem 3.4: Acceleration on an Inclined Plane

Consider a block sliding down an inclined plane with an angle of inclination α .



Solution 3.4

To find the magnitude of acceleration down the incline, we project the gravitational force $\vec{F}_g$ along the incline direction:

$$\begin{aligned}\text{acceleration} &= \left\| \vec{F}_{g\parallel} \right\| \\ &= \left\| \vec{F}_g \right\| |\cos \theta| \\ &= \left\| \vec{F}_g \right\| \sin \alpha = 9.8 \sin \alpha \quad (\text{m/s}^2)\end{aligned}$$

Problem 3.5: Vector Projection

Find the projection of $\vec{u} = \langle 3, 2, 2 \rangle$ onto $\vec{v} = \langle 1, -1, 0 \rangle$.

Solution 3.5

Using the vector projection formula:


$$\begin{aligned}\vec{u}_{\parallel} &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \vec{v} \\ &= \frac{3(1) + 2(-1) + 2(0)}{1^2 + (-1)^2 + 0^2} \langle 1, -1, 0 \rangle \\ &= \frac{3 - 2 + 0}{1 + 1 + 0} \langle 1, -1, 0 \rangle \\ &= \frac{1}{2} \langle 1, -1, 0 \rangle \\ &= \left\langle \frac{1}{2}, -\frac{1}{2}, 0 \right\rangle\end{aligned}$$

3.2 Determinants and Cross Products

Definition 3.4: Vector Cross Product

Let $\vec{u}, \vec{v} \in \mathbb{R}^3$. The **cross product** $\vec{u} \times \vec{v}$ is a vector in $\mathbb{R}^3$ satisfying the following three geometric properties:

1. **Orthogonality:** $\vec{u} \times \vec{v} \perp \vec{u}$ and $\vec{u} \times \vec{v} \perp \vec{v}$.
2. **Magnitude:** $\|\vec{u} \times \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin \theta$, which equals the area of the parallelogram formed by $\vec{u}$ and $\vec{v}$.
3. **Orientation:** The ordered triple $(\vec{u}, \vec{v}, \vec{u} \times \vec{v})$ is right-handed (positively oriented).



From Property (2), we have for the standard basis vectors:

$$\|\vec{i} \times \vec{i}\| = \|\vec{i}\| \|\vec{i}\| \sin 0^\circ = 0 \implies \vec{i} \times \vec{i} = \vec{0}$$

And by the Right-Hand Rule:

$$\vec{i} \times \vec{j} = \vec{k}$$


Definition 3.5: Determinants

For a 2×2 matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, the determinant is defined as:

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$$

This is denoted as $\det(A)$ or $|A|$ for a 2×2 matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$.

For a 2×2 matrix, the determinant represents the area of the parallelogram spanned by $\langle a, c \rangle$ and $\langle b, d \rangle$ (the column vectors) or $\langle a, b \rangle$ and $\langle c, d \rangle$.




For a 3×3 matrix $B = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$, the determinant is defined as:

$$\det \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

$$= aei + bfg + cdh - ceg - bdi - afh$$

$\det \begin{pmatrix} \vec{u} \\ \vec{v} \\ \vec{w} \end{pmatrix}$ represents the volume of the parallelepiped spanned by $\vec{u}$, $\vec{v}$, and $\vec{w}$.



The cross product of two vectors $\vec{u} = \langle u_1, u_2, u_3 \rangle$ and $\vec{v} = \langle v_1, v_2, v_3 \rangle$ can be computed using the determinant of a 3×3 matrix:

$$\vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} = \vec{i} \begin{vmatrix} u_2 & u_3 \\ v_2 & v_3 \end{vmatrix} - \vec{j} \begin{vmatrix} u_1 & u_3 \\ v_1 & v_3 \end{vmatrix} + \vec{k} \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix}$$

Problem 3.6: Cross Product

Let $\vec{v} = \langle 3, 2, -1 \rangle$, $\vec{w} = \langle 4, 1, -3 \rangle$. Find $\vec{v} \times \vec{w}$.

Solution 3.6

$$\begin{aligned}\vec{v} \times \vec{w} &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 3 & 2 & -1 \\ 4 & 1 & -3 \end{vmatrix} \\ &= \vec{i} \begin{vmatrix} 2 & -1 \\ 1 & -3 \end{vmatrix} - \vec{j} \begin{vmatrix} 3 & -1 \\ 4 & -3 \end{vmatrix} + \vec{k} \begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix} \\ &= \vec{i}(2(-3) - (-1)(1)) - \vec{j}(3(-3) - (-1)(4)) + \vec{k}(3(1) - 2(4)) \\ &= \vec{i}(-6 + 1) - \vec{j}(-9 + 4) + \vec{k}(3 - 8) \\ &= \vec{i}(-5) - \vec{j}(-5) + \vec{k}(-5) \\ &= -5\vec{i} + 5\vec{j} - 5\vec{k}\end{aligned}$$

3.2.1 Properties of the Cross Product

- $\vec{v} \times \vec{w} \perp \vec{v}$ and $\vec{v} \times \vec{w} \perp \vec{w}$. This implies that $\vec{v} \cdot (\vec{v} \times \vec{w}) = 0$ and $\vec{w} \cdot (\vec{v} \times \vec{w}) = 0$.
- $\{\vec{v}, \vec{w}, \vec{v} \times \vec{w}\}$ forms a right-handed system.
- $\|\vec{v} \times \vec{w}\|$ represents the area of the parallelogram spanned by $\vec{v}$ and $\vec{w}$.

3.2.2 Algebraic Rules of the Cross Product

- Anti-commutativity: $\vec{v} \times \vec{w} = -(\vec{w} \times \vec{v})$
- Parallel vectors: $\vec{v} \times \vec{w} = \vec{0} \iff \vec{v} = \lambda\vec{w} \text{ or } \vec{w} = \vec{0}$.
- Scalar multiplication: $\lambda(\vec{v} \times \vec{w}) = (\lambda\vec{v}) \times \vec{w} = \vec{v} \times (\lambda\vec{w})$
- Distributivity: $(\vec{u} + \vec{v}) \times \vec{w} = \vec{u} \times \vec{w} + \vec{v} \times \vec{w}$
- Standard basis relations:

$$\vec{i} \times \vec{j} = \vec{k}, \quad \vec{j} \times \vec{k} = \vec{i}, \quad \vec{k} \times \vec{i} = \vec{j}$$


Hence, we may expand the cross product of two vectors $\vec{v} = \langle v_1, v_2, v_3 \rangle$ and $\vec{w} = \langle w_1, w_2, w_3 \rangle$ component-wise:

$$\begin{aligned}
\vec{v} \times \vec{w} &= (v_1 \vec{i} + v_2 \vec{j} + v_3 \vec{k}) \times (w_1 \vec{i} + w_2 \vec{j} + w_3 \vec{k}) \\
&= v_1 w_1 (\vec{i} \times \vec{i}) + v_1 w_2 (\vec{i} \times \vec{j}) + v_1 w_3 (\vec{i} \times \vec{k}) \\
&\quad + v_2 w_1 (\vec{j} \times \vec{i}) + v_2 w_2 (\vec{j} \times \vec{j}) + v_2 w_3 (\vec{j} \times \vec{k}) \\
&\quad + v_3 w_1 (\vec{k} \times \vec{i}) + v_3 w_2 (\vec{k} \times \vec{j}) + v_3 w_3 (\vec{k} \times \vec{k}) \\
&= v_1 w_2 \vec{k} - v_1 w_3 \vec{j} - v_2 w_1 \vec{k} + v_2 w_3 \vec{i} + v_3 w_1 \vec{j} - v_3 w_2 \vec{i} \\
&= (v_2 w_3 - v_3 w_2) \vec{i} + (v_3 w_1 - v_1 w_3) \vec{j} + (v_1 w_2 - v_2 w_1) \vec{k}
\end{aligned}$$

3.3 Scalar Triple Product

Given $\vec{u}, \vec{v}, \vec{w} \in \mathbb{R}^3$, the quantity $\vec{u} \cdot (\vec{v} \times \vec{w})$ is defined as the **scalar triple product**.

The absolute value $|\vec{u} \cdot (\vec{v} \times \vec{w})|$ gives the geometric volume of the parallelepiped spanned by $\vec{u}$, $\vec{v}$, and $\vec{w}$:



The sign of $\vec{u} \cdot (\vec{v} \times \vec{w})$ indicates the orientation of the vector basis.

$\vec{u} \cdot (\vec{v} \times \vec{w}) > 0$ if the ordered triple $(\vec{u}, \vec{v}, \vec{w})$ forms a **right-handed system** (i.e., $\vec{u}$ points to the same side of the plane spanned by $\vec{v}$ and $\vec{w}$ as $\vec{v} \times \vec{w}$).

4 Discussion Week 2: Quadric Surfaces and Traces

Definition 4.1: Quadric Surface

A quadric surface is defined by a quadratic equation in three variables x , y , and z :

$$Ax^2 + By^2 + Cz^2 + Dxy + Eyz + Fxz + ax + by + cz + d = 0$$


, where $A, B, C, D, E, F, a, b, c, d$ are constants, and at least one of A, B, C, D, E, F is non-zero.

Definition 4.2: Trace

A trace is the intersection of a quadric surface with a plane, usually by taking slices parallel to the coordinate planes.


Note that when our surface can be written by $z = f(x, y)$, the traces form level sets of f .

Example 4.1. Consider the quadratic equation $z = x^2 - y^2$. Now take the traces in $z = -1$, $z = 0$, and $z = 1$. Make a sketch of the traces in $x = -1$, $x = 0$, and $x = 1$. In general, the traces in $x = k$ when k is a constant are hyperbolas with equations $z = k^2 - y^2$.




Now take the traces in $z = -1$, $z = 0$, and $z = 1$.


Trace at $z = -1$




$$y^2 - x^2 = 1$$




Trace at $z = 0$




$$y = \pm x$$



Trace at $z = 1$




$$x^2 - y^2 = 1$$




Make a sketch of the traces in $x = -1$, $x = 0$, and $x = 1$. Slicing the surface with vertical planes $x = k$ parallel to the yz -plane yields parabolic trace curves.


Trace at $x = -1$




$$z = 1 - y^2$$




Trace at $x = 0$




$$z = -y^2$$



Trace at $x = 1$



$$z = 1 - y^2$$



Ellipsoids are another family of quadric surfaces. The general equation of an ellipsoid is given by:

$$\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 + \left(\frac{z}{c}\right)^2 = 1$$

5 Planes and Surfaces

5.1 Determining a Plane

A plane $\mathcal{P}$ in $\mathbb{R}^3$ can be uniquely determined by any of the following geometric data:

- A point Q on the plane and a normal vector $\vec{n} \perp \mathcal{P}$.
- Three non-collinear points.
- A general linear equation $ax + by + cz + d = 0$.
- Two intersecting lines (or two non-parallel direction vectors).
- Two parallel distinct lines.
- A line $\mathcal{L}$ and a point $Q \notin \mathcal{L}$.

5.2 Mathematical Formulations of a Plane

Case 1: Plane passing through the origin $O(0, 0, 0)$

A vector $\vec{v} = \langle x, y, z \rangle$ lies in $\mathcal{P} \iff \vec{v} \perp \vec{n}$, where $\vec{n} = \langle a, b, c \rangle$:

$$\vec{v} \in \mathcal{P} \iff \vec{v} \cdot \vec{n} = 0 \iff ax + by + cz = 0$$

In set notation:

$$\mathcal{P} = \{ \langle x, y, z \rangle \mid ax + by + cz = 0 \}$$

Case 2: Plane passing through a point

$Q(x_0, y_0, z_0)$

Let Q be a known point on $\mathcal{P}$ with position vector $\overrightarrow{OQ} = \langle x_0, y_0, z_0 \rangle$. Any point $W(x, y, z)$ on the plane satisfies:

$$(\overrightarrow{OW} - \overrightarrow{OQ}) \perp \vec{n} \iff (\overrightarrow{OW} - \overrightarrow{OQ}) \cdot \vec{n} = 0$$


Expanding this dot product yields the point-normal form:

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$$

or equivalently:

$$ax + by + cz = ax_0 + by_0 + cz_0$$

We can always scale the coefficients a, b, c, d in $ax + by + cz = d$ by $\frac{1}{\|\vec{n}\|}$ to obtain a unit normal vector $\hat{n} = \frac{\langle a, b, c \rangle}{\sqrt{a^2 + b^2 + c^2}}$.



5.3 Properties of Normal Vectors

For any set of vectors $\{\vec{u}, \vec{v}, \vec{w}\}$ where $\vec{v} \nparallel \vec{w}$, the following statements are equivalent:

- $\vec{u} \perp \vec{v}$ and $\vec{u} \perp \vec{w}$.
- $\vec{u} \parallel (\vec{v} \times \vec{w})$.
- $\vec{u} \perp \mathcal{P}$, where $\mathcal{P} = \text{Span}\{\vec{v}, \vec{w}\}$.

Problem 5.1: Finding Normal Vectors

Find a normal vector $\vec{n}$ to the plane $\mathcal{P}$ given by the equation $(x - 1) - (y - 1) = 0$.

Solution 5.1

The equation simplifies to $x - y = 0$. We can extract or compute the normal vector in two ways:

Method 1 (Direct Inspection): From the general form $ax + by + cz = d$, the coefficients of x, y, z directly give the normal vector:

$$\vec{n} = \langle 1, -1, 0 \rangle$$

Method 2 (Cross Product of Parallel Plane Vectors): Two vectors parallel to the plane $\mathcal{P}$ are $\vec{v}_1 = \langle 0, 0, 1 \rangle$ and $\vec{v}_2 = \langle 1, 1, 0 \rangle$. Taking their cross product gives:

$$\vec{n} = \vec{v}_1 \times \vec{v}_2 = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{vmatrix} = \vec{i}(0 - 1) - \vec{j}(0 - 1) + \vec{k}(0 - 0) = \langle -1, 1, 0 \rangle$$

Note that $\langle -1, 1, 0 \rangle = -\langle 1, -1, 0 \rangle$, which represents the same normal line direction.

Problem 5.2: Finding a Plane From Three Points

Find the equation of the plane containing the points $P(1, 0, 1)$, $Q(0, 2, 0)$, and $R(2, 1, 0)$.

Solution 5.2


Step 1: Construct two displacement vectors in the plane

$$\overrightarrow{PQ} = \langle 0 - 1, 2 - 0, 0 - 1 \rangle = \langle -1, 2, -1 \rangle$$

$$\overrightarrow{PR} = \langle 2 - 1, 1 - 0, 0 - 1 \rangle = \langle 1, 1, -1 \rangle$$

Step 2: Compute the normal vector via cross product

$$\vec{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -1 & 2 & -1 \\ 1 & 1 & -1 \end{vmatrix} = \langle -1, -2, -3 \rangle$$



Step 3: Write the plane equation using point $P(1, 0, 1)$

$$-1(x - 1) - 2(y - 0) - 3(z - 1) = 0$$

Simplifying yields:

$$x + 2y + 3z = 4$$

6 Spheres and Planes

6.1 Determining a Sphere

A sphere in $\mathbb{R}^3$ can be uniquely determined by any of the following sets of data:

- A center point $(a, b, c) \in \mathbb{R}^3$ and a radius $R \in \mathbb{R}^{>0}$.
- Two points representing the endpoints of a diameter.
- Four non-coplanar points.

- A plane tangent to the sphere together with three points lying on the sphere.

6.2 Plane-Sphere Intersections

Example 6.1. Consider the intersection of the sphere $x^2 + y^2 + z^2 = 4$ with the line $\ell(t) = (t, 0, 1)$ passing through $(1, 0, 1)$ and $(-1, 0, 1)$.

Substituting $\ell(t)$ into the sphere equation yields:

$$\begin{aligned} t^2 + 0^2 + 1^2 &= 4 \\ t^2 + 1 &= 4 \\ t^2 = 3 &\implies t = \pm\sqrt{3} \end{aligned}$$

Thus, the line intersects the sphere at $\ell(\sqrt{3}) = (\sqrt{3}, 0, 1)$ and $\ell(-\sqrt{3}) = (-\sqrt{3}, 0, 1)$.

6.3 Traces of Surfaces

Definition 6.1: Traces

A **trace** of a surface is the intersection curve obtained by intersecting the surface with a parallel coordinate plane of the form $x = k$, $y = k$, or $z = k$.

Example 6.2 (Traces of a Sphere). Consider the sphere given by:

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = R^2$$


Taking the trace with the plane $z = k$ yields the equation:

$$(x - a)^2 + (y - b)^2 = R^2 - (k - c)^2$$

Depending on the plane distance k , we have three distinct cases:

- **Case 1** ($|k - c| = R$): The trace is a single point at (a, b, k) .
- **Case 2** ($|k - c| < R$): The trace is a circle of radius $\sqrt{R^2 - (k - c)^2}$ centered at (a, b, k) .

- **Case 3** ($|k - c| > R$): The trace is empty (no real intersection).



Example 6.3. Find the trace of the plane $\mathcal{P} = \{(x, y, z) \mid -2x + 6y - 5z = 4\}$ with the plane $x = -1$:

$$-2(-1) + 6y - 5z = 4$$

$$2 + 6y - 5z = 4$$

$$6y - 5z = 2 \implies z = \frac{6}{5}y - \frac{2}{5}$$

The trace is a straight line in the plane $x = -1$.

6.4 Quadric Surfaces


A general **quadric surface** in $\mathbb{R}^3$ is given by the second-degree equation:

$$Ax^2 + By^2 + Cz^2 + Dxy + Eyz + Fzx + ax + by + cz + d = 0$$

When $D = E = F = 0$, the surface is in **standard position** (axes aligned with coordinate axes).

Example 6.4. The surface $x^2 + z^2 = 1$ is a circular cylinder aligned along the y -axis with radius 1.


- $z = 0$ trace: $x^2 = 1 \implies x = \pm 1$ (parallel lines along y).
- $x = 0$ trace: $z^2 = 1 \implies z = \pm 1$ (parallel lines along y).



Example 6.5. Consider the surface given by $x^2 + 4y^2 - z^2 = -1$.

Traces:

- $x = 0$ **trace:** $4y^2 - z^2 = -1 \implies z = \pm\sqrt{4y^2 + 1}$. As $|y| \rightarrow \infty$, $z \approx \pm 2y$ (Hyperbola).
- $y = 0$ **trace:** $x^2 - z^2 = -1 \implies z = \pm\sqrt{x^2 + 1}$. As $|x| \rightarrow \infty$, $z \approx \pm x$ (Hyperbola).
- $z = k$ **trace:** $x^2 + 4y^2 = k^2 - 1$:
 - $|k| = 1$: Single point at $(0, 0, \pm 1)$.
 - $|k| < 1$: No trace (empty set).
 - $|k| > 1$: Ellipse given by $x^2 + 4y^2 = k^2 - 1$.



Why do cross-sectional ellipses fall on the surface?

For any fixed height $z = k$ with $|k| > 1$, substituting $z = k$ fixes the value $k^2 - 1 > 0$ on the right side of $x^2 + 4y^2 = k^2 - 1$. This traces a locus of points (x, y, k) whose horizontal distance components continuously satisfy the surface equation for that slice.

7 Vector-Valued Functions

7.1 Coordinate Systems in 3D Space

Specifying a point or giving directions in space can be accomplished in multiple ways:

- Giving a direction and a distance.
- Using Cartesian coordinates $(d_1, d_2, d_3) = (x, y, z)$.
- Specifying an angle relative to a reference axis along with a magnitude.
- Referencing a relative landmark.
- Using geographic coordinates like latitude and longitude.

7.1.1 Polar Coordinates (2D)

In the xy -plane, a point is specified by its radial distance r from the origin and an angle θ relative to the positive x -axis: (r, θ) .

$$r = \sqrt{x^2 + y^2} \quad (r \geq 0)$$

While $\theta = \arctan\left(\frac{y}{x}\right)$ is commonly used, it is undefined when $x = 0$. The foundational algebraic transformation between systems is given by:

$$x = r \cos \theta$$

$$y = r \sin \theta$$

7.1.2 Cylindrical Coordinates

Cylindrical coordinates (r, θ, z) extend polar coordinates into 3D space by preserving the standard Cartesian z -axis:

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$z = z$$

7.1.3 Spherical Coordinates

Spherical coordinates are represented as (ρ, θ, ϕ) , where ρ is the distance from the origin, θ is the azimuthal angle in the xy -plane, and ϕ is the polar angle measured down from the positive z -axis.

$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$


$$z = \rho \cos \phi$$

The standard domain boundaries are:

$$\rho \geq 0$$

$$0 \leq \theta < 2\pi$$

$$0 \leq \phi \leq \pi$$



7.2 Coordinate System Transformations

Problem 7.1: Conversion Between Coordinate Systems

Convert the surface equation $x^2 + \frac{y^2}{a^2} = z^2$ into cylindrical and spherical coordinates.

Solution 7.1

Cylindrical Coordinates:

Substituting $x = r \cos \theta$, $y = r \sin \theta$, and $z = z$ into the original equation yields:

$$\begin{aligned} (r \cos \theta)^2 + \frac{(r \sin \theta)^2}{a^2} &= z^2 \\ r^2 \cos^2 \theta + \frac{r^2 \sin^2 \theta}{a^2} &= z^2 \\ r^2 \left(\cos^2 \theta + \frac{\sin^2 \theta}{a^2} \right) &= z^2 \end{aligned}$$

Spherical Coordinates:

Substituting $x = \rho \sin \phi \cos \theta$, $y = \rho \sin \phi \sin \theta$, and $z = \rho \cos \phi$ into the original equation yields:

$$\begin{aligned} (\rho \sin \phi \cos \theta)^2 + \frac{(\rho \sin \phi \sin \theta)^2}{a^2} &= (\rho \cos \phi)^2 \\ \rho^2 \sin^2 \phi \cos^2 \theta + \frac{\rho^2 \sin^2 \phi \sin^2 \theta}{a^2} &= \rho^2 \cos^2 \phi \\ \rho^2 \sin^2 \phi \left(\cos^2 \theta + \frac{\sin^2 \theta}{a^2} \right) &= \rho^2 \cos^2 \phi \end{aligned}$$

Assuming $\rho \neq 0$, we divide both sides by ρ^2 :

$$\begin{aligned}\sin^2 \phi \left(\cos^2 \theta + \frac{\sin^2 \theta}{a^2} \right) &= \cos^2 \phi \\ \cos^2 \theta + \frac{\sin^2 \theta}{a^2} &= \frac{\cos^2 \phi}{\sin^2 \phi} \\ \cos^2 \theta + \frac{\sin^2 \theta}{a^2} &= \cot^2 \phi\end{aligned}$$

7.3 Vector-Valued Functions


Definition 7.1: Vector-Valued Functions

A vector-valued function maps one or more real parameters to a space vector:

$$\vec{r}(t) = \langle x(t), y(t), z(t) \rangle, \quad \vec{r} : I \rightarrow \mathbb{R}^3$$

where $I \subseteq \mathbb{R}$ is an interval. This traces out a spatial curve parametrized by t .

Example 7.1. Consider the vector function $\vec{r}(t) = \langle \cos t, \sin t, 1 \rangle$ for $0 \leq t < 2\pi$. This describes a unit circle in 3D space elevated to the horizontal plane $z = 1$.



Problem 7.2: Finding Collisions and Intersections

Let $\vec{r}_1(t) = \left\langle t^2 + \frac{t}{2}, t - t^3 \right\rangle$ and $\vec{r}_2(t) = \langle t, t^2 \rangle$. Find all collision points and structural intersections between the curves.

Solution 7.2

1. Collision Points (Same location at same time t):

Setting $\vec{r}_1(t) = \vec{r}_2(t)$ gives:

$$\left\langle t^2 + \frac{t}{2}, t - t^3 \right\rangle = \langle t, t^2 \rangle$$

Equating the individual components gives the system:

$$\begin{aligned} t^2 + \frac{t}{2} &= t \\ t - t^3 &= t^2 \end{aligned}$$

Solving the first equation for t :

$$\begin{aligned} t^2 + \frac{t}{2} - t &= 0 \\ t^2 - \frac{t}{2} &= 0 \\ t \left(t - \frac{1}{2} \right) &= 0 \implies t = 0 \quad \text{or} \quad t = \frac{1}{2} \end{aligned}$$

Now, we verify both parameters in the second component equation $t - t^3 = t^2$:

$$\begin{aligned} \text{For } t = 0 : \quad 0 - (0)^3 &= (0)^2 \implies 0 = 0 \quad (\text{Valid}) \\ \text{For } t = \frac{1}{2} : \quad \frac{1}{2} - \left(\frac{1}{2} \right)^3 &= \frac{1}{2} - \frac{1}{8} = \frac{3}{8} \\ \left(\frac{1}{2} \right)^2 &= \frac{1}{4} = \frac{2}{8} \neq \frac{3}{8} \quad (\text{Invalid}) \end{aligned}$$

Therefore, $t = 0$ is the only valid collision time. Evaluating $\vec{r}_1(0)$:

$$\vec{r}_1(0) = \left\langle 0^2 + \frac{0}{2}, 0 - 0^3 \right\rangle = \langle 0, 0 \rangle$$

The collision point is $(0, 0)$.

2. Intersection Points (Same location at potentially different times t and

s):

Setting $\vec{r}_1(t) = \vec{r}_2(s)$ gives:

$$\left\langle t^2 + \frac{t}{2}, t - t^3 \right\rangle = \langle s, s^2 \rangle$$

Equating components yields:

$$s = t^2 + \frac{t}{2} \tag{1}$$

$$t - t^3 = s^2 \tag{2}$$

Substituting equation (1) into equation (2):

$$\begin{aligned} t - t^3 &= \left(t^2 + \frac{t}{2} \right)^2 \\ t - t^3 &= t^4 + t^3 + \frac{t^2}{4} \\ 0 &= t^4 + 2t^3 + \frac{t^2}{4} - t \\ 0 &= t \left(t^3 + 2t^2 + \frac{t}{4} - 1 \right) \end{aligned}$$

Setting each factor to zero:

$$t = 0 \implies s = 0^2 + \frac{0}{2} = 0$$

$\implies$ Intersection point $(0, 0)$

$$t^3 + 2t^2 + \frac{t}{4} - 1 = 0 \implies t \approx 0.576$$

Solving for s when $t \approx 0.576$:

$$s = (0.576)^2 + \frac{0.576}{2}$$

$$s \approx 0.332 + 0.288 \approx 0.620$$

Evaluating the spatial coordinates using $\vec{r}_2(s)$:


$$x = s \approx 0.620$$

$$y = s^2 \approx (0.620)^2 \approx 0.384$$


Hence, the two spatial intersection points are $(0, 0)$ and $(0.620, 0.384)$.

7.4 Gallery of Parametric Space Curves


Below are standard parametric 3D space curves used in multivariable trajectory analysis:



$$\vec{r}(t) = \langle \cos t, \sin t, t \rangle$$



$$\vec{r}(s) = \langle \cos s, \sin s, \sin(4s) \rangle$$



$$\vec{r}(t) = \langle \cos t, \sin t, 4 \sin t \rangle$$

7.5 Parameterization of Curves and Surfaces

Problem 7.3: Parameterization of a Curve on a Surface

Find a parameterization for the curve given by the intersection of the surfaces $y = \sqrt{x^2 + z^2}$ and $x = z^2$.

Solution 7.3

Notice that for any choice of $z \in \mathbb{R}$, the value of x is uniquely determined by $x = z^2$.

We can set $z = t$ as our parameter. Then,

$$x = t^2$$

$$z = t$$

$$y = \sqrt{(t^2)^2 + t^2} = \sqrt{t^4 + t^2} = |t|\sqrt{t^2 + 1}$$

Thus, a vector parameterization of the curve is:

$$\vec{r}(t) = \langle t^2, |t|\sqrt{t^2 + 1}, t \rangle, \quad t \in \mathbb{R}$$

7.6 Calculus of Vector-Valued Functions

Calculus operations on vector-valued functions are defined component-wise.

For a vector-valued function $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$, limits, derivatives, and integrals are computed as follows:

$$\lim_{t \rightarrow t_0} \vec{r}(t) = \left\langle \lim_{t \rightarrow t_0} x(t), \lim_{t \rightarrow t_0} y(t), \lim_{t \rightarrow t_0} z(t) \right\rangle$$

$$\vec{r}'(t) = \langle x'(t), y'(t), z'(t) \rangle$$

$$\int_a^b \vec{r}(t) dt = \left\langle \int_a^b x(t) dt, \int_a^b y(t) dt, \int_a^b z(t) dt \right\rangle$$


Example 7.2 (Tangent Vectors and Parameterizing Tangent Lines). Given the circular helix $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$.

Differentiating component-wise gives:

$$\vec{r}'(t) = \langle -\sin t, \cos t, 1 \rangle$$

$$\vec{r}'(0) = \langle 0, 1, 1 \rangle$$

The derivative $\vec{r}'(0)$ yields a vector representing the tangent direction of the curve at $t = 0$.



To parameterize the tangent line $\ell(t)$ to the helix at $t = 0$:

$$\vec{r}(0) = \langle 1, 0, 0 \rangle$$

Thus,

$$\begin{aligned} \ell(t) &= \vec{r}(0) + t\vec{r}'(0) \\ &= \langle 1, 0, 0 \rangle + t\langle 0, 1, 1 \rangle = \langle 1, t, t \rangle \end{aligned}$$

Problem 7.4: Derivative and Singular Points of a Cycloid

Find the derivative of the vector-valued function $\vec{r}(t) = \langle t - \sin t, 1 - \cos t \rangle$ and evaluate it at $t = \frac{\pi}{2}, \pi, 2\pi$.

Solution 7.4

Differentiating each component with respect to t :


$$\begin{aligned} \vec{r}'(t) &= \left\langle \frac{d}{dt}(t - \sin t), \frac{d}{dt}(1 - \cos t) \right\rangle \\ &= \langle 1 - \cos t, \sin t \rangle \end{aligned}$$

Evaluating at specific parameter values:

$$\vec{r}'\left(\frac{\pi}{2}\right) = \left\langle 1 - \cos \frac{\pi}{2}, \sin \frac{\pi}{2} \right\rangle = \langle 1, 1 \rangle$$

$$\vec{r}'(\pi) = \langle 1 - \cos \pi, \sin \pi \rangle = \langle 2, 0 \rangle$$

$$\vec{r}'(2\pi) = \langle 1 - \cos 2\pi, \sin 2\pi \rangle = \langle 0, 0 \rangle$$



Notice from the graph of the cycloid that at $t = 2\pi$, the curve forms a sharp sharp corner (cusp). At this point, $\vec{r}'(2\pi) = \langle 0, 0 \rangle$. Although the vector function $\vec{r}(t)$ is differentiable everywhere, the tangent line direction is undefined at the cusp because the derivative vector vanishes.

7.7 Properties of Vector Derivatives & Integration

Problem 7.5: Orthogonality Property for Constant-Length Vector Functions

If $\|\vec{r}(t)\| = c$ for some constant c , what does this imply about the relationship between $\vec{r}(t)$ and its derivative $\vec{r}'(t)$?

Solution 7.5

Since $\|\vec{r}(t)\| = c$, taking the square of both sides yields:

$$\|\vec{r}(t)\|^2 = c^2 \implies \vec{r}(t) \cdot \vec{r}(t) = c^2$$

Differentiating both sides with respect to t using the product rule for dot products:

$$\begin{aligned}\frac{d}{dt}(\vec{r}(t) \cdot \vec{r}(t)) &= \frac{d}{dt}(c^2) \\ \vec{r}'(t) \cdot \vec{r}(t) + \vec{r}(t) \cdot \vec{r}'(t) &= 0 \\ 2\vec{r}(t) \cdot \vec{r}'(t) &= 0 \implies \vec{r}(t) \cdot \vec{r}'(t) = 0\end{aligned}$$

Therefore, $\vec{r}(t) \perp \vec{r}'(t)$ for all t . (The position vector is always orthogonal to its velocity vector).

Product Rule Verification for Dot Products

Given two vector-valued functions $\vec{r}_1(t), \vec{r}_2(t) \in \mathbb{R}^2$:

$$\begin{aligned}\frac{d}{dt}[\vec{r}_1(t) \cdot \vec{r}_2(t)] &= \frac{d}{dt}[x_1(t)x_2(t) + y_1(t)y_2(t)] \\ &= x_1'(t)x_2(t) + x_1(t)x_2'(t) + y_1'(t)y_2(t) + y_1(t)y_2'(t) \\ &= \vec{r}_1'(t) \cdot \vec{r}_2(t) + \vec{r}_1(t) \cdot \vec{r}_2'(t)\end{aligned}$$

Problem 7.6: Initial Value Problem for Motion

A particle moves with velocity $\vec{v}(t) = \langle 3t^2 - 2, \cos(\pi t), \sqrt{t} \rangle$. Find its position vector $\vec{r}(t)$, given the initial condition $\vec{r}(0) = \langle 2, 1, 1 \rangle$.

Solution 7.6

Integrating $\vec{v}(t)$ component-wise:

$$\begin{aligned}\vec{r}(t) &= \int \vec{v}(t) dt \\ &= \left\langle \int (3t^2 - 2) dt, \int \cos(\pi t) dt, \int t^{1/2} dt \right\rangle \\ &= \left\langle t^3 - 2t + C_1, \frac{1}{\pi} \sin(\pi t) + C_2, \frac{2}{3} t^{3/2} + C_3 \right\rangle\end{aligned}$$

Applying the initial condition $\vec{r}(0) = \langle 2, 1, 1 \rangle$:

$$\vec{r}(0) = \langle C_1, C_2, C_3 \rangle = \langle 2, 1, 1 \rangle \implies C_1 = 2, C_2 = 1, C_3 = 1$$

Therefore,

$$\vec{r}(t) = \left\langle t^3 - 2t + 2, \frac{1}{\pi} \sin(\pi t) + 1, \frac{2}{3} t^{3/2} + 1 \right\rangle$$

Note on Circular Motion: If given acceleration $\vec{r}''(t) = \langle -\cos t, -\sin t, 0 \rangle$, to uniquely guarantee that $\vec{r}(t)$ describes a circle centered at the origin, appropriate initial conditions for both position $\vec{r}(0)$ and velocity $\vec{r}'(0)$ must be specified.

7.8 Arc Length of Curves

The arc length s of a smooth curve $\vec{r}(t) = \langle x(t), y(t) \rangle$ on $t \in [a, b]$ is derived as the limit of polygonal approximations:

$$s = \lim_{\Delta t \rightarrow 0} \sum \sqrt{\left(\frac{\Delta x}{\Delta t}\right)^2 + \left(\frac{\Delta y}{\Delta t}\right)^2} \Delta t = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_a^b \|\vec{r}'(t)\| dt$$

Example 7.3 (Arc Length of a Helix). Find the arc length of $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$ for $t \in [0, 2\pi]$.

First, evaluate the speed $\|\vec{r}'(t)\|$:

$$\vec{r}'(t) = \langle -\sin t, \cos t, 1 \rangle \implies \|\vec{r}'(t)\| = \sqrt{(-\sin t)^2 + \cos^2 t + 1^2} = \sqrt{2}$$

Thus, the arc length is:

$$s = \int_0^{2\pi} \sqrt{2} dt = 2\pi\sqrt{2}$$

Example 7.4 (Arc Length of an Ellipse and Elliptic Integrals). Consider the ellipse $x^2 + \frac{y^2}{4} = 1$.

Method 1 (Cartesian Parameterization - Upper Half): Setting $x = t$ and $y =$

$2\sqrt{1-t^2}$ for $t \in [-1, 1]$ yields:

$$\vec{r}(t) = \left\langle t, 2\sqrt{1-t^2} \right\rangle \implies \vec{r}'(t) = \left\langle 1, -\frac{2t}{\sqrt{1-t^2}} \right\rangle$$

$$\|\vec{r}'(t)\| = \sqrt{1 + \frac{4t^2}{1-t^2}} = \sqrt{\frac{1+3t^2}{1-t^2}}$$

The arc length of the upper half is given by $\int_{-1}^1 \sqrt{\frac{1+3t^2}{1-t^2}} dt$.

Method 2 (Trigonometric Parameterization - Full Ellipse): Setting $x = \cos t$ and $y = 2 \sin t$ for $t \in [0, 2\pi]$:

$$\vec{r}'(t) = \langle -\sin t, 2 \cos t \rangle \implies \|\vec{r}'(t)\| = \sqrt{\sin^2 t + 4 \cos^2 t} = \sqrt{4 - 3 \sin^2 t}$$

The total perimeter of the ellipse is:

$$s = \int_0^{2\pi} \sqrt{4 - 3 \sin^2 t} dt$$

Definition 7.2: Elliptic Integral

An elliptic integral is a type of integral that cannot be expressed in terms of elementary functions.

Elliptic Integrals: It turns out these integrals cannot be evaluated in terms of elementary functions. They belong to a class of functions known as **elliptic integrals**, which require numerical methods to evaluate.

8 Scalar-Valued Functions of Several Variables

Definition 8.1: Scalar Function

A scalar-valued function of several variables maps a vector of multiple inputs to a single real output $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$.

Examples:

- Temperature distribution: $T(\text{longitude}, \text{latitude}, \text{time}) : \mathbb{R}^3 \rightarrow \mathbb{R}$.
- Precipitation depth: $P(\text{longitude}, \text{latitude}) : \mathbb{R}^2 \rightarrow \mathbb{R}$.
- Domain and Range example: For $f(x, y) = y\sqrt{x-1}$,

Domain: $\{(x, y) \in \mathbb{R}^2 : x \geq 1\}$, Range: $\mathbb{R}$

8.1 Visualizing Multivariable Functions: Traces and Level Sets

The graph of a function $f(x, y)$ is defined as the surface $\{(x, y, z) \in \mathbb{R}^3 : z = f(x, y)\}$.

Example 8.1 (Analyzing Traces of $f(x, y) = y \sin x$). To visualize $f(x, y) = y \sin x$, we examine slices along constant coordinates (traces):

y -traces ($y = c$): For $y = 0 \implies z = 0$; for $y = 1 \implies z = \sin x$; for $y = -1 \implies z = -\sin x$. (Sinusoidal curves scaled by y).


x -traces ($x = c$): For $x = 0 \implies z = 0$; for $x = \frac{\pi}{2} \implies z = y$; for $x = \pi \implies z = 0$. (Lines whose slopes depend on $\sin x$).

8.2 Level Sets and Contour Maps

For a function $f : D \rightarrow \mathbb{R}$ ($D \subseteq \mathbb{R}^2$), the **level set** (or level curve) at height c is defined as $\{(x, y) \in \mathbb{R}^2 : f(x, y) = c\}$. A collection of such level curves forms a **contour map**.

Example 8.2 (Comparing Level Curves). Consider three functions and their level sets $f(x, y) = c$:


1. $f(x, y) = \sqrt{x^2 + y^2}$: Level sets are equally spaced concentric circles $x^2 + y^2 = c^2$ (a cone).
2. $g(x, y) = x^2 + y^2$: Level sets are circles $x^2 + y^2 = c$ spaced closer together as c increases (a paraboloid).
3. $h(x, y) = (x^2 + y^2)^{1/3}$: Level sets are circles $x^2 + y^2 = c^3$ spaced further apart as c increases.




8.3 Visualizing Higher Dimensions via Slices

- A sphere in $\mathbb{R}^3$ defined by $x^2 + y^2 + z^2 = \rho^2$ can be sliced along $z = c$ to produce 2D circular cross-sections in $\mathbb{R}^2$.
- A 3-sphere in $\mathbb{R}^4$ defined by $x^2 + y^2 + z^2 + w^2 = R^2$ can be visualized through its 3D cross-sections ($w = c$), which appear as concentric 2-spheres in $\mathbb{R}^3$ whose radii expand from 0 to R and then contract back to 0.

3D Sphere Slices ($z = c$)



4D Hypersphere Slices ($w = c$ in $\mathbb{R}^3$)




9 Limits and Continuity of Functions of Several Variables

For single-variable functions, we say that the limit of a function $f(x)$ exists such that

$$\lim_{x \rightarrow a} f(x) = L \iff \forall \epsilon > 0, \exists \delta > 0 \text{ such that } 0 < |x - a| < \delta \implies |f(x) - L| < \epsilon$$

As shown below, we can visualize the ϵ - δ definition of a limit for a single-variable function $f(x) = \ln x$ at $x = a$.



Now consider the case of functions of two variables. We say that the limit of a function $f(x, y)$ exists such that

$$\begin{aligned} \lim_{(x,y) \rightarrow (a,b)} f(x, y) = L &\iff \forall \epsilon > 0, \exists \delta > 0 \text{ such that } 0 < \sqrt{(x - a)^2 + (y - b)^2} < \delta \\ &\implies |f(x, y) - L| < \epsilon \end{aligned}$$

Similarly, we can visualize the ϵ - δ definition of a limit for a two-variable function $f(x, y)$.




Figure 1: Visualization of the δ -disk in the xy -plane and its projection onto the surface $z = f(x, y)$, mapped horizontally to $f(a, b) \pm \epsilon$ on the z -axis.

Problem 9.1: Limit of a Function of Two Variables

Find, if it exists, $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$.

Solution 9.1

First, we consider approaching $(0, 0)$ along the x -axis, i.e., $y = 0$:

$$\lim_{x \rightarrow 0} \frac{x(0)}{x^2 + 0^2} = 0$$

Next, we consider approaching $(0, 0)$ along the y -axis, i.e., $x = 0$:

$$\lim_{y \rightarrow 0} \frac{0(y)}{0^2 + y^2} = 0$$

Next, we consider approaching $(0, 0)$ along the line $y = x$:

$$\lim_{x \rightarrow 0} \frac{x(x)}{x^2 + x^2} = \lim_{x \rightarrow 0} \frac{x^2}{2x^2} = \lim_{x \rightarrow 0} \frac{1}{2} = \frac{1}{2}$$

Since approaching along different paths yields different limiting values, the limit does not exist.

Systematic Path Approaches Checking paths individually can be cumbersome. Instead, we can systematically analyze general path families:

1. Linear Paths ($y = mx$):

$$\lim_{x \rightarrow 0} f(x, mx) = \lim_{x \rightarrow 0} \frac{x(mx)}{x^2 + (mx)^2} = \lim_{x \rightarrow 0} \frac{mx^2}{x^2(1 + m^2)} = \frac{m}{1 + m^2}$$

The limit depends explicitly on the slope m . Hence, the limit does not exist.

2. Polar Coordinates ($x = r \cos \theta, y = r \sin \theta$):

$$\lim_{r \rightarrow 0} f(r \cos \theta, r \sin \theta) = \lim_{r \rightarrow 0} \frac{r^2 \cos \theta \sin \theta}{r^2} = \cos \theta \sin \theta$$

The limit depends on the angle θ , confirming the limit does not exist.

Definition 9.1: Squeeze Theorem for Two Variables

If $g(x, y) \leq f(x, y) \leq h(x, y)$ for all (x, y) in a disk centered at (a, b) (except possibly at (a, b) itself), and

$$\lim_{(x,y) \rightarrow (a,b)} g(x, y) = \lim_{(x,y) \rightarrow (a,b)} h(x, y) = L$$

then $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$.

Definition 9.2: Continuity of Functions of Several Variables

A function $f(x, y)$ is *continuous* at (a, b) if:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$$

Furthermore, $f(x, y)$ is continuous on a domain D if it is continuous at every point $(a, b) \in D$.

Example 9.1 (Continuity of a Composite Function). Consider $f(x, y) = \arctan\left(\frac{y}{x}\right)$.

The inner function $g(x, y) = \frac{y}{x}$ is continuous everywhere except on the line $x = 0$. Since $h(t) = \arctan(t)$ is continuous on $\mathbb{R}$, the composite function $f(x, y) = h(g(x, y))$ is continuous at all points $(x, y) \in \mathbb{R}^2$ where $x \neq 0$.