MAT8021, Algebraic Topology

Assignment 7

Due in-class on Tuesday, April 6

- 1. Find all (2,3)-shuffles α and give formulas for the associated shuffle maps $f_{\alpha} : \Delta[5] \to \Delta[2] \times \Delta[3]$.
- 2. Find recursive formulas for $\dim_{\mathbb{Z}/2} H_k((\mathbb{RP}^2)^n; \mathbb{Z}/2)$ in terms of k and n.
- 3. Find a pair of chain complexes C_* and D_* such that the tensor product chain complex $C_* \otimes D_*$ does not satisfy the Künneth formula, i.e., there is some n such that

$$H_n(C_* \otimes D_*) \ncong \bigoplus_{p+q=n} H_p(C_*) \otimes H_q(D_*) \oplus \bigoplus_{p+q=n-1} (H_p(C_*), H_q(D_*))$$

- 4. Suppose G is a topological group and X is a topological space with a continuous map $G \times X \to X$ which is an action of G. Show that $H_*(X)$ becomes a left module over the Pontrjagin ring $H_*(G)$.
- 5. Find the homology of the complex Grassmannian $Gr_{\mathbb{C}}(3,5)$.
- 6. There is a continuous map from one Grassmannian Gr(k, n) to the next Gr(k, n+1) by sending a plane $V \subset \mathbb{R}^n$ to the plane

$$\{(0, x_1, \dots, x_n) \mid (x_1, \dots, x_n) \in V\}$$

Show that the image consists of a union of Schubert cells, and find the dimension of the smallest cell not in the image.