

# **22nd Westlake Math Colloquium | Yifei Zhu: Topology of stratified singular moduli spaces for gapless quantum mechanical systems**

**Time:** 16:00-17:00, Friday, Nov 11, 2022

**Venue:** E4-233, Yungu Campus & ZOOM

**ZOOM ID:** 863 6606 6486

**PASSCODE:** 738489


**Host:** Dr. Xing Gu, Institute for Theoretical Sciences, Westlake University

**Speaker :** Dr. Yifei Zhu, an assistant professor at the Department of Mathematics of the Southern University of Science and Technology. His research interests are in algebraic topology and related fields, particularly in its connections to algebraic geometry and number theory via objects such as formal groups, elliptic curves, and modular forms.

**Title:** Topology of stratified singular moduli spaces for gapless quantum mechanical systems

**Abstract:** This talk presents an external application of the algebraic topology of moduli spaces. In condensed matter physics, the Hamiltonian of a quantum mechanical system takes a mathematical form of a square matrix, with parameters functions on the 3D momentum space. Such a matrix satisfies the Hermitian symmetry, so that its eigenvalues are real and represent observed energies. We will discuss this space of parameters for Hamiltonians, especially its degeneracy locus where eigenvalues occur with multiplicities. Such a locus gives rise to exceptional properties in the larger scale, with applications to the design of sensing and absorbing devices. We focus on certain non-Hermitian Hamiltonians, the imaginary parts of whose eigenvalues model energy exchange of open systems. Their parameter space possesses intriguing topology, with a stratification of non-isolated singularities, which affords interesting phenomena such as the so-called bulk-edge correspondence. The associated algebraic invariants enable classifications and predictions for phases of matter. This work is in collaboration with C. T. Chan, Jing Hu, Hongwei Jia, Xiaoping Ouyang, Yixiao Wang, Yixin Xiao, Ruo-Yang Zhang, and Zhao-Qing Zhang.

# **Topology of stratified singular moduli spaces for gapless quantum mechanical systems**




Yifei Zhu (SUSTech)

Westlake Math Colloquium, November 2022

## Phases of matter: a motivating example

Ice, water, and vapor are different forms of H<sub>2</sub>O:



## Phases of matter: a motivating example


Ice, water, and vapor are different forms of H<sub>2</sub>O:



The real line of **temperatures** (in °C) minus the two **transition** points has **3 path components**. Forms of H<sub>2</sub>O connected continuously by a path are considered to be in the same **phase**.


## Phases of matter: a motivating example

Ice, water, and vapor are different forms of H<sub>2</sub>O:




The real line of **temperatures** (in °C) minus the two **transition points** has **3 path components**. Forms of H<sub>2</sub>O connected continuously by a path are considered to be in the same **phase**.

If we also allow **pressure** to vary, then there are only two phases of H<sub>2</sub>O.



## Phases of matter: a motivating example


Ice, water, and vapor are different forms of  $\text{H}_2\text{O}$ :



The real line of **temperatures** (in  $^{\circ}\text{C}$ ) minus the two **transition points** has **3 path components**. Forms of  $\text{H}_2\text{O}$  connected continuously by a path are considered to be in the same **phase**.

If we also allow **pressure** to vary, then there are only two phases of  $\text{H}_2\text{O}$ .

## A mathematical framework




There is a space  $\mathcal{M}$  of “systems” with a “singular” locus  $\Delta \subset \mathcal{M}$ , and we are interested in  $\pi_0(\mathcal{M} - \Delta)$  or, more generally, the **homotopy type** of  $\mathcal{M} - \Delta$ .

## Moduli problems: a basic example

Fix a positive integer  $n$ . Let  $\mathcal{M}_n$  be the space of configurations of  $n$  points on the real line  $\mathbb{R}$ .


## Moduli problems: a basic example

Fix a positive integer  $n$ . Let  $\mathcal{M}_n$  be the space of configurations of  $n$  points on the real line  $\mathbb{R}$ .



## Moduli problems: a basic example


Fix a positive integer  $n$ . Let  $\mathcal{M}_n$  be the space of configurations of  $n$  points on the real line  $\mathbb{R}$ .



The position of the  $i$ 'th point is a function  $x^i: \mathcal{M}_n \rightarrow \mathbb{R}$ . Together these position functions define an isomorphism  $\mathcal{M}_n \cong \mathbb{R}^n$ .

## Moduli problems: a basic example

Fix a positive integer  $n$ . Let  $\mathcal{M}_n$  be the space of configurations of  $n$  points on the real line  $\mathbb{R}$ .




The position of the  $i$ 'th point is a function  $x^i: \mathcal{M}_n \rightarrow \mathbb{R}$ . Together these position functions define an isomorphism  $\mathcal{M}_n \cong \mathbb{R}^n$ .

Here  $\mathcal{M}_n$  is a moduli space, and so far there is no interesting topology since it is contractible.

## Moduli problems: a basic example

Fix a positive integer  $n$ . Let  $\mathcal{M}_n$  be the space of configurations of  $n$  points on the real line  $\mathbb{R}$ .




The position of the  $i$ 'th point is a function  $x^i: \mathcal{M}_n \rightarrow \mathbb{R}$ . Together these position functions define an isomorphism  $\mathcal{M}_n \cong \mathbb{R}^n$ .

Here  $\mathcal{M}_n$  is a moduli space, and so far there is no interesting topology since it is contractible.

Let  $\Delta \subset \mathcal{M}_n$  be the locus of  $n$ -tuples  $x = (x^1, \dots, x^n)$  in which not all  $x^i$  are distinct. Configurations in  $\mathcal{M}_n - \Delta$  satisfy a “gap condition,” and now there is nontrivial topology:  $\mathcal{M}_n - \Delta$  has  $n!$  contractible components.

## Moduli problems: a basic example

Fix a positive integer  $n$ . Let  $\mathcal{M}_n$  be the space of configurations of  $n$  points on the real line  $\mathbb{R}$ .



The position of the  $i$ 'th point is a function  $x^i: \mathcal{M}_n \rightarrow \mathbb{R}$ . Together these position functions define an isomorphism  $\mathcal{M}_n \cong \mathbb{R}^n$ .

Here  $\mathcal{M}_n$  is a moduli space, and so far there is no interesting topology since it is contractible.

Let  $\Delta \subset \mathcal{M}_n$  be the locus of  $n$ -tuples  $x = (x^1, \dots, x^n)$  in which not all  $x^i$  are distinct. Configurations in  $\mathcal{M}_n - \Delta$  satisfy a “gap condition,” and now there is nontrivial topology:  $\mathcal{M}_n - \Delta$  has  $n!$  contractible components.

A gapped configuration  $x \in \mathcal{M}_n - \Delta$  determines a permutation  $\sigma(x) \in \text{Sym}_n$ . In fact,  $\sigma$  induces an isomorphism  $\pi_0(\mathcal{M}_n - \Delta) \cong \text{Sym}_n$  of groups.

## Moduli problems: key common features

- There are discrete parameters, e.g., the positive integer  $n$ .

## Moduli problems: key common features

- There are discrete parameters, e.g., the positive integer  $n$ .
- There is a singular locus  $\Delta$  in the moduli space  $\mathcal{M}$ :
  - away from  $\Delta$ , the parametrized objects satisfy a nonsingular (or gap) condition;

## Moduli problems: key common features

- There are discrete parameters, e.g., the positive integer  $n$ .
- There is a singular locus  $\Delta$  in the moduli space  $\mathcal{M}$ :
  - away from  $\Delta$ , the parametrized objects satisfy a nonsingular (or gap) condition;
  - moving across  $\Delta$ , gap closes and reopens, which signifies a phase transition.

## Moduli problems: key common features

- There are discrete parameters, e.g., the positive integer  $n$ .
- There is a singular locus  $\Delta$  in the moduli space  $\mathcal{M}$ :
  - away from  $\Delta$ , the parametrized objects satisfy a nonsingular (or gap) condition;
  - moving across  $\Delta$ , gap closes and reopens, which signifies a phase transition.
- There are interesting functions on  $\mathcal{M}$  which encode geometric information about the parametrized objects, e.g.,  $|x^1 - x^3|$ , the distance between the 1st and 3rd points.  
(Also think about moduli spaces of elliptic curves.)

## Moduli problems: key common features

- There are discrete parameters, e.g., the positive integer  $n$ .
- There is a singular locus  $\Delta$  in the moduli space  $\mathcal{M}$ :
  - away from  $\Delta$ , the parametrized objects satisfy a nonsingular (or gap) condition;
  - moving across  $\Delta$ , gap closes and reopens, which signifies a phase transition.
- There are interesting functions on  $\mathcal{M}$  which encode geometric information about the parametrized objects, e.g.,  $|x^1 - x^3|$ , the distance between the 1st and 3rd points.  
(Also think about moduli spaces of elliptic curves.)
- There is a complete invariant of the path component, which is an isomorphism to a known or computable set.

## Moduli problems: key common features

- There are discrete parameters, e.g., the positive integer  $n$ .
- There is a singular locus  $\Delta$  in the moduli space  $\mathcal{M}$ :
  - away from  $\Delta$ , the parametrized objects satisfy a nonsingular (or gap) condition;
  - moving across  $\Delta$ , gap closes and reopens, which signifies a phase transition.
- There are interesting functions on  $\mathcal{M}$  which encode geometric information about the parametrized objects, e.g.,  $|x^1 - x^3|$ , the distance between the 1st and 3rd points.  
(Also think about moduli spaces of elliptic curves.)
- There is a complete invariant of the path component, which is an isomorphism to a known or computable set.  
Such a complete invariant is not present in all situations.


## Why do we care about moduli spaces for physical systems?



Holography featured prominently in recent Olympic ceremonies

Holography is made possible via **exceptional** optical devices and materials.

## Why do we care about moduli spaces for physical systems?



Holography featured prominently in recent Olympic ceremonies

Holography is made possible via **exceptional** optical devices and materials.

Moduli spaces of physical systems, especially their singular loci, play a pivotal role in designing such.

## Why do we care about moduli spaces for physical systems?



Holography featured prominently in recent Olympic ceremonies

Holography is made possible via **exceptional** optical devices and materials.

Moduli spaces of physical systems, especially their singular loci, play a pivotal role in designing such.

Design materials that can “do wonders”, which cannot be found in nature, e.g., invisibility cloaks.

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

In mathematical terms, as a first example, let us consider a **2-band** Hamiltonian with **Hermitian** and **parity–time** symmetries.

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

In mathematical terms, as a first example, let us consider a **2-band** Hamiltonian with **Hermitian** and **parity-time** symmetries.

Let  $\vec{k} = (k_x, k_y, k_z)$  in the momentum space (or  $\vec{k}$ -space).

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

In mathematical terms, as a first example, let us consider a **2-band** Hamiltonian with **Hermitian** and **parity-time** symmetries.

Let  $\vec{k} = (k_x, k_y, k_z)$  in the momentum space (or  $\vec{k}$ -space). Consider the  $2 \times 2$  matrix

$$H_2(\vec{k}) = f_0(\vec{k})\mathbf{1} + \vec{f}(\vec{k}) \cdot \vec{\sigma}$$

where **1** is the identity matrix and  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  are the Pauli matrices

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \sigma_2 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

In mathematical terms, as a first example, let us consider a **2-band** Hamiltonian with **Hermitian** and **parity-time** symmetries.

Let  $\vec{k} = (k_x, k_y, k_z)$  in the momentum space (or  $\vec{k}$ -space). Consider the  $2 \times 2$  matrix

$$H_2(\vec{k}) = f_0(\vec{k})\mathbf{1} + \vec{f}(\vec{k}) \cdot \vec{\sigma}$$

where **1** is the identity matrix and  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  are the Pauli matrices

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \sigma_2 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

We have

$$H_2 = \begin{bmatrix} f_0 & 0 \\ 0 & f_0 \end{bmatrix} + \begin{bmatrix} 0 & f_1 \\ f_1 & 0 \end{bmatrix} + \begin{bmatrix} if_2 & 0 \\ 0 & -if_2 \end{bmatrix} + \begin{bmatrix} f_3 & 0 \\ 0 & -f_3 \end{bmatrix}$$

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

In mathematical terms, as a first example, let us consider a **2-band** Hamiltonian with **Hermitian** and **parity-time** symmetries.

Let  $\vec{k} = (k_x, k_y, k_z)$  in the momentum space (or  $\vec{k}$ -space). Consider the  $2 \times 2$  matrix


$$H_2(\vec{k}) = f_0(\vec{k})\mathbf{1} + \vec{f}(\vec{k}) \cdot \vec{\sigma}$$

where **1** is the identity matrix and  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  are the Pauli matrices

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \sigma_2 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

We have

$$H_2 = \begin{bmatrix} f_0 & 0 \\ 0 & f_0 \end{bmatrix} + \begin{bmatrix} 0 & f_1 \\ f_1 & 0 \end{bmatrix} + \begin{bmatrix} if_2 & 0 \\ 0 & -if_2 \end{bmatrix} + \begin{bmatrix} f_3 & 0 \\ 0 & -f_3 \end{bmatrix}$$

  
PT symmetry  
(complex conjugation)

## Moduli spaces for quantum mechanical systems

In quantum mechanics, the **Hamiltonian** of a system is an operator corresponding to the total energy of that system.

In mathematical terms, as a first example, let us consider a **2-band** Hamiltonian with **Hermitian** and **parity-time** symmetries.

Let  $\vec{k} = (k_x, k_y, k_z)$  in the momentum space (or  $\vec{k}$ -space). Consider the  $2 \times 2$  matrix


$$H_2(\vec{k}) = f_0(\vec{k})\mathbf{1} + \vec{f}(\vec{k}) \cdot \vec{\sigma}$$

where **1** is the identity matrix and  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  are the Pauli matrices

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \sigma_2 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

We have

$$H_2 = \begin{bmatrix} f_0 & 0 \\ 0 & f_0 \end{bmatrix} + \begin{bmatrix} 0 & f_1 \\ f_1 & 0 \end{bmatrix} + \begin{bmatrix} if_2 & 0 \\ 0 & -if_2 \end{bmatrix} + \begin{bmatrix} f_3 & 0 \\ 0 & -f_3 \end{bmatrix} = \begin{bmatrix} f_0 + f_3 & f_1 \\ f_1 & f_0 - f_3 \end{bmatrix}$$

  
PT symmetry  
(complex conjugation)

## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue.

## Moduli spaces for quantum mechanical systems


We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$

## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$




## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$

This gives the moduli space (not quite).  
A  $\mathbb{Z}_2$ -symmetry is present:



## Moduli spaces for quantum mechanical systems


We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$

This gives the moduli space (not quite).

A  $\mathbb{Z}_2$ -symmetry is present:

$$\begin{aligned} \omega_+ &= \sqrt{f_1^2 + f_3^2} & \vec{\phi}_+ &= \begin{bmatrix} f_3 + \sqrt{-} \\ f_1 \end{bmatrix} \\ \omega_- &= -\sqrt{-} & \vec{\phi}_- &= \begin{bmatrix} f_3 - \sqrt{-} \\ f_1 \end{bmatrix} \end{aligned}$$



## Moduli spaces for quantum mechanical systems


We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$

This gives the moduli space (not quite).

A  $\mathbb{Z}_2$ -symmetry is present:

$$\begin{array}{ll} \omega_+ = \sqrt{f_1^2 + f_3^2} & \vec{\phi}_+ = \begin{bmatrix} f_3 + \sqrt{-1}f_1 \\ f_1 \end{bmatrix} \\ \omega_- = -\sqrt{-1}f_3 & \vec{\phi}_- = \begin{bmatrix} f_3 - \sqrt{-1}f_1 \\ f_1 \end{bmatrix} \end{array} \xrightarrow{\text{normalize}} \begin{array}{ll} \omega_+ = 1 & \vec{\phi}_+ = \begin{bmatrix} \cos \theta + 1 \\ \sin \theta \end{bmatrix} \\ \omega_- = -1 & \vec{\phi}_- = \begin{bmatrix} \cos \theta - 1 \\ \sin \theta \end{bmatrix} \end{array}$$



## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$

This gives the moduli space (not quite).


A  $\mathbb{Z}_2$ -symmetry is present:

$$\begin{aligned} \omega_+ &= \sqrt{f_1^2 + f_3^2} & \vec{\phi}_+ &= \begin{bmatrix} f_3 + \sqrt{-1} \\ f_1 \end{bmatrix} & \text{normalize} \rightsquigarrow \\ \omega_- &= -\sqrt{-1} & \vec{\phi}_- &= \begin{bmatrix} f_3 - \sqrt{-1} \\ f_1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \omega_+ &= 1 \\ \omega_- &= -1 \end{aligned}$$

$$\begin{aligned} \vec{\phi}_+ &= \begin{bmatrix} \cos \theta + 1 \\ \sin \theta \end{bmatrix} \\ \vec{\phi}_- &= \begin{bmatrix} \cos \theta - 1 \\ \sin \theta \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \hat{\phi}_+ &= \begin{bmatrix} \cos(\theta/2) \\ \sin(\theta/2) \end{bmatrix} = |\phi_+\rangle \\ \hat{\phi}_- &= \begin{bmatrix} -\sin(\theta/2) \\ \cos(\theta/2) \end{bmatrix} = |\phi_-\rangle \end{aligned}$$



## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$

This gives the moduli space (not quite).

A  $\mathbb{Z}_2$ -symmetry is present:


$$\begin{aligned} \omega_+ &= \sqrt{f_1^2 + f_3^2} & \vec{\phi}_+ &= \begin{bmatrix} f_3 + \sqrt{-1} \\ f_1 \end{bmatrix} & \text{normalize} \rightsquigarrow \\ \omega_- &= -\sqrt{-1} & \vec{\phi}_- &= \begin{bmatrix} f_3 - \sqrt{-1} \\ f_1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \omega_+ &= 1 \\ \omega_- &= -1 \end{aligned}$$

$$\begin{aligned} \vec{\phi}_+ &= \begin{bmatrix} \cos \theta + 1 \\ \sin \theta \end{bmatrix} \\ \vec{\phi}_- &= \begin{bmatrix} \cos \theta - 1 \\ \sin \theta \end{bmatrix} \end{aligned} \quad \text{normalize} \rightsquigarrow$$

$$\begin{aligned} \hat{\phi}_+ &= \begin{bmatrix} \cos(\theta/2) \\ \sin(\theta/2) \end{bmatrix} = |\phi_+\rangle \\ \hat{\phi}_- &= \begin{bmatrix} -\sin(\theta/2) \\ \cos(\theta/2) \end{bmatrix} = |\phi_-\rangle \end{aligned}$$


$$\implies H_2 = \mathbf{1} - 2|\phi_-\rangle\langle\phi_-|$$



## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$



This gives the moduli space (not quite).

A  $\mathbb{Z}_2$ -symmetry is present:

$$\begin{aligned} \omega_+ &= \sqrt{f_1^2 + f_3^2} & \vec{\phi}_+ &= \begin{bmatrix} f_3 + \sqrt{-1} \\ f_1 \end{bmatrix} & \text{normalize} \rightsquigarrow \\ \omega_- &= -\sqrt{-1} & \vec{\phi}_- &= \begin{bmatrix} f_3 - \sqrt{-1} \\ f_1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \omega_+ &= 1 & \vec{\phi}_+ &= \begin{bmatrix} \cos \theta + 1 \\ \sin \theta \end{bmatrix} & \text{normalize} \rightsquigarrow \\ \omega_- &= -1 & \vec{\phi}_- &= \begin{bmatrix} \cos \theta - 1 \\ \sin \theta \end{bmatrix} & \hat{\phi}_+ = \begin{bmatrix} \cos(\theta/2) \\ \sin(\theta/2) \end{bmatrix} = |\phi_+\rangle \\ & & & & \hat{\phi}_- = \begin{bmatrix} -\sin(\theta/2) \\ \cos(\theta/2) \end{bmatrix} = |\phi_-\rangle \end{aligned}$$


$$\implies H_2 = \mathbf{1} - 2|\phi_-\rangle\langle\phi_-|$$

Observe that  $|\phi_-\rangle \mapsto -|\phi_-\rangle$  does not change  $H_2$ .

## Moduli spaces for quantum mechanical systems

We compute for which values of parameters  $H_2$  has a **doubled** eigenvalue. For this purpose, we may assume  $f_0 = 0$  and get the characteristic polynomial

$$\begin{vmatrix} f_3 - \omega & f_1 \\ f_1 & -f_3 - \omega \end{vmatrix} = \omega^2 - f_1^2 - f_3^2.$$



This gives the moduli space (not quite).

A  $\mathbb{Z}_2$ -symmetry is present:

$$\begin{aligned} \omega_+ &= \sqrt{f_1^2 + f_3^2} & \vec{\phi}_+ &= \begin{bmatrix} f_3 + \sqrt{-1} \\ f_1 \end{bmatrix} & \text{normalize} \rightsquigarrow \\ \omega_- &= -\sqrt{-1} & \vec{\phi}_- &= \begin{bmatrix} f_3 - \sqrt{-1} \\ f_1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \omega_+ &= 1 & \vec{\phi}_+ &= \begin{bmatrix} \cos \theta + 1 \\ \sin \theta \end{bmatrix} & \text{normalize} \rightsquigarrow & \hat{\phi}_+ = \begin{bmatrix} \cos(\theta/2) \\ \sin(\theta/2) \end{bmatrix} = |\phi_+\rangle \\ \omega_- &= -1 & \vec{\phi}_- &= \begin{bmatrix} \cos \theta - 1 \\ \sin \theta \end{bmatrix} & \hat{\phi}_- = \begin{bmatrix} -\sin(\theta/2) \\ \cos(\theta/2) \end{bmatrix} = |\phi_-\rangle \end{aligned}$$

$$\implies H_2 = \mathbf{1} - 2|\phi_-\rangle\langle\phi_-|$$

Observe that  $|\phi_-\rangle \mapsto -|\phi_-\rangle$  does not change  $H_2$ .

Thus the moduli space  $\mathcal{M}_2 = \mathrm{SO}(2)/\mathbb{Z}_2 \cong S^1$  and its “topological charge” (a homotopy invariant) is  $\pi_1(\mathcal{M}_2) \cong \mathbb{Z}$ .

## Moduli spaces for quantum mechanical systems

Taking

$$f_1(\vec{k}) = k_x k_z$$

$$f_3(\vec{k}) = \pm k_x^2 + k_y^2 \pm k_z^2 - 4$$


# Moduli spaces for quantum mechanical systems

Taking

$$f_1(\vec{k}) = k_x k_z$$

$$f_3(\vec{k}) = \pm k_x^2 + k_y^2 \pm k_z^2 - 4$$

we have in the  $\vec{k}$ -space




# Moduli spaces for quantum mechanical systems

Taking


$$f_1(\vec{k}) = k_x k_z$$

$$f_3(\vec{k}) = \pm k_x^2 + k_y^2 \pm k_z^2 - 4$$

we have in the  $\vec{k}$ -space



and in the moduli space




# Moduli spaces for quantum mechanical systems

Taking


$$f_1(\vec{k}) = k_x k_z$$

$$f_3(\vec{k}) = \pm k_x^2 + k_y^2 \pm k_z^2 - 4$$

we have in the  $\vec{k}$ -space




and in the moduli space



The **winding number**  $w_\Gamma$  of the loop  $\Gamma$  equals 2.


# Moduli spaces for quantum mechanical systems

The winding number serves as a topological classifier:



## Moduli spaces for quantum mechanical systems


The winding number serves as a topological classifier:



More generally, for all  $n \geq 2$ , Wu et al. computed topological charges for  $n$ -band Hermitian Hamiltonians with PT symmetry and found that  $\pi_1(\mathcal{M}_n)$  is **non-Abelian** when  $n \geq 3$ .

## Moduli spaces for quantum mechanical systems

The winding number serves as a topological classifier:




More generally, for all  $n \geq 2$ , Wu et al. computed topological charges for  $n$ -band Hermitian Hamiltonians with PT symmetry and found that  $\pi_1(\mathcal{M}_n)$  is **non-Abelian** when  $n \geq 3$ . For example,

$$\mathcal{M}_3 = \text{SO}(3)/D_2$$

where  $D_2$  = the three-dimensional “dihedral” crystallographic point group  
 $\cong$  the Klein four-group

## Moduli spaces for quantum mechanical systems

The winding number serves as a topological classifier:



More generally, for all  $n \geq 2$ , Wu et al. computed topological charges for  $n$ -band Hermitian Hamiltonians with PT symmetry and found that  $\pi_1(\mathcal{M}_n)$  is **non-Abelian** when  $n \geq 3$ . For example,


$$\mathcal{M}_3 = \text{SO}(3)/D_2$$

where  $D_2$  = the three-dimensional “dihedral” crystallographic point group  
 $\cong$  the Klein four-group

and  $\pi_1(\mathcal{M}_3) \cong Q = \{\pm 1, \pm i, \pm j, \pm k\}$ , the quaternion group.

## Moduli spaces for quantum mechanical systems

The winding number serves as a topological classifier:



More generally, for all  $n \geq 2$ , Wu et al. computed topological charges for  $n$ -band Hermitian Hamiltonians with PT symmetry and found that  $\pi_1(\mathcal{M}_n)$  is **non-Abelian** when  $n \geq 3$ . For example,

$$\mathcal{M}_3 = \text{SO}(3)/D_2$$

where  $D_2$  = the three-dimensional “dihedral” crystallographic point group  
 $\cong$  the Klein four-group

and  $\pi_1(\mathcal{M}_3) \cong Q = \{\pm 1, \pm i, \pm j, \pm k\}$ , the quaternion group.

(We have  $\pi_1(\text{SO}(3)) \cong \pi_1(\mathbb{R}P^3) \cong \mathbb{Z}_2$ ,  $\text{SU}(2) \cong S^3$  its 2-fold universal cover.)

## Moduli spaces for non-Hermitian Hamiltonians

Beyond the well-studied Hermitian symmetry, non-Hermitian Hamiltonians possess eigenvalues with **imaginary part** that represents **energy exchanges** with surrounding environment.

## Moduli spaces for non-Hermitian Hamiltonians

Beyond the well-studied Hermitian symmetry, non-Hermitian Hamiltonians possess eigenvalues with **imaginary part** that represents **energy exchanges** with surrounding environment.

Joint with the physics group at HKUST led by Che Ting Chan, especially Hongwei Jia and Jing Hu, we investigated moduli spaces for 2-band and 3-band Hamiltonians with the following symmetries:

$$\eta H \eta^{-1} = \overline{H^t} \quad \text{and} \quad [PT, H] = 0$$

pseudo-Hermiticity

parity-time symmetry

where

$$\eta = \begin{bmatrix} I_{n-1} & 0 \\ 0 & -1 \end{bmatrix} \quad \text{is a Riemannian metric form}$$

$PT$  = complex conjugation operator

## Moduli spaces for non-Hermitian Hamiltonians

Beyond the well-studied Hermitian symmetry, non-Hermitian Hamiltonians possess eigenvalues with **imaginary part** that represents **energy exchanges** with surrounding environment.

Joint with the physics group at HKUST led by Che Ting Chan, especially Hongwei Jia and Jing Hu, we investigated moduli spaces for 2-band and 3-band Hamiltonians with the following symmetries:

$$\eta H \eta^{-1} = \overline{H^t} \quad \text{and} \quad [PT, H] = 0$$

pseudo-Hermiticity parity-time symmetry

where

$$\eta = \begin{bmatrix} I_{n-1} & 0 \\ 0 & -1 \end{bmatrix} \quad \text{is a Riemannian metric form}$$

$PT$  = complex conjugation operator

Here emerge **non-isolated, stratified** singular loci, making our systems **gapless** and their topology much intriguing.


## Moduli spaces for non-Hermitian Hamiltonians: 2-band systems

In the generic 2-band case, we give complete invariants.

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad (\text{Recall Hermitian } \begin{bmatrix} f_3 & f_1 \\ f_1 & -f_3 \end{bmatrix})$$

## Moduli spaces for non-Hermitian Hamiltonians: 2-band systems

In the generic 2-band case, we give **complete invariants**.




$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad (\text{Recall Hermitian } \begin{bmatrix} f_3 & f_1 \\ f_1 & -f_3 \end{bmatrix})$$

I, III: real eigenstates

II, IV: complex eigenstates

## Moduli spaces for non-Hermitian Hamiltonians: 2-band systems

In the generic 2-band case, we give **complete invariants**.



$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad (\text{Recall Hermitian } \begin{bmatrix} f_3 & f_1 \\ f_1 & -f_3 \end{bmatrix})$$

I, III: real eigenstates


II, IV: complex eigenstates

**Stratified singular locus:**

- EL (exceptional line): doubled eigenstate (defective degeneracies)
- CIP (complete intersection point):  $\eta$ -orthogonal eigenstates (non-defective degeneracy)

# Moduli spaces for non-Hermitian Hamiltonians: 2-band systems

In the generic 2-band case, we give **complete invariants**.




$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad (\text{Recall Hermitian } \begin{bmatrix} f_3 & f_1 \\ f_1 & -f_3 \end{bmatrix})$$

I, III: real eigenstates


**Stratified singular locus:**

- EL (exceptional line): doubled eigenstate (defective degeneracies)
- CIP (complete intersection point):  $\eta$ -orthogonal eigenstates (non-defective degeneracy)



# Moduli spaces for non-Hermitian Hamiltonians: 2-band systems

In the generic 2-band case, we give **complete invariants**.




$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad (\text{Recall Hermitian } \begin{bmatrix} f_3 & f_1 \\ f_1 & -f_3 \end{bmatrix})$$

I, III: real eigenstates


**Stratified singular locus:**

- EL (exceptional line): doubled eigenstate (defective degeneracies)
- CIP (complete intersection point):  $\eta$ -orthogonal eigenstates (non-defective degeneracy)



# Moduli spaces for non-Hermitian Hamiltonians: 2-band systems

In the generic 2-band case, we give **complete invariants**.




$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad (\text{Recall Hermitian } \begin{bmatrix} f_3 & f_1 \\ f_1 & -f_3 \end{bmatrix})$$

I, III: real eigenstates

**Stratified singular locus:**



- EL (exceptional line): doubled eigenstate (defective degeneracies)
- CIP (complete intersection point):  $\eta$ -orthogonal eigenstates (non-defective degeneracy)



The moduli space  $\mathcal{M}_2$  can be identified as  $S^1 \vee S^1 \vee S^1$  doubly covering  $S^1 \vee S^1$ . Thus  $\pi_1(\mathcal{M}_2)$  is a free subgroup of  $F(a, \beta)$  on 3 generators. This gives the **gapless** system a **physically meaningful**, non-Abelian topological charge.


## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

The 3-band case is more complex and exotic. Here is a sample of portraits for the stratified singular moduli spaces.





$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$



$$\begin{bmatrix} f_1 f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$







$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system


## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

The 3-band case is more complex and exotic. Here is a sample of portraits for the stratified singular moduli spaces.



$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix} \quad \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

The central figure within these configurations is the so-called **swallowtail catastrophe**.



$$\begin{bmatrix} f_1 f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$


$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system

## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

The singularity of a swallowtail arises in the **discriminant** surface of a generic **degree-4 polynomial**. Here, it is the characteristic polynomial of  $H(f_1, f_2, f_3)$ .

$$\begin{vmatrix} -\omega & -f_1 & -f_2 & -f_3 \\ 1 & -\omega & 0 & 0 \\ 0 & 1 & -\omega & 0 \\ 0 & 0 & 1 & -\omega \end{vmatrix} = \omega^4 + f_1 \omega^2 + f_2 \omega + f_3$$



$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

The singularity of a swallowtail arises in the **discriminant** surface of a generic **degree-4 polynomial**. Here, it is the characteristic polynomial of  $H(f_1, f_2, f_3)$ .

$$\begin{vmatrix} -\omega & -f_1 & -f_2 & -f_3 \\ 1 & -\omega & 0 & 0 \\ 0 & 1 & -\omega & 0 \\ 0 & 0 & 1 & -\omega \end{vmatrix} = \omega^4 + f_1 \omega^2 + f_2 \omega + f_3$$

$$\Delta(f_1, f_2, f_3) = 4f_1^3f_2^2 + 27f_2^4 - 16f_1^4f_3 - 144f_1f_2^2f_3 + 128f_1^2f_3^2 - 256f_3^3$$

<https://yifeizhu.github.io/swtl.mp4>



$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

The singularity of a swallowtail arises in the **discriminant** surface of a generic **degree-4 polynomial**. Here, it is the characteristic polynomial of  $H(f_1, f_2, f_3)$ .


$$\begin{vmatrix} -\omega & -f_1 & -f_2 & -f_3 \\ 1 & -\omega & 0 & 0 \\ 0 & 1 & -\omega & 0 \\ 0 & 0 & 1 & -\omega \end{vmatrix} = \omega^4 + f_1 \omega^2 + f_2 \omega + f_3$$

$$\Delta(f_1, f_2, f_3) = 4f_1^3f_2^2 + 27f_2^4 - 16f_1^4f_3 - 144f_1f_2^2f_3 + 128f_1^2f_3^2 - 256f_3^3$$


<https://yifeizhu.github.io/swtl.mp4>




Discriminant surfaces are ruled (in fact, developable).



1908 quadratic ( $\Delta = b^2 - 4ac$ )



cubic




$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Illustration from Klein's lecture notes by Hellinger.

## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

As the singular loci of moduli spaces for polynomials, swallowtail and other catastrophes are important and well-studied objects in [dynamical systems](#) and [algebraic geometry](#). Arnold famously related their [complements](#) to [braid groups](#) and computed their cohomology, establishing a connection to [topology](#) as well.




V. I. Arnold's tombstone at the Novodevichy Cemetery in Moscow

## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

As the singular loci of moduli spaces for polynomials, swallowtail and other catastrophes are important and well-studied objects in [dynamical systems](#) and [algebraic geometry](#). Arnold famously related their [complements](#) to [braid groups](#) and computed their cohomology, establishing a connection to [topology](#) as well.

However, [moduli spaces for Hamiltonians](#) carry additional structures and are more complex:

- Physicists desire classifications for the behavior of [eigenstates](#) along a loop across/encircling the stratified non-isolated singularity (e.g., Berry phase of adiabatic transformation, close and open of gaps).
- Over [the reals](#), we know less even on the mathematical side.



Reg I and Reg II: PT-exact phases  
Reg III: PT-broken phase



V. I. Arnold's tombstone at the Novodevichy Cemetery in Moscow

## **Moduli spaces for non-Hermitian Hamiltonians: 3-band systems**

Despite the delicate physical meanings of this moduli space for Hamiltonians, the question of the desired classification appears to be purely mathematical.

## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

Despite the delicate physical meanings of this moduli space for Hamiltonians, the question of the desired classification appears to be purely mathematical.

We obtained results for **intersection homotopy classes** of loops near the swallowtail, i.e., deformations of loops conserving **topological charges**, with electric-circuit experimental realizations.

## Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

Despite the delicate physical meanings of this moduli space for Hamiltonians, the question of the desired classification appears to be purely mathematical.

We obtained results for **intersection homotopy classes** of loops near the swallowtail, i.e., deformations of loops conserving **topological charges**, with electric-circuit experimental realizations.




Example: transition from two EL3s  
to NIL and NL

<https://yifeizhu.github.io/loop.mp4>

# Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

Despite the delicate physical meanings of this moduli space for Hamiltonians, the question of the desired classification appears to be purely mathematical.

We obtained results for **intersection homotopy classes** of loops near the swallowtail, i.e., deformations of loops conserving **topological charges**, with electric circuit realizations.



Example: transition from two EL3s to NIL and NL

<https://yifeizhu.github.io/loop.mp4>

EL3 = triple exceptional line

NIL = nodal intersection line


NL = nodal line

a1, b1: loops on the spherical cross-section of the swallowtail

# Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

Despite the delicate physical meanings of this moduli space for Hamiltonians, the question of the desired classification appears to be purely mathematical.

We obtained results for **intersection homotopy classes** of loops near the swallowtail, i.e., deformations of loops conserving **topological charges**, with electric circuit realizations.



Example: transition from two EL3s to NIL and NL

<https://yifeizhu.github.io/loop.mp4>


EL3 = triple exceptional line  
NIL = nodal intersection line  
NL = nodal line

a1, b1: loops on the spherical cross-section of the swallowtail  
a2, b2: trajectories of **eigenvalues**

# Moduli spaces for non-Hermitian Hamiltonians: 3-band systems

Despite the delicate physical meanings of this moduli space for Hamiltonians, the question of the desired classification appears to be purely mathematical.

We obtained results for **intersection homotopy classes** of loops near the swallowtail, i.e., deformations of loops conserving **topological charges**, with electric circuit realizations.



Example: transition from two EL3s to NIL and NL

<https://yifeizhu.github.io/loop.mp4>

EL3 = triple exceptional line  
NIL = nodal intersection line  
NL = nodal line

a1, b1: loops on the spherical cross-section of the swallowtail

a2, b2: trajectories of **eigenvalues**

a3, b3: trajectories of **eigenframe**, experimental (above) and theoretical (below)

## Implications and ramifications

- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.

## Implications and ramifications

- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.
- We have been experimentally investigating the **bulk–edge correspondence** for hypersurface singularities stemmed from our theoretical analysis with the swallowtail moduli space,

## Implications and ramifications


- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.
- We have been experimentally investigating the **bulk–edge correspondence** for hypersurface singularities stemmed from our theoretical analysis with the swallowtail moduli space, i.e., the topology of **bulk states (moduli space)** determines the numerology of **edge states (parametrized system)**.

## Implications and ramifications

- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.
- We have been experimentally investigating the **bulk–edge correspondence** for hypersurface singularities stemmed from our theoretical analysis with the swallowtail moduli space, i.e., the topology of **bulk states (moduli space)** determines the numerology of **edge states (parametrized system)**. There has not been a rigorous mathematical explanation for such a correspondence, but it is reminiscent of the **geometric Langlands duality**.

## Implications and ramifications


- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.
- We have been experimentally investigating the **bulk–edge correspondence** for hypersurface singularities stemmed from our theoretical analysis with the swallowtail moduli space, i.e., the topology of **bulk states (moduli space)** determines the numerology of **edge states (parametrized system)**. There has not been a rigorous mathematical explanation for such a correspondence, but it is reminiscent of the **geometric Langlands duality**.
- Along a resolution of singularities, homotopy type of the (complex) swallowtail complement has a precise relationship to the **homotopy groups of spheres** in specific dimensions.



The unfurled swallowtail over the ordinary swallowtail

## Implications and ramifications


- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.
- We have been experimentally investigating the **bulk–edge correspondence** for hypersurface singularities stemmed from our theoretical analysis with the swallowtail moduli space, i.e., the topology of **bulk states (moduli space)** determines the numerology of **edge states (parametrized system)**. There has not been a rigorous mathematical explanation for such a correspondence, but it is reminiscent of the **geometric Langlands duality**.
- Along a resolution of singularities, homotopy type of the (complex) swallowtail complement has a precise relationship to the **homotopy groups of spheres** in specific dimensions. The rich geometric (and physical) structure of this stratified space can be exploited to introduce additional structures in  $\{\pi_i(S^{d-\mu-2})\}$ .





The unfurled swallowtail over the ordinary swallowtail



## Implications and ramifications


- Our  $S^1 \vee S^1 \vee S^1$  classification for 2-band systems predicts a new kind of non-Hermitian gapless phase of matter, with **topologically protected edge states**.
- We have been experimentally investigating the **bulk-edge correspondence** for hypersurface singularities stemmed from our theoretical analysis with the swallowtail moduli space, i.e., the topology of **bulk states (moduli space)** determines the numerology of **edge states (parametrized system)**. There has not been a rigorous mathematical explanation for such a correspondence, but it is reminiscent of the **geometric Langlands duality**.
- Along a resolution of singularities, homotopy type of the (complex) swallowtail complement has a precise relationship to the **homotopy groups of spheres** in specific dimensions. The rich geometric (and physical) structure of this stratified space can be exploited to introduce additional structures in  $\{\pi_i(S^{d-\mu-2})\}$ .
- We have been zooming in to **local details** of combinations of swallowtails (and other basic types of singularities) in order to pass from local invariants to **global (and complete) invariants** via fuller power of algebraic topology.



The unfurled swallowtail over the ordinary swallowtail







## Credits and references

- Daniel S. Freed, *Lectures on field theory and topology*, American Mathematical Society, 2019
- Phase diagram of water (simplified) from Wikipedia (author of the original work: Cmglee)
- Beijing Winter Olympics picture: <http://en.kremlin.ru/events/president/news/67715>
- Quadric surfaces diagrams for  $\vec{k}$ -spaces of Hermitian systems and winding number comparison diagrams adapted from  
QuanSheng Wu, Alexey A. Soluyanov, and Tomáš Bzdušek, *Non-Abelian band topology in noninteracting metals*, Science, 2019
- Order parameter space diagrams for non-Hermitian systems from  
Hongwei Jia, Ruo-Yang Zhang, Jing Hu, Yixin Xiao, Yifei Zhu, and C. T. Chan, *Topological classification for intersection singularities of exceptional surfaces in pseudo-Hermitian systems*, preprint, 2022
- 3D plots for 3-band-system moduli spaces drawn with Mathematica, middle-right pair by Hongwei Jia
- Swallowtail 3D video: Oliver Labs, <https://yifeizhu.github.io/swtl.mp4>
- Sketch of ruled swallowtail surface from Felix Klein's lecture notes by Ernst David Hellinger, 1907–1909.
- Photo of Arnold's tombstone from  
Boris A. Khesin and Serge L. Tabachnikov, ed., *Arnold: Swimming against the tide*, American Mathematical Society, 2014
- Spherical swallowtail plot, loop transition video and diagrams from  
Jing Hu, Ruo-Yang Zhang, Yixiao Wang, Xiaoping Ouyang, Yifei Zhu, Hongwei Jia, and C. T. Chan, *Non-Hermitian swallowtail catastrophe revealing transitions across diverse topological singularities*, preprint, 2022
- Unfurled swallowtail plot from  
V. I. Arnold, *Singularities of caustics and wave fronts*, Springer, 1990
- Swallowtail ensemble and detail plots drawn with Mathematica by Pingya Feng