

Moduli, moduli, moduli:

Portraits of moduli spaces



Yifei Zhu


Southern University of Science and Technology

2023.7.25

What is a moduli space?

A moduli space is a space of parameters, that is, a set of parameters with extra structure. These parameters label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a space of parameters, that is, a set of parameters with extra structure. These parameters label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a **space** of parameters, that is, a **set** of parameters **with extra structure**. These parameters label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a **space** of parameters, that is, a **set** of parameters **with extra structure**. These **parameters** label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a **space** of parameters, that is, a **set** of parameters **with extra structure**. These **parameters** label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a **space** of parameters, that is, a **set** of parameters **with extra structure**. These **parameters** label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a **space** of parameters, that is, a **set** of parameters **with extra structure**. These **parameters** label objects we would like to study, often in a continuous fashion.


Mental picture:



What is a moduli space?

A **moduli space** is a **space** of parameters, that is, a **set** of parameters **with extra structure**. These **parameters** label objects we would like to study, often in a continuous fashion.

Mental picture:



Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous family of objects, or how an object varies as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object varies as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own **structure**, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually **and as a whole**.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the second-order nature.

Why would you care about moduli spaces?

- Instead of dealing with an isolated or static object, we would really like to understand a continuous **family** of objects, or how an object **varies** as the parameters on which it depends change.
- For these purposes, it is often fruitful to study this collection of parameters as a space on its own right.
- While the objects in question have structure, the parameter space also has its own structure, often rich and distinct from that of the objects.
- Understanding the moduli space offers in turn understanding of the objects individually and as a whole.
- In this sense, studying moduli spaces is of the **second-order** nature.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a stratification of the moduli stack of formal groups by heights and primes, chromatic homotopy theory organizes generalized cohomology theories according to their capabilities to detect periodic families of elements in the homotopy groups of spheres.
 - In the opposite direction, homotopical methods, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a stratification of the moduli stack of formal groups by heights and primes, chromatic homotopy theory organizes generalized cohomology theories according to their capabilities to detect periodic families of elements in the homotopy groups of spheres.
 - In the opposite direction, homotopical methods, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a stratification of the moduli stack of formal groups by heights and primes, chromatic homotopy theory organizes generalized cohomology theories according to their capabilities to detect periodic families of elements in the homotopy groups of spheres.
 - In the opposite direction, homotopical methods, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, chromatic homotopy theory organizes generalized cohomology theories according to their capabilities to detect periodic families of elements in the homotopy groups of spheres.
 - In the opposite direction, homotopical methods, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, chromatic homotopy theory organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, homotopical methods, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, homotopical methods, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **homotopy theory** **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **homotopy theory** **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of **infinity categories**, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **homotopy theory** **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **homotopy theory**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with chromatic homotopy groups of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with **chromatic homotopy groups** of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with **chromatic homotopy groups** of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed a sheaf further binding arithmetic and topological data.

First portraits: Context and motivations

- **p -adic arithmetic geometry**  **topology**
 - In one direction, through a **stratification of the moduli stack of formal groups by heights and primes**, **chromatic homotopy theory** organizes generalized cohomology theories according to their capabilities to detect **periodic families of elements in the homotopy groups of spheres**.
 - In the opposite direction, **homotopical methods**, especially the theory of infinity categories, have enabled constructions previously ad hoc or impossible in algebra and number theory, leading to recent advances in p -adic Hodge theory and algebraic K-theory.
- [Joint with Guozhen Wang et al.] Building upon both computational and conceptual progresses with **chromatic homotopy groups** of spaces and cohomology operations in connection with explicit arithmetic moduli of algebro-geometric objects, we have constructed **a sheaf further binding arithmetic and topological data**.

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\text{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- **Objects to parametrize:** Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\text{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\text{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\text{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\text{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\text{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

$$\begin{array}{ccccccc}
\mathbb{G} & \xleftarrow{\quad} & \pi^*\mathbb{G} & \xrightarrow[\cong]{\alpha} & i^*G & \xrightarrow{\quad} & G \\
\downarrow & & \downarrow \llcorner & & \downarrow \lrcorner & & \downarrow \\
\mathrm{Spf} R & \xleftarrow{\pi} & \mathrm{Spec} R/\mathfrak{m} & \xlongequal{\quad} & \mathrm{Spec} R/\mathfrak{m} & \xrightarrow{i} & \mathrm{Spec} k
\end{array}$$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\mathrm{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

$$\begin{array}{ccccccc}
\mathbb{G} & \xleftarrow{\quad} & \pi^*\mathbb{G} & \xrightarrow[\cong]{\alpha} & i^*G & \xrightarrow{\quad} & G \\
\downarrow & & \downarrow \llcorner & & \downarrow \lrcorner & & \downarrow \\
\mathrm{Spf} R & \xleftarrow{\pi} & \mathrm{Spec} R/\mathfrak{m} & \xlongequal{\quad} & \mathrm{Spec} R/\mathfrak{m} & \xrightarrow{i} & \mathrm{Spec} k
\end{array}$$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\mathrm{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebro-geometric setup

- Objects to parametrize: Deformations of a formal group with level structure

G = formal group of height $h < \infty$ over a perfect field k of characteristic $p > 0$

R = complete local ring with residue field k , nilpotent maximal ideal \mathfrak{m} , and natural projection $\pi: R \rightarrow R/\mathfrak{m}$

deformation of G/k to $R := (\mathbb{G}, i, \alpha)$ with \mathbb{G} a formal group over R , $i: k \hookrightarrow R/\mathfrak{m}$, and $\alpha: \pi^*\mathbb{G} \xrightarrow{\cong} i^*G$

$$\begin{array}{ccccccc}
\mathbb{G} & \xleftarrow{\quad} & \pi^*\mathbb{G} & \xrightarrow[\cong]{\alpha} & i^*G & \xrightarrow{\quad} & G \\
\downarrow & & \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow \\
\mathrm{Spf} R & \xleftarrow{\pi} & \mathrm{Spec} R/\mathfrak{m} & \xlongequal{\quad} & \mathrm{Spec} R/\mathfrak{m} & \xrightarrow{i} & \mathrm{Spec} k
\end{array}$$

deformation of G with a level- $\Gamma_0(p^n)$ structure := (\mathbb{G}, \mathbb{H}) with \mathbb{H} a cyclic degree- p^n subgroup

$= \psi: \mathbb{G} \rightarrow \mathbb{G}/\mathbb{H}$ over an extension of R
which lifts the relative Frobenius $\mathrm{Frob}^n: G \rightarrow G^{(p^n)}$

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are topologically realized by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are topologically realized by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are topologically realized by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are topologically realized by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are topologically realized by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are topologically realized by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are **topologically realized** by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all levels literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are **topologically realized** by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all **levels** literally:

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are **topologically realized** by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all **levels** literally:

$$\begin{array}{c} \mathcal{LT}_n \\ \downarrow \\ \mathcal{LT}_0 = \coprod_{\mathbb{Z}} \mathbb{B}^{h-1} \end{array}$$

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are **topologically realized** by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all **levels** literally:

$$\begin{array}{ccc} \mathcal{LT}_n & & \\ \downarrow & & \\ \mathcal{LT}_0 & = & \coprod_{\mathbb{Z}} \mathbb{B}^{h-1} \\ \text{Gross–Hopkins} & & \\ \text{period} & & \\ \downarrow & & \\ & & \mathbb{P}^{h-1} \end{array}$$

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures

Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are **topologically realized** by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all **levels** literally:

$$\begin{array}{c} \mathcal{LT}_\infty \\ \downarrow \\ \mathcal{LT}_n \\ \downarrow \\ \mathcal{LT}_0 = \coprod_{\mathbb{Z}} \mathbb{B}^{h-1} \\ \searrow^{\text{Gross–Hopkins period}} \\ \mathbb{P}^{h-1} \end{array}$$

First portraits: Algebraic and topological moduli spaces

- Moduli spaces for deformations of a formal group with level structures


Fix a formal group G/k . Its level- $\Gamma_0(p^n)$ deformations are classified:

- [Lubin–Tate '66] for $n = 0$, by an affine formal scheme $\mathrm{Spf} A_0$ of dimension $h - 1$ over the Witt ring \mathcal{O}_F of k (with $F = \max.$ unrf. ext. of p -adic comp. of k).
- [Strickland '97] for each $n > 0$, by an affine formal scheme $\mathrm{Spf} A_n$ of finite rank over A_0 .

[Goerss–Hopkins–Miller, Strickland '98] These moduli spaces are **topologically realized** by the Morava E-theory spectrum $E = E_h(G/k)$, i.e.,

$$A_n \cong E^0(B\Sigma_{p^n})/I_{\mathrm{tr}}$$

[Fargues '08, Weinstein '15] As a family of spaces indexed by n , they stack into a tower with all **levels** literally:



First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of rings of E -power operations. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of rings of E -power operations. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of rings of E -power operations. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of rings of E -power operations. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of rings of E -power operations. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of rings of E -power operations. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of **rings of E -power operations**. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A sheaf over the Lubin–Tate tower

- Koszul complexes for computing unstable chromatic homotopy of spheres

[Devinatz–Hopkins '04, Wang '15] There is a homology-to-homotopy SS converging to the v_n -periodic homotopy groups of the q -dimensional sphere

$$H_c^*(\mathcal{G}_h; E_*^\wedge \Phi_h(S^q)) \implies v_n^{-1} \pi_* S^q$$

whose E_2 -page is the continuous group cohomology of the h 'th Morava stabilizer group \mathcal{G}_h with coefficients in the completed E -homology of the Bousfield–Kuhn functor Φ_h applied to the q -sphere.

This can be viewed as a homotopy-fixed-point SS with symmetry group \mathcal{G}_h .

[Behrens–Rezk '20, Rezk, Zhu '17] To compute its E_2 -page, the \mathcal{G}_h -modules of E -homology groups can be computed from a certain Koszul complex of **rings of E -power operations**. Moreover, these rings can be derived from A_n in the Lubin–Tate tower, so that with this modular interpretation, the Koszul complex has explicit formulas.

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Filtration (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^\wedge \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\bar{\mathbb{F}}_p)[[v_1]] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\bar{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array} \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I'$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^\wedge \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[[v_1]] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array} \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I'$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^{\wedge} \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\bar{\mathbb{F}}_p)[[v_1]] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\bar{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{c} \Omega S^1 \\ \downarrow \\ \Omega^3 S^3 \\ \downarrow \\ \Omega^5 S^5 \\ \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^* \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[[v_1]] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{aligned} E^0 &\xrightarrow{\psi} E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 &\mapsto v'_1 \quad \alpha_1 \mapsto \alpha'_1 \end{aligned}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^{\wedge} \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^{\wedge} \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^{\wedge} \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^* \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^* \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^{\wedge} \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{array}{ccc} E^0 & \xrightarrow{\psi} & E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 \mapsto v'_1 & & \alpha_1 \mapsto \alpha'_1 \end{array}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \vdots & & & & & & \\ A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 & & \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p & & \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Filtration** (in the case of $h = 2$ for simplicity)
 - Have a sequence of unstable spheres
 - Applying $E_0^* \Phi_2(-)$, get a sequence of Koszul complexes

$$A_0 = W(\overline{\mathbb{F}}_p)[\![v_1]\!] \cong E^0(\text{pt})$$

$$\begin{aligned} A_1 &= W(\overline{\mathbb{F}}_p)[\alpha_1, \alpha'_1]/(\alpha_1 \alpha'_1 - p) \\ &\cong E^0(B\Sigma_p)/I_{\text{tr}} \end{aligned}$$

$$\begin{aligned} E^0 &\xrightarrow{\psi} E^0(B\Sigma_p)/I \xrightarrow{\psi} E^0(B(\Sigma_p \wr \Sigma_p))/I' \\ v_1 &\mapsto v'_1 \quad \alpha_1 \mapsto \alpha'_1 \end{aligned}$$

$$\begin{array}{ccccccc} \Omega S^1 & & & & & & \\ \downarrow & & & & & & \\ \Omega^3 S^3 & & & & & & \\ \downarrow & & & & & & \\ \Omega^5 S^5 & & & & & & \\ \downarrow & & & & & & \\ \vdots & & & & & & \\ \end{array} \quad \begin{array}{ccccc} A_0 & \xrightarrow{-1} & A_1 & \xrightarrow{1} & A_1/A_0 \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p \\ A_0 & \xrightarrow{-\alpha_1} & A_1 & \xrightarrow{\alpha'_1} & A_1/A_0 \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p \\ A_0 & \xrightarrow{-\alpha_1^2} & A_1 & \xrightarrow{\alpha'^2_1} & A_1/A_0 \\ \downarrow \text{id} & & \downarrow \alpha_1 & & \downarrow p \\ \vdots & & \vdots & & \vdots \end{array}$$

- More concretely, this forms a sheaf of functions over the moduli, filtered by orders m of pole at the irreducible component cut out by α_1 , which precisely correspond to dimensions $q = 2m + 1$ of the spheres.

| c_4 | c_6 | c_4^2 | $c_4 c_6$ | Δ, c_4^3 | $c_4^2 c_6$ | $c_4 \Delta, c_4^4$ | $c_6 \Delta, c_4^3 c_6$ |
|-----------|--------------------------|-----------------|--------------------------|--------------------------|-----------------|---------------------|-------------------------|
| $a_1 a_3$ | $9a_3^2$ | $a_1 a_3 c_4$ | $9a_3^2 c_4$ | $a_1 a_3 c_4^2$ | $9a_3^2 c_4^2$ | $a_1 a_3 c_4^3$ | $9a_3^2 c_4^3$ |
| x_0^2 | $3a_3^2$ | $a_1^2 a_3^2$ | $3a_3^2 c_4$ | $a_1^2 a_3^2 c_4$ | $3a_3^2 c_4^2$ | $a_1^2 a_3^2 c_4^2$ | $3a_3^2 c_4^3$ |
| a_3^2 | $a_2 x_0^3 - 2a_4 x_0^2$ | $a_3^2 c_4$ | $27a_3^4 \sim a_3^2 c_6$ | $a_3^2 c_4^2$ | $a_3^2 c_6 c_4$ | $a_3^2 c_4^3$ | |
| | x_0^4 | $a_1 a_3^3 (?)$ | $9a_3^4$ | $a_1 a_3^3 c_4$ | $9a_3^4 c_4$ | $a_1 a_3^3 c_4^2$ | |
| | | x_0^5 | $3a_3^4$ | $a_1^2 a_3^4$ | $3a_3^4 c_4$ | $a_1^2 a_3^4 c_4$ | |
| | | | a_3^4 | $a_2 x_0^6 - 5a_4 x_0^5$ | $a_3^4 c_4$ | $a_3^4 c_6$ | |

Part of calculations at $p = 3$

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Symmetries**
 - The sequence of Koszul complexes is equivariant with respect to the action of the Morava stabilizer group $\mathcal{G}_h \cong \mathcal{O}_D^\times$ (with $D / F = \text{cent. div. alg. of inv. } 1/h$).
 - Both $GL_h(F)$ (its congruence subgroups K_n) and D^\times act on the Lubin–Tate tower, realizing the Jacques–Langlands correspondence.
 - [Faltings, Fargues '08] There is an equivariant isomorphism between the Lubin–Tate tower and another Drinfeld tower (parametrizing shtukas).

First portraits: A filtered, equivariant, quasi-syntomic sheaf


- **Symmetries**
 - The sequence of Koszul complexes is equivariant with respect to the action of the Morava stabilizer group $\mathcal{G}_h \cong \mathcal{O}_D^\times$ (with $D / F = \text{cent. div. alg. of inv. } 1/h$).
 - Both $GL_h(F)$ (its congruence subgroups K_n) and D^\times act on the Lubin–Tate tower, realizing the Jacques–Langlands correspondence.
 - [Faltings, Fargues '08] There is an equivariant isomorphism between the Lubin–Tate tower and another Drinfeld tower (parametrizing shtukas).

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Symmetries**
 - The sequence of Koszul complexes is equivariant with respect to the action of the Morava stabilizer group $\mathcal{G}_h \cong \mathcal{O}_D^\times$ (with $D / F =$ cent. div. alg. of inv. $1/h$).
 - Both $GL_h(F)$ (its congruence subgroups K_n) and D^\times act on the Lubin–Tate tower, realizing the Jacques–Langlands correspondence.
 - [Faltings, Fargues ’08] There is an equivariant isomorphism between the Lubin–Tate tower and another Drinfeld tower (parametrizing shtukas).

First portraits: A filtered, equivariant, quasi-syntomic sheaf


- **Symmetries**
 - The sequence of Koszul complexes is equivariant with respect to the action of the Morava stabilizer group $\mathcal{G}_h \cong \mathcal{O}_D^\times$ (with $D / F = \text{cent. div. alg. of inv. } 1/h$).
 - Both $\text{GL}_h(F)$ (its congruence subgroups K_n) and D^\times act on the Lubin–Tate tower, realizing the **Jacques–Langlands correspondence**.
 - [Faltings, Fargues '08] There is an equivariant isomorphism between the Lubin–Tate tower and another Drinfeld tower (parametrizing shtukas).



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Symmetries**

- The sequence of Koszul complexes is equivariant with respect to the action of the Morava stabilizer group $\mathcal{G}_h \cong \mathcal{O}_D^\times$ (with $D / F = \text{cent. div. alg. of inv. } 1/h$).
- Both $\text{GL}_h(F)$ (its congruence subgroups K_n) and D^\times act on the Lubin–Tate tower, realizing the **Jacques–Langlands correspondence**.
- [Faltings, Fargues '08] There is an equivariant isomorphism between the Lubin–Tate tower and another Drinfeld tower (parametrizing shtukas).




First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Symmetries**

- The sequence of Koszul complexes is equivariant with respect to the action of the Morava stabilizer group $\mathcal{G}_h \cong \mathcal{O}_D^\times$ (with $D / F = \text{cent. div. alg. of inv. } 1/h$).
- Both $\text{GL}_h(F)$ (its congruence subgroups K_n) and D^\times act on the Lubin–Tate tower, realizing the **Jacques–Langlands correspondence**.
- [Faltings, Fargues ’08] There is an equivariant isomorphism between the Lubin–Tate tower and another Drinfeld tower (parametrizing shtukas).

$$\begin{array}{ccccccc} & \cdots & A_0 & \xleftarrow{\text{id}} & A_0 & \xleftarrow{\text{id}} & A_0 \\ & \downarrow & \downarrow -b^2 & & \downarrow & & \downarrow -1 \\ & \cdots & A_1 & \xleftarrow{b} & A_1 & \xleftarrow{b} & A_1 \\ & \downarrow & \downarrow b' & & \downarrow & & \downarrow 1 \\ & \cdots & A_1/A_0 & \xleftarrow{p} & A_1/A_0 & \xleftarrow{p} & A_1/A_0 \end{array}$$



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Topologies
 - [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of quasi-syntomic rings and sheaves on the category of quasi-regular semi-perfectoid rings, valued in any presentable ∞ -category.
 - These correspond to moduli spaces at finite levels and at the infinite level, respectively.

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Topologies
 - [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of **quasi-syntomic rings** and sheaves on the category of quasi-regular semi-perfectoid rings, valued in any presentable ∞ -category.
 - These correspond to moduli spaces at finite levels and at the infinite level, respectively.

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Topologies
 - [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of **quasi-syntomic rings** and sheaves on the category of **quasi-regular semi-perfectoid rings**, valued in any presentable ∞ -category.
 - These correspond to moduli spaces at finite levels and at the infinite level, respectively.

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Topologies
 - [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of **quasi-syntomic rings** and sheaves on the category of **quasi-regular semi-perfectoid rings**, valued in any presentable ∞ -category.
 - These correspond to moduli spaces at finite levels and at the infinite level, respectively.


First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Topologies
 - [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of **quasi-syntomic rings** and sheaves on the category of **quasi-regular semi-perfectoid rings**, valued in any presentable ∞ -category.
 - These correspond to moduli spaces at **finite levels** and at the **infinite level**, respectively.

First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Topologies**


- [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of **quasi-syntomic rings** and sheaves on the category of **quasi-regular semi-perfectoid rings**, valued in any presentable ∞ -category.
- These correspond to moduli spaces at **finite levels** and at the **infinite level**, respectively.



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Topologies**


- [Bhatt–Morrow–Scholze ’19] There is an equivalence between sheaves on the category of **quasi-syntomic rings** and sheaves on the category of **quasi-regular semi-perfectoid rings**, valued in any presentable ∞ -category.
- These correspond to moduli spaces at **finite levels** and at the **infinite level**, respectively.



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- **Topologies**

- [Bhatt–Morrow–Scholze ’19] There is an equivalence between **sheaves** on the category of **quasi-syntomic rings** and **sheaves** on the category of **quasi-regular semi-perfectoid rings**, valued in any presentable ∞ -category.
- These correspond to moduli spaces at **finite levels** and at the **infinite level**, respectively.



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Besides as an additional structure on the moduli spaces, a further goal is to leverage the interplay of symmetries on this sheaf for computations, namely, homotopy-fixed-point spectral sequences for unstable chromatic homotopy groups.



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Besides as an additional structure on the moduli spaces, a further goal is to leverage the interplay of symmetries on this sheaf for computations, namely, homotopy-fixed-point spectral sequences for unstable chromatic homotopy groups.



First portraits: A filtered, equivariant, quasi-syntomic sheaf

- Besides as an additional structure on the moduli spaces, a further goal is to leverage the interplay of symmetries on this sheaf for computations, namely, homotopy-fixed-point spectral sequences for unstable chromatic homotopy groups.



Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding continuous evolution of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via exceptional optical devices.

Moduli spaces of physical systems, especially their singular loci, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding continuous evolution of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via exceptional optical devices.

Moduli spaces of physical systems, especially their singular loci, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding **continuous evolution** of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via exceptional optical devices.

Moduli spaces of physical systems, especially their singular loci, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding **continuous evolution** of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via **exceptional** optical devices.



Moduli spaces of physical systems, especially their singular loci, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding **continuous evolution** of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via **exceptional** optical devices.



Moduli spaces of physical systems, especially their **singular loci**, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding **continuous evolution** of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via **exceptional** optical devices.



Moduli spaces of physical systems, especially their **singular loci**, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding **continuous evolution** of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via **exceptional** optical devices.



Moduli spaces of physical systems, especially their **singular loci**, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Context and motivations

- Topological classifications for physical systems

Understanding **continuous evolution** of physical systems at the micro or quantum scale has a real impact on the larger-scale properties of materials. For example, holography is made possible via **exceptional** optical devices.



Moduli spaces of physical systems, especially their **singular loci**, play a pivotal role in designing such. Topological classifications enable physicists to fine-tune and create materials that can “do wonders” and cannot be found in nature, e.g., making invisible cloaks and other absorption devices.

Image credit: Natalya Burova / Getty Images

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally Hermitian matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally Hermitian matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - **Physical systems root in symmetries.** For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally Hermitian matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally Hermitian matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand degeneracies of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand **degeneracies** of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Algebraic setup

- Objects to parametrize: Hamiltonians with symmetries
 - Physical systems root in symmetries. For example, quantum mechanical systems can be described by their Hamiltonians, whose mathematical bearings are conventionally **Hermitian** matrices. Here, Hermiticity guarantees that the eigenvalues are real, corresponding to the fact that energies of the systems are observed to be real.
 - More recently, physicists have begun to model open systems by relaxing the Hermitian symmetry to allow eigenvalues with a nonzero imaginary part. This imaginary part measures energy exchange between the system and its surrounding environment. Still, some sorts of symmetry need to be imposed on the matrices to make them physically meaningful.
 - The size of the matrices corresponds to the number of energy band gaps. It is critical to understand **degeneracies** of eigenvalues and eigenvectors, across which the gaps close and open.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity–time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the non-isolated singular loci in the base spaces for these gapless 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the **non-isolated** singular loci in the base spaces for these **gapless** 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Moduli spaces for non-Hermitian Hamiltonians

- [Joint with Hongwei Jia et al.] By imposing the parity-time symmetry and a pseudo-Hermitian symmetry (with respect to a Lorentz-like transformation), we completely classified a generic family of 2-band systems as well as partially for certain 3-band systems:

$$H_2 = \begin{bmatrix} f_3 & f_2 \\ -f_2 & -f_3 \end{bmatrix} \quad H_3 = \begin{bmatrix} 1 - f_1 - f_2 & f_1 & f_2 \\ -f_1 & f_1 - f_3 & f_3 \\ -f_2 & f_3 & f_2 - f_3 \end{bmatrix}$$

where the parameters f_i are functions on the 3D momentum space $\{\mathbf{k}_x, \mathbf{k}_y, \mathbf{k}_z\}$. Moreover, my physics collaborators experimentally realized the above 3-band systems by circuits and verified our mathematical models.

- The moduli spaces keep track of eigen-energies (roots of the characteristic polynomial) as well as their corresponding eigenstates (the eigenvectors). Thus, they are “stratified vector bundles.” Interestingly, the stratification of the **non-isolated** singular loci in the base spaces for these **gapless** 3-band systems reveals transitions among diverse exceptional physical states, providing a test ground for exotic phenomena and anomalous effects.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A stratified space is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that

- each stratum $S \in \Sigma$ is a smooth manifold and
- if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.

A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.

- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A **stratified morphism** is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A stratified vector bundle consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Stratified vector bundles

- [Goresky–MacPherson '80, Pflaum '01, Ross '23, ...] A **stratified space** is a reasonable topological space X together with a reasonable partition Σ into reasonable subspaces such that
 - each stratum $S \in \Sigma$ is a smooth manifold and
 - if $S, T \in \Sigma$ are two strata with $S \cap \bar{T} \neq \emptyset$, then $S \subset \bar{T}$.A stratified morphism is a continuous map between stratified spaces that preserves the stratifications and restricts to be smooth over each stratum.
- [Ross '23] A **stratified vector bundle** consists of two stratified spaces (E, Σ_E) and (B, Σ_B) together with a stratified morphism $p: E \rightarrow B$ such that
 - for each $S \in \Sigma_B$, $E|_S := p^{-1}(S) \in \Sigma_E$ and $p: E|_S \rightarrow S$ is a smooth vector bundle,
 - the scalar multiplication map $\mu: \mathbb{R} \times E \rightarrow E$ is a stratified morphism.
- Besides our “eigenvectors fibrations” [Arnold '99, *Polymathematics*], other interesting examples include tangent bundles to stratified spaces and equivariant vector bundles.

Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “swallowtail catastrophe” [Thom, Arnold]:

$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

mechanical wave system

$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$

Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “swallowtail catastrophe” [Thom, Arnold]:

$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system


$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$

Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “[swallowtail catastrophe](#)” [Thom, Arnold]:



$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system


$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$

Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “[swallowtail catastrophe](#)” [Thom, Arnold]:



$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system



$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$

Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “[swallowtail catastrophe](#)” [Thom, Arnold]:



$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system


$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$


Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “[swallowtail catastrophe](#)” [Thom, Arnold]:




$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

A mechanical wave system



$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$




$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$


Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “[swallowtail catastrophe](#)” [Thom, Arnold]:




$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


A mechanical wave system



$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$




$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$



$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$


Second portraits: Swallowtails and more

- Portraits of base spaces for gapless systems, the central figure within these configurations being the “[swallowtail catastrophe](#)” [Thom, Arnold]:




$$\begin{bmatrix} 0 & -f_1 & -f_2 & -f_3 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$



A mechanical wave system



$$\begin{bmatrix} 1-f_1-f_2 & f_1 & f_2 \\ -f_1 & f_1-f_3 & f_3 \\ -f_2 & f_3 & f_2-f_3 \end{bmatrix}$$



$$\begin{bmatrix} 1 & f_1 & f_2 \\ -f_1 & -1 & f_3 \\ -f_2 & f_3 & -1 \end{bmatrix}$$



$$\begin{bmatrix} f_1f_2 & f_1 & f_2 \\ -f_1 & f_1 & f_3 \\ -f_2 & f_3 & f_2 \end{bmatrix}$$

Second portraits: Swallowtails and more

Further work in progress:

- Monodromy groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”
- Mathematical models and experimental realizations for “anomalous bulk–edge correspondence.”
This may be related to index theory for manifolds with fibered boundary [Yamashita ’20].

Second portraits: Swallowtails and more



Further work in progress:

- Monodromy groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”
- Mathematical models and experimental realizations for “anomalous bulk–edge correspondence.”
This may be related to index theory for manifolds with fibered boundary [Yamashita ’20].

Second portraits: Swallowtails and more

Further work in progress:

- **Monodromy** groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”




- Mathematical models and experimental realizations for “anomalous bulk–edge correspondence.”
This may be related to index theory for manifolds with fibered boundary [Yamashita ’20].

Second portraits: Swallowtails and more

Further work in progress:

- **Monodromy** groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”




- Mathematical models and experimental realizations for “anomalous bulk–edge correspondence.”
This may be related to index theory for manifolds with fibered boundary [Yamashita ’20].

Second portraits: Swallowtails and more

Further work in progress:

- **Monodromy** groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”




- Mathematical models and experimental realizations for “anomalous bulk–edge correspondence.”
This may be related to index theory for manifolds with fibered boundary [Yamashita ’20].

Second portraits: Swallowtails and more

Further work in progress:

- **Monodromy** groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”




- Mathematical models and experimental realizations for “anomalous **bulk-edge correspondence**. ”
This may be related to **index theory** for manifolds with fibered boundary [Yamashita ’20].

Second portraits: Swallowtails and more

Further work in progress:

- **Monodromy** groups of the stratified vector bundles in question computed by intersection homology/homotopy [Goresky–MacPherson, Gajer ’96].
What happens “upstairs?”



- Mathematical models and experimental realizations for “anomalous bulk-edge correspondence.”
This may be related to **index theory** for manifolds **with fibered boundary** [Yamashita ’20].

Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.

Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.

Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.

Third portraits: Context and motivations


- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.

Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.


Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.




Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.




Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.




Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.



Third portraits: Context and motivations

- From topological data analysis to topological deep learning:
 - Using persistent homology, Carlsson, Ishkhanov, de Silva, and Zomorodian qualitatively analyzed approximately 4.5×10^6 high-contrast local patches of natural images obtained by van Hateren and van der Schaaf and previously studied by Lee, Mumford, and Petersen. In their 2008 article, they discovered that as vectors of pixels, the image data were unevenly distributed over a Klein bottle within the 7-dimensional Euclidean sphere! We may view the Klein bottle as a moduli space for local image data.
 - A decade later, Love, Filippenko, Maroulas, and Carlsson have made the Klein bottle as a topological input for designing convolutional layers in neural networks that learn image data. Moreover, they have incorporated the tangent bundle of a Klein bottle into TCNNs for learning video data. Both learnings achieved higher accuracies with smaller training sets.



Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.

Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.

Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.

Third portraits: Moduli spaces for speech data


- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.

Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.


Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.




Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.




Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.




Third portraits: Moduli spaces for speech data

- [Joint with Pingyao Feng et al.] Motivated by the works of Carlsson and his collaborators', in consultation with Meng Yu of Tencent AI Lab, we have been investigating analogous questions for speech signals, with the additional tool of time-delay embedding for turning time series data to point clouds in Euclidean spaces.
 - For phonetic data, linguists created a charted “moduli space” of vowels:
 - Using speech files from SpeechBox, our topological approach achieved an average accuracy exceeding 95% in classifying voiced and voiceless consonants via machine learning.
 - A main goal remains to use topological methods to reveal moduli spaces for speech signals. We have also analyzed other types of time series data, such as animal behaviors in connection with biomedical engineering.



Thank you.



Credit: Europe-Spain/Alamy Stock Photo