

# Quasi-elliptic cohomology

Zhen Huan

Sun Yat-sen University

November 30, 2017

Plan.

- Motivation and construction
- The power operation
- The orthogonal  $G$ -spectrum

## An old idea of Witten

[Landweber]

The elliptic cohomology of a space  $X$  is related to the  $\mathbb{T}$ -equivariant K-theory of  $LX = \mathbb{C}^\infty(S^1, X)$  with the circle  $\mathbb{T}$  acting on  $LX$  by rotating loops.

It's surprisingly difficult to make this precise.

## Why?

In application, one needs to consider the case that a group  $G$  acts on  $X$ . In this case the loop space  $LX$  has rich structures as an orbifold.

I will show the relation between Tate K-theory and the loop space, which in fact bring a new theory, quasi-elliptic cohomology.

## An old idea of Witten

[Landweber]

The elliptic cohomology of a space  $X$  is related to the  $\mathbb{T}$ -equivariant K-theory of  $LX = \mathbb{C}^\infty(S^1, X)$  with the circle  $\mathbb{T}$  acting on  $LX$  by rotating loops.

It's surprisingly difficult to make this precise.

## Why?

In application, one needs to consider the case that a group  $G$  acts on  $X$ . In this case the loop space  $LX$  has rich structures as an orbifold.

I will show the relation between Tate K-theory and the loop space, which in fact bring a new theory, quasi-elliptic cohomology.

## An old idea of Witten

[Landweber]

The elliptic cohomology of a space  $X$  is related to the  $\mathbb{T}$ -equivariant K-theory of  $LX = \mathbb{C}^\infty(S^1, X)$  with the circle  $\mathbb{T}$  acting on  $LX$  by rotating loops.

It's surprisingly difficult to make this precise.

## Why?

In application, one needs to consider the case that a group  $G$  acts on  $X$ . In this case the loop space  $LX$  has rich structures as an orbifold.

I will show the relation between Tate K-theory and the loop space, which in fact bring a new theory, quasi-elliptic cohomology.

## An old idea of Witten

[Landweber]

The elliptic cohomology of a space  $X$  is related to the  $\mathbb{T}$ -equivariant K-theory of  $LX = \mathbb{C}^\infty(S^1, X)$  with the circle  $\mathbb{T}$  acting on  $LX$  by rotating loops.

It's surprisingly difficult to make this precise.

## Why?

In application, one needs to consider the case that a group  $G$  acts on  $X$ . In this case the loop space  $LX$  has rich structures as an orbifold.

I will show the relation between Tate K-theory and the loop space, which in fact bring a new theory, quasi-elliptic cohomology.

# The category of bibundles

## A bibundle from $\mathbb{H}$ to $\mathbb{G}$

[Schommer-Pries][Lerman]

a smooth manifold  $P$  together with

- the structure maps:

- $\tau : P \longrightarrow \mathbb{G}_0;$

- a surjective submersion  $\sigma : P \longrightarrow \mathbb{H}_0.$

- The action maps in  $Man_{\mathbb{G}_0 \times \mathbb{H}_0}$

- $\mathbb{G}_1 \times_{\tau} P \longrightarrow P;$

- $P \times_{\sigma} \mathbb{H}_1 \longrightarrow P$

such that

- $g_1 \cdot (g_2 \cdot p) = (g_1 g_2) \cdot p$  for all  $(g_1, g_2, p) \in \mathbb{G}_1 \times_{\tau} \mathbb{G}_1 \times_{\tau} P;$
- $(p \cdot h_1) \cdot h_2 = p \cdot (h_1 h_2)$  for all  $(p, h_1, h_2) \in P \times_{\sigma} \mathbb{H}_1 \times_{\sigma} \mathbb{H}_1;$
- $p \cdot u_H(\sigma(p)) = p$  and  $u_G(\tau(p)) \cdot p = p$  for all  $p \in P.$
- $g \cdot (p \cdot h) = (g \cdot p) \cdot h$  for all  $(g, p, h) \in \mathbb{G}_1 \times_{\tau} P \times_{\sigma} \mathbb{H}_1.$
- $\mathbb{G}_1 \times_{\tau} P \longrightarrow P \times_{\sigma} \mathbb{H}_1$   
 $(g, p) \mapsto (g \cdot p, p)$  is an isomorphism.

## Bibundle Map

[Schommer-Pries][Lerman]

a map  $P \longrightarrow P'$  over  $\mathbb{H}_0 \times \mathbb{G}_0$  commuting with the  $\mathbb{G}$ - and  $\mathbb{H}$ -action.

# The category of bibundles

## A bibundle from $\mathbb{H}$ to $\mathbb{G}$

[Schommer-Pries][Lerman]

a smooth manifold  $P$  together with

- the structure maps:
  - $\tau : P \longrightarrow \mathbb{G}_0$ ;
  - a surjective submersion  $\sigma : P \longrightarrow \mathbb{H}_0$ .
- The action maps in  $Man_{\mathbb{G}_0 \times \mathbb{H}_0}$ 
  - $\mathbb{G}_1 \times_{\sigma, \tau} P \longrightarrow P$ ;
  - $P \times_{\sigma, \tau} \mathbb{H}_1 \longrightarrow P$

such that

1.  $g_1 \cdot (g_2 \cdot p) = (g_1 g_2) \cdot p$  for all  $(g_1, g_2, p) \in \mathbb{G}_1 \times_{\sigma, \tau} \mathbb{G}_1 \times_{\sigma, \tau} P$ ;
2.  $(p \cdot h_1) \cdot h_2 = p \cdot (h_1 h_2)$  for all  $(p, h_1, h_2) \in P \times_{\sigma, \tau} \mathbb{H}_1 \times_{\sigma, \tau} \mathbb{H}_1$ ;
3.  $p \cdot u_H(\sigma(p)) = p$  and  $u_G(\tau(p)) \cdot p = p$  for all  $p \in P$ .
4.  $g \cdot (p \cdot h) = (g \cdot p) \cdot h$  for all  $(g, p, h) \in \mathbb{G}_1 \times_{\sigma, \tau} P \times_{\sigma, \tau} \mathbb{H}_1$ .
5.  $\mathbb{G}_1 \times_{\sigma, \tau} P \longrightarrow P \times_{\sigma, \tau} P$   $(g, p) \mapsto (g \cdot p, p)$  is an isomorphism.

## Bibundle Map

[Schommer-Pries][Lerman]

a map  $P \longrightarrow P'$  over  $\mathbb{H}_0 \times \mathbb{G}_0$  commuting with the  $\mathbb{G}$ - and  $\mathbb{H}$ -action.



# The category of bibundles

## A bibundle from $\mathbb{H}$ to $\mathbb{G}$

[Schommer-Pries][Lerman]

a smooth manifold  $P$  together with

- the structure maps:
  - $\tau : P \longrightarrow \mathbb{G}_0$ ;
  - a surjective submersion  $\sigma : P \longrightarrow \mathbb{H}_0$ .
- The action maps in  $Man_{\mathbb{G}_0 \times \mathbb{H}_0}$ 
  - $\mathbb{G}_1 \times_{\sigma, \tau} P \longrightarrow P$ ;
  - $P \times_{\sigma, \tau} \mathbb{H}_1 \longrightarrow P$

such that

1.  $g_1 \cdot (g_2 \cdot p) = (g_1 g_2) \cdot p$  for all  $(g_1, g_2, p) \in \mathbb{G}_1 \times_{\sigma, \tau} \mathbb{G}_1 \times_{\sigma, \tau} P$ ;
2.  $(p \cdot h_1) \cdot h_2 = p \cdot (h_1 h_2)$  for all  $(p, h_1, h_2) \in P \times_{\sigma, \tau} \mathbb{H}_1 \times_{\sigma, \tau} \mathbb{H}_1$ ;
3.  $p \cdot u_H(\sigma(p)) = p$  and  $u_G(\tau(p)) \cdot p = p$  for all  $p \in P$ .
4.  $g \cdot (p \cdot h) = (g \cdot p) \cdot h$  for all  $(g, p, h) \in \mathbb{G}_1 \times_{\sigma, \tau} P \times_{\sigma, \tau} \mathbb{H}_1$ .
5.  $\mathbb{G}_1 \times_{\sigma, \tau} P \longrightarrow P \times_{\sigma, \tau} P$   $(g, p) \mapsto (g \cdot p, p)$  is an isomorphism.

## Bibundle Map

[Schommer-Pries][Lerman]

a map  $P \longrightarrow P'$  over  $\mathbb{H}_0 \times \mathbb{G}_0$  commuting with the  $\mathbb{G}$ - and  $\mathbb{H}$ -action.

# The category of bibundles

## A bibundle from $\mathbb{H}$ to $\mathbb{G}$

[Schommer-Pries][Lerman]

a smooth manifold  $P$  together with

- the structure maps:
  - $\tau : P \longrightarrow \mathbb{G}_0$ ;
  - a surjective submersion  $\sigma : P \longrightarrow \mathbb{H}_0$ .
- The action maps in  $Man_{\mathbb{G}_0 \times \mathbb{H}_0}$ 
  - $\mathbb{G}_1 \times_{\sigma, \tau} P \longrightarrow P$ ;
  - $P \times_{\sigma, \tau} \mathbb{H}_1 \longrightarrow P$

such that

1.  $g_1 \cdot (g_2 \cdot p) = (g_1 g_2) \cdot p$  for all  $(g_1, g_2, p) \in \mathbb{G}_1 \times_{\sigma, \tau} \mathbb{G}_1 \times_{\sigma, \tau} P$ ;
2.  $(p \cdot h_1) \cdot h_2 = p \cdot (h_1 h_2)$  for all  $(p, h_1, h_2) \in P \times_{\sigma, \tau} \mathbb{H}_1 \times_{\sigma, \tau} \mathbb{H}_1$ ;
3.  $p \cdot u_H(\sigma(p)) = p$  and  $u_G(\tau(p)) \cdot p = p$  for all  $p \in P$ .
4.  $g \cdot (p \cdot h) = (g \cdot p) \cdot h$  for all  $(g, p, h) \in \mathbb{G}_1 \times_{\sigma, \tau} P \times_{\sigma, \tau} \mathbb{H}_1$ .
5.  $\mathbb{G}_1 \times_{\sigma, \tau} P \longrightarrow P \times_{\sigma, \tau} P$   $(g, p) \mapsto (g \cdot p, p)$  is an isomorphism.

## Bibundle Map

[Schommer-Pries][Lerman]

a map  $P \longrightarrow P'$  over  $\mathbb{H}_0 \times \mathbb{G}_0$  commuting with the  $\mathbb{G}$ - and  $\mathbb{H}$ -action.

# The Loop Spaces of Interest

Example ( $Loop_1(X // G) := Bibun(S^1 // *, X // G)$ )

- Objects:

$$\mathcal{P} := \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\}$$

- $\pi : \text{principal } G\text{-bundle over } S^1$
- $f : G\text{-equivariant;}$
- Morphism  $\mathcal{P} \rightarrow \mathcal{P}'$ :  $G\text{-bundle map } \alpha : P \rightarrow P'$

$$\begin{array}{ccccc} S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\ & \swarrow \pi' & \downarrow \alpha & \nearrow f' & \\ & & P' & & \end{array}$$

Example (Another Model  $Loop_2(X // G)$ )

- Objects:  $(\sigma, \gamma)$ 
  - $\sigma \in G$
  - $\gamma : \mathbb{R} \rightarrow X$  smooth  $\gamma(s+1) = \gamma(s) \cdot \sigma$
- Morphism  $(\sigma, \gamma) \rightarrow (\sigma', \gamma')$ :  $\alpha : \mathbb{R} \rightarrow G$  smooth,  $\gamma'(s) = \gamma(s)\alpha(s)$ .

# The Loop Spaces of Interest

Example ( $Loop_1(X // G) := Bibun(S^1 // *, X // G)$ )

- Objects:

$$\mathcal{P} := \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\}$$

- $\pi : \text{principal } G\text{-bundle over } S^1$
- $f : G\text{-equivariant;}$
- Morphism  $\mathcal{P} \rightarrow \mathcal{P}'$ :  $G\text{-bundle map } \alpha : P \rightarrow P'$

$$\begin{array}{ccccc} S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\ & \swarrow \pi' & \downarrow \alpha & \searrow f' & \\ & & P' & & \end{array}$$

Example (Another Model  $Loop_2(X // G)$ )

- Objects:  $(\sigma, \gamma)$ 
  - $\sigma \in G$
  - $\gamma : \mathbb{R} \rightarrow X$  smooth  $\gamma(s+1) = \gamma(s) \cdot \sigma$
- Morphism  $(\sigma, \gamma) \rightarrow (\sigma', \gamma')$ :  $\alpha : \mathbb{R} \rightarrow G$  smooth,  $\gamma'(s) = \gamma(s)\alpha(s)$ .

# The Loop Spaces of Interest

## Example ( $Loop_1(X // G) := Bibun(S^1 // *, X // G)$ )

- Objects:

$$\mathcal{P} := \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\}$$

- $\pi : \text{principal } G\text{-bundle over } S^1$
- $f : G\text{-equivariant;}$
- Morphism  $\mathcal{P} \rightarrow \mathcal{P}'$ :  $G\text{-bundle map } \alpha : P \rightarrow P'$

$$\begin{array}{ccccc} S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\ & \swarrow \pi' & \downarrow \alpha & \searrow f' & \\ & & P' & & \end{array}$$

## Example (Another Model $Loop_2(X // G)$ )

- Objects:  $(\sigma, \gamma)$ 
  - $\sigma \in G$
  - $\gamma : \mathbb{R} \rightarrow X$  smooth  $\gamma(s+1) = \gamma(s) \cdot \sigma$
- Morphism  $(\sigma, \gamma) \rightarrow (\sigma', \gamma')$ :  $\alpha : \mathbb{R} \rightarrow G$  smooth,  $\gamma'(s) = \gamma(s)\alpha(s)$ .

# The Loop Spaces of Interest

## Example ( $Loop_1(X // G) := Bibun(S^1 // *, X // G)$ )

- Objects:

$$\mathcal{P} := \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\}$$

- $\pi : \text{principal } G\text{-bundle over } S^1$
- $f : G\text{-equivariant;}$
- Morphism  $\mathcal{P} \rightarrow \mathcal{P}'$ :  $G\text{-bundle map } \alpha : P \rightarrow P'$

$$\begin{array}{ccccc} S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\ & \swarrow \pi' & \downarrow \alpha & \searrow f' & \\ & & P' & & \end{array}$$

## Example (Another Model $Loop_2(X // G)$ )

- Objects:  $(\sigma, \gamma)$ 
  - $\sigma \in G$
  - $\gamma : \mathbb{R} \rightarrow X$  smooth  $\gamma(s+1) = \gamma(s) \cdot \sigma$
- Morphism  $(\sigma, \gamma) \rightarrow (\sigma', \gamma')$ :  $\alpha : \mathbb{R} \rightarrow G$  smooth,  $\gamma'(s) = \gamma(s)\alpha(s)$ .

## Example ( $Loop_1^{ext}(X // G)$ )

- the same objects as  $Loop_1(X // G)$ ;
- $(t, \alpha) : \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\} \longrightarrow \{S^1 \xleftarrow{\pi'} P' \xrightarrow{f'} X\}$ 
  - $t \in \mathbb{T}$
  - $\alpha : P \longrightarrow P'$ :  $G$ -bundle map

$$\begin{array}{ccccc}
 S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\
 t \downarrow & & \downarrow \alpha & \nearrow f' & \\
 S^1 & \xleftarrow{\pi'} & P' & & 
 \end{array}$$

## Example ( $Loop_2^{ext}(X // G)$ )

- the same objects as  $Loop_2(X // G)$ ;
- A morphism

$$(\alpha, t) : (\sigma, \gamma) \longrightarrow (\sigma', \gamma')$$

- $\alpha : \mathbb{R} \longrightarrow G$  smooth
- $t \in \mathbb{R}$
- $\gamma'(s) = \gamma(s - t)\alpha(s - t).$

## Example ( $Loop_1^{ext}(X // G)$ )

- the same objects as  $Loop_1(X // G)$ ;
- $(t, \alpha) : \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\} \longrightarrow \{S^1 \xleftarrow{\pi'} P' \xrightarrow{f'} X\}$ 
  - $t \in \mathbb{T}$
  - $\alpha : P \longrightarrow P'$ :  $G$ -bundle map

$$\begin{array}{ccccc}
 S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\
 t \downarrow & & \downarrow \alpha & \nearrow f' & \\
 S^1 & \xleftarrow{\pi'} & P' & & 
 \end{array}$$

## Example ( $Loop_2^{ext}(X // G)$ )

- the same objects as  $Loop_2(X // G)$ ;
- A morphism

$$(\alpha, t) : (\sigma, \gamma) \longrightarrow (\sigma', \gamma')$$

- $\alpha : \mathbb{R} \longrightarrow G$  smooth
- $t \in \mathbb{R}$
- $\gamma'(s) = \gamma(s - t)\alpha(s - t).$



## Example ( $Loop_1^{ext}(X // G)$ )

- the same objects as  $Loop_1(X // G)$ ;
- $(t, \alpha) : \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\} \longrightarrow \{S^1 \xleftarrow{\pi'} P' \xrightarrow{f'} X\}$ 
  - $t \in \mathbb{T}$
  - $\alpha : P \longrightarrow P'$ :  $G$ -bundle map

$$\begin{array}{ccccc}
 S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\
 t \downarrow & & \downarrow \alpha & \nearrow f' & \\
 S^1 & \xleftarrow{\pi'} & P' & & 
 \end{array}$$

## Example ( $Loop_2^{ext}(X // G)$ )

- the same objects as  $Loop_2(X // G)$ ;
- A morphism

$$(\alpha, t) : (\sigma, \gamma) \longrightarrow (\sigma', \gamma')$$

- $\alpha : \mathbb{R} \longrightarrow G$  smooth
- $t \in \mathbb{R}$
- $\gamma'(s) = \gamma(s - t)\alpha(s - t).$

## Example ( $Loop_1^{ext}(X // G)$ )

- the same objects as  $Loop_1(X // G)$ ;
- $(t, \alpha) : \{S^1 \xleftarrow{\pi} P \xrightarrow{f} X\} \longrightarrow \{S^1 \xleftarrow{\pi'} P' \xrightarrow{f'} X\}$ 
  - $t \in \mathbb{T}$
  - $\alpha : P \longrightarrow P'$ :  $G$ -bundle map

$$\begin{array}{ccccc}
 S^1 & \xleftarrow{\pi} & P & \xrightarrow{f} & X \\
 t \downarrow & & \downarrow \alpha & \nearrow f' & \\
 S^1 & \xleftarrow{\pi'} & P' & & 
 \end{array}$$

## Example ( $Loop_2^{ext}(X // G)$ )

- the same objects as  $Loop_2(X // G)$ ;
- A morphism

$$(\alpha, t) : (\sigma, \gamma) \longrightarrow (\sigma', \gamma')$$

- $\alpha : \mathbb{R} \longrightarrow G$  smooth
- $t \in \mathbb{R}$
- $\gamma'(s) = \gamma(s - t)\alpha(s - t).$

# Good Groupoid or Not?

## A skeleton of $Loop_2^{\text{ext}}(X // G)$

- $\mathcal{L}_g X$ : the space of objects  $(g, \gamma)$  in  $Loop_2(X // G)$ .
- $L_g G = \{\alpha : \mathbb{R} \longrightarrow G \mid \alpha(s+1) = g^{-1}\alpha(s)g\}$ , the gauge group of the principal  $G$ -bundle  $P_g := \mathbb{R} \times G / (s+1, a) \sim (s, ga)$  over  $S^1$ ;
- $L_g G \rtimes \mathbb{T}$ :  $(\alpha, t) \cdot (\alpha', t') := (s \mapsto \alpha(s)\alpha'(s+t), t+t')$ .
- For any  $(\alpha, t) \in L_g G \rtimes \mathbb{T}$ , and  $\gamma \in \mathcal{L}_g X$ ,  
 $\gamma \cdot (\alpha, t) := (s \mapsto \gamma(s-t) \cdot \alpha(s-t))$ .

$\coprod_{g \in \pi_0 G / \text{conj}} \mathcal{L}_g X // L_g G \rtimes \mathbb{T}$  is a skeleton of  $Loop_2^{\text{ext}}(X // G)$ .

## Problem

$L_g G \rtimes \mathbb{T}$  is an infinite dimensional topological group when  $G$  is not finite.

# Good Groupoid or Not?

## A skeleton of $Loop_2^{\text{ext}}(X // G)$

- $\mathcal{L}_g X$ : the space of objects  $(g, \gamma)$  in  $Loop_2(X // G)$ .
- $L_g G = \{\alpha : \mathbb{R} \longrightarrow G \mid \alpha(s+1) = g^{-1}\alpha(s)g\}$ , the gauge group of the principal  $G$ -bundle  $P_g := \mathbb{R} \times G / (s+1, a) \sim (s, ga)$  over  $S^1$ ;
- $L_g G \rtimes \mathbb{T}$ :  $(\alpha, t) \cdot (\alpha', t') := (s \mapsto \alpha(s)\alpha'(s+t), t+t')$ .
- For any  $(\alpha, t) \in L_g G \rtimes \mathbb{T}$ , and  $\gamma \in \mathcal{L}_g X$ ,  
 $\gamma \cdot (\alpha, t) := (s \mapsto \gamma(s-t) \cdot \alpha(s-t))$ .

$\coprod_{g \in \pi_0 G / \text{conj}} \mathcal{L}_g X // L_g G \rtimes \mathbb{T}$  is a skeleton of  $Loop_2^{\text{ext}}(X // G)$ .

## Problem

$L_g G \rtimes \mathbb{T}$  is an infinite dimensional topological group when  $G$  is not finite.

# Good Groupoid or Not?

## A skeleton of $Loop_2^{\text{ext}}(X // G)$

- $\mathcal{L}_g X$ : the space of objects  $(g, \gamma)$  in  $Loop_2(X // G)$ .
- $L_g G = \{\alpha : \mathbb{R} \longrightarrow G \mid \alpha(s+1) = g^{-1}\alpha(s)g\}$ , the gauge group of the principal  $G$ -bundle  $P_g := \mathbb{R} \times G / (s+1, a) \sim (s, ga)$  over  $S^1$ ;
- $L_g G \rtimes \mathbb{T}$ :  $(\alpha, t) \cdot (\alpha', t') := (s \mapsto \alpha(s)\alpha'(s+t), t+t')$ .
- For any  $(\alpha, t) \in L_g G \rtimes \mathbb{T}$ , and  $\gamma \in \mathcal{L}_g X$ ,  
 $\gamma \cdot (\alpha, t) := (s \mapsto \gamma(s-t) \cdot \alpha(s-t))$ .

$\coprod_{g \in \pi_0 G / \text{conj}} \mathcal{L}_g X // L_g G \rtimes \mathbb{T}$  is a skeleton of  $Loop_2^{\text{ext}}(X // G)$ .

## Problem

$L_g G \rtimes \mathbb{T}$  is an infinite dimensional topological group when  $G$  is not finite.

# Good Groupoid or Not?

## A skeleton of $Loop_2^{\text{ext}}(X // G)$

- $\mathcal{L}_g X$ : the space of objects  $(g, \gamma)$  in  $Loop_2(X // G)$ .
- $L_g G = \{\alpha : \mathbb{R} \longrightarrow G \mid \alpha(s+1) = g^{-1}\alpha(s)g\}$ , the gauge group of the principal  $G$ -bundle  $P_g := \mathbb{R} \times G / (s+1, a) \sim (s, ga)$  over  $S^1$ ;
- $L_g G \rtimes \mathbb{T}$ :  $(\alpha, t) \cdot (\alpha', t') := (s \mapsto \alpha(s)\alpha'(s+t), t+t')$ .
- For any  $(\alpha, t) \in L_g G \rtimes \mathbb{T}$ , and  $\gamma \in \mathcal{L}_g X$ ,  
 $\gamma \cdot (\alpha, t) := (s \mapsto \gamma(s-t) \cdot \alpha(s-t))$ .

$\coprod_{g \in \pi_0 G / \text{conj}} \mathcal{L}_g X // L_g G \rtimes \mathbb{T}$  is a skeleton of  $Loop_2^{\text{ext}}(X // G)$ .

## Problem

$L_g G \rtimes \mathbb{T}$  is an infinite dimensional topological group when  $G$  is not finite.

# Good Groupoid or Not?

## A skeleton of $Loop_2^{\text{ext}}(X // G)$

- $\mathcal{L}_g X$ : the space of objects  $(g, \gamma)$  in  $Loop_2(X // G)$ .
- $L_g G = \{\alpha : \mathbb{R} \longrightarrow G \mid \alpha(s+1) = g^{-1}\alpha(s)g\}$ , the gauge group of the principal  $G$ -bundle  $P_g := \mathbb{R} \times G / (s+1, a) \sim (s, ga)$  over  $S^1$ ;
- $L_g G \rtimes \mathbb{T}$ :  $(\alpha, t) \cdot (\alpha', t') := (s \mapsto \alpha(s)\alpha'(s+t), t+t')$ .
- For any  $(\alpha, t) \in L_g G \rtimes \mathbb{T}$ , and  $\gamma \in \mathcal{L}_g X$ ,  
 $\gamma \cdot (\alpha, t) := (s \mapsto \gamma(s-t) \cdot \alpha(s-t))$ .

$\coprod_{g \in \pi_0 G / \text{conj}} \mathcal{L}_g X // L_g G \rtimes \mathbb{T}$  is a skeleton of  $Loop_2^{\text{ext}}(X // G)$ .

## Problem

$L_g G \rtimes \mathbb{T}$  is an infinite dimensional topological group when  $G$  is not finite.

# Good Groupoid or Not?

## A skeleton of $Loop_2^{\text{ext}}(X // G)$

- $\mathcal{L}_g X$ : the space of objects  $(g, \gamma)$  in  $Loop_2(X // G)$ .
- $L_g G = \{\alpha : \mathbb{R} \longrightarrow G \mid \alpha(s+1) = g^{-1}\alpha(s)g\}$ , the gauge group of the principal  $G$ -bundle  $P_g := \mathbb{R} \times G / (s+1, a) \sim (s, ga)$  over  $S^1$ ;
- $L_g G \rtimes \mathbb{T}$ :  $(\alpha, t) \cdot (\alpha', t') := (s \mapsto \alpha(s)\alpha'(s+t), t+t')$ .
- For any  $(\alpha, t) \in L_g G \rtimes \mathbb{T}$ , and  $\gamma \in \mathcal{L}_g X$ ,  
 $\gamma \cdot (\alpha, t) := (s \mapsto \gamma(s-t) \cdot \alpha(s-t))$ .

$\coprod_{g \in \pi_0 G / \text{conj}} \mathcal{L}_g X // L_g G \rtimes \mathbb{T}$  is a skeleton of  $Loop_2^{\text{ext}}(X // G)$ .

## Problem

$L_g G \rtimes \mathbb{T}$  is an infinite dimensional topological group when  $G$  is not finite.



# Quasi-elliptic cohomology

Consider the subgroupoid  $\Lambda(X//G)$  instead

$$\Lambda(X//G) := \coprod_{g \in G_{conj}^{tors}} X^g // \Lambda_G(g)$$

$G_{conj}^{tors}$ : a set of representatives of  $G$ –conjugacy classes in  $G^{tors}$ ;

$$\Lambda_G(g) = C_G(g) \times \mathbb{R} / \langle (g, -1) \rangle$$

$\Lambda_G(g)$  acts on  $X^g$  by

$$[h, t] \cdot x := h \cdot x.$$

$QEll$  as equivariant  $K$ –theories

$$QEll_G(X) \cong \prod_{g \in G_{conj}^{tors}} K_{\Lambda_G(g)}(X^g)$$

Relation with Tate  $K$ –theory

$$QEll_G^*(X) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong K_{Tate}^*(X//G).$$

# Quasi-elliptic cohomology

Consider the subgroupoid  $\Lambda(X//G)$  instead

$$\Lambda(X//G) := \coprod_{g \in G_{conj}^{tors}} X^g // \Lambda_G(g)$$

$G_{conj}^{tors}$ : a set of representatives of  $G$ -conjugacy classes in  $G^{tors}$ ;

$$\Lambda_G(g) = C_G(g) \times \mathbb{R} / \langle (g, -1) \rangle$$

$\Lambda_G(g)$  acts on  $X^g$  by

$$[h, t] \cdot x := h \cdot x.$$

$QEll$  as equivariant  $K$ -theories

$$QEll_G(X) \cong \prod_{g \in G_{conj}^{tors}} K_{\Lambda_G(g)}(X^g)$$

Relation with Tate  $K$ -theory

$$QEll_G^*(X) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong K_{Tate}^*(X//G).$$

# Quasi-elliptic cohomology

Consider the subgroupoid  $\Lambda(X//G)$  instead

$$\Lambda(X//G) := \coprod_{g \in G_{conj}^{tors}} X^g // \Lambda_G(g)$$

$G_{conj}^{tors}$ : a set of representatives of  $G$ -conjugacy classes in  $G^{tors}$ ;

$$\Lambda_G(g) = C_G(g) \times \mathbb{R} / \langle (g, -1) \rangle$$

$\Lambda_G(g)$  acts on  $X^g$  by

$$[h, t] \cdot x := h \cdot x.$$

$QEll$  as equivariant  $K$ -theories

$$QEll_G(X) \cong \prod_{g \in G_{conj}^{tors}} K_{\Lambda_G(g)}(X^g)$$

Relation with Tate  $K$ -theory

$$QEll_G^*(X) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong K_{Tate}^*(X//G).$$

# Setting up the theory

## Restriction

$$\phi : X // G \longrightarrow Y // H \implies \Lambda(\phi) : \Lambda(X // G) \longrightarrow \Lambda(Y // H)$$

$$\begin{array}{ccc} QEII^*(Y // H) & \xrightarrow{\phi^*} & QEII^*(X // G) \\ \pi_{\phi(\tau)} \downarrow & & \downarrow \pi_\tau \\ K_{\Lambda_H(\phi(\tau))}^*(Y^{\phi(\tau)}) & \xrightarrow{\phi_\Lambda^*} & K_{\Lambda_G(\tau)}^*(X^\tau) \end{array}$$

## Künneth Map

$$K_{\Lambda_G(\sigma)}(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}(Y^\tau) \longrightarrow K_{\Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau)}((X \times Y)^{(\sigma, \tau)}) \cong K_{\Lambda_{G \times H}(\sigma, \tau)}((X \times Y)^{(\sigma, \tau)}) \text{ where}$$

$$\Lambda_{G \times H}(\sigma, \tau) \cong \Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau): \text{ pullback of } \Lambda_G(\sigma) \xrightarrow{\pi} \mathbb{T} \xleftarrow{\pi} \Lambda_H(\tau).$$

$$QEII_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEII_H^*(Y) := \prod_{\sigma \in G_{conj}^{tors}, \tau \in H_{conj}^{tors}} K_{\Lambda_G(\sigma)}^*(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}^*(Y^\tau).$$

$$\text{The Künneth map: } QEII_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEII_H^*(Y) \longrightarrow QEII_{G \times H}^*(X \times Y).$$

# Setting up the theory

## Restriction

$$\phi : X // G \longrightarrow Y // H \implies \Lambda(\phi) : \Lambda(X // G) \longrightarrow \Lambda(Y // H)$$

$$QEII^*(Y // H) \xrightarrow{\phi^*} QEII^*(X // G)$$

$$\pi_{\phi(\tau)} \downarrow$$

$$\pi_\tau \downarrow$$

$$K_{\Lambda_H(\phi(\tau))}^*(Y^{\phi(\tau)}) \xrightarrow{\phi_\Lambda^*} K_{\Lambda_G(\tau)}^*(X^\tau)$$

## Künneth Map

$$K_{\Lambda_G(\sigma)}(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}(Y^\tau) \longrightarrow K_{\Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau)}((X \times Y)^{(\sigma, \tau)}) \cong K_{\Lambda_{G \times H}(\sigma, \tau)}((X \times Y)^{(\sigma, \tau)}) \text{ where}$$

$$\Lambda_{G \times H}(\sigma, \tau) \cong \Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau): \text{ pullback of } \Lambda_G(\sigma) \xrightarrow{\pi} \mathbb{T} \xleftarrow{\pi} \Lambda_H(\tau).$$

$$QEII_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEII_H^*(Y) := \prod_{\sigma \in G_{conj}^{tors}, \tau \in H_{conj}^{tors}} K_{\Lambda_G(\sigma)}^*(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}^*(Y^\tau).$$

$$\text{The Künneth map: } QEII_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEII_H^*(Y) \longrightarrow QEII_{G \times H}^*(X \times Y).$$

# Setting up the theory

## Restriction

$$\phi : X // G \longrightarrow Y // H \implies \Lambda(\phi) : \Lambda(X // G) \longrightarrow \Lambda(Y // H)$$

$$\begin{array}{ccc} QEll^*(Y // H) & \xrightarrow{\phi^*} & QEll^*(X // G) \\ \pi_{\phi(\tau)} \downarrow & & \downarrow \pi_\tau \\ K_{\Lambda_H(\phi(\tau))}^*(Y^{\phi(\tau)}) & \xrightarrow{\phi_\Lambda^*} & K_{\Lambda_G(\tau)}^*(X^\tau) \end{array}$$

## Künneth Map

$$K_{\Lambda_G(\sigma)}(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}(Y^\tau) \longrightarrow K_{\Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau)}((X \times Y)^{(\sigma, \tau)}) \cong K_{\Lambda_{G \times H}(\sigma, \tau)}((X \times Y)^{(\sigma, \tau)}) \text{ where}$$

$$\Lambda_{G \times H}(\sigma, \tau) \cong \Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau): \text{ pullback of } \Lambda_G(\sigma) \xrightarrow{\pi} \mathbb{T} \xleftarrow{\pi} \Lambda_H(\tau).$$

$$QEll_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEll_H^*(Y) := \prod_{\sigma \in G_{conj}^{tors}, \tau \in H_{conj}^{tors}} K_{\Lambda_G(\sigma)}^*(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}^*(Y^\tau).$$

$$\text{The Künneth map: } QEll_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEll_H^*(Y) \longrightarrow QEll_{G \times H}^*(X \times Y).$$

# Setting up the theory

## Restriction

$$\phi : X // G \longrightarrow Y // H \implies \Lambda(\phi) : \Lambda(X // G) \longrightarrow \Lambda(Y // H)$$

$$\begin{array}{ccc} QEll^*(Y // H) & \xrightarrow{\phi^*} & QEll^*(X // G) \\ \pi_{\phi(\tau)} \downarrow & & \downarrow \pi_\tau \\ K_{\Lambda_H(\phi(\tau))}^*(Y^{\phi(\tau)}) & \xrightarrow{\phi_\Lambda^*} & K_{\Lambda_G(\tau)}^*(X^\tau) \end{array}$$

## Künneth Map

$$K_{\Lambda_G(\sigma)}(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}(Y^\tau) \longrightarrow K_{\Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau)}((X \times Y)^{(\sigma, \tau)}) \cong K_{\Lambda_{G \times H}(\sigma, \tau)}((X \times Y)^{(\sigma, \tau)}) \text{ where}$$

$$\Lambda_{G \times H}(\sigma, \tau) \cong \Lambda_G(\sigma) \times_{\mathbb{T}} \Lambda_H(\tau): \text{ pullback of } \Lambda_G(\sigma) \xrightarrow{\pi} \mathbb{T} \xleftarrow{\pi} \Lambda_H(\tau).$$

$$QEll_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEll_H^*(Y) := \prod_{\sigma \in G_{conj}^{tors}, \tau \in H_{conj}^{tors}} K_{\Lambda_G(\sigma)}^*(X^\sigma) \otimes_{\mathbb{Z}[q^\pm]} K_{\Lambda_H(\tau)}^*(Y^\tau).$$

$$\text{The Künneth map: } QEll_G^*(X) \hat{\otimes}_{\mathbb{Z}[q^\pm]} QEll_H^*(Y) \longrightarrow QEll_{G \times H}^*(X \times Y).$$

# Setting up the theory

## The change-of-group isomorphism

- $H$ : closed subgroup of  $G$ ;
- $X$ :  $H$ -space;
- $\phi : H \longrightarrow G$  is the inclusion.

## Theorem

*The change-of-group map  $\rho_H^G$  is an isomorphism.*

$$\rho_H^G : QEll_G^*(G \times_H X) \xrightarrow{\phi^*} QEll_H^*(G \times_H X) \xrightarrow{i^*} QEll_H^*(X)$$

- $\phi^*$ : the restriction map
- $i : X \longrightarrow G \times_H X : i(x) = [e, x]$ .

## Induced map

$$\mathcal{I}_H^G : QEll(X // H) \xrightarrow{\cong} QEll((G \times_H X) // G) \longrightarrow QEll(X // G)$$

- the first map: the change-of-group isomorphism
- the second: the finite covering  $\Lambda(G \times_H X // G) \longrightarrow \Lambda(X // G)$   
 $\text{obj } (\sigma, [g, x]) \mapsto (\sigma, gx); \text{ mor } ([g', t], (\sigma, [g, x])) \mapsto ([g', t], (gx, \sigma)).$



# Setting up the theory

## The change-of-group isomorphism

- $H$ : closed subgroup of  $G$ ;
- $X$ :  $H$ -space;
- $\phi : H \longrightarrow G$  is the inclusion.

## Theorem

*The change-of-group map  $\rho_H^G$  is an isomorphism.*

$$\rho_H^G : QEll_G^*(G \times_H X) \xrightarrow{\phi^*} QEll_H^*(G \times_H X) \xrightarrow{i^*} QEll_H^*(X)$$

- $\phi^*$ : the restriction map
- $i : X \longrightarrow G \times_H X : i(x) = [e, x]$ .

## Induced map

$$\mathcal{I}_H^G : QEll(X // H) \xrightarrow{\cong} QEll((G \times_H X) // G) \longrightarrow QEll(X // G)$$

- the first map: the change-of-group isomorphism
- the second: the finite covering  $\Lambda(G \times_H X // G) \longrightarrow \Lambda(X // G)$   
 $\text{obj } (\sigma, [g, x]) \mapsto (\sigma, gx); \text{ mor } ([g', t], (\sigma, [g, x])) \mapsto ([g', t], (gx, \sigma)).$

# Setting up the theory

## The change-of-group isomorphism

- $H$ : closed subgroup of  $G$ ;
- $X$ :  $H$ -space;
- $\phi : H \longrightarrow G$  is the inclusion.

## Theorem

*The change-of-group map  $\rho_H^G$  is an isomorphism.*

$$\rho_H^G : QEll_G^*(G \times_H X) \xrightarrow{\phi^*} QEll_H^*(G \times_H X) \xrightarrow{i^*} QEll_H^*(X)$$

- $\phi^*$ : the restriction map
- $i : X \longrightarrow G \times_H X : i(x) = [e, x]$ .

## Induced map

$$\mathcal{I}_H^G : QEll(X // H) \xrightarrow{\cong} QEll((G \times_H X) // G) \longrightarrow QEll(X // G)$$

- the first map: the change-of-group isomorphism
- the second: the finite covering  $\Lambda(G \times_H X // G) \longrightarrow \Lambda(X // G)$   
obj  $(\sigma, [g, x]) \mapsto (\sigma, gx)$ ; mor  $([g', t], (\sigma, [g, x])) \mapsto ([g', t], (gx, \sigma))$ .

# Power Operation of Equivariant cohomology theories

## Power Operation of K-theory

[Atiyah]

$$P_n : K(X) \longrightarrow K_{\Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Power Operation of equivariant K-theory

[Atiyah]

$$P_n : K_G(X) \longrightarrow K_{G \wr \Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Wreath product $G \wr \Sigma_n$

$$(g_1, \dots, g_n, \sigma) \cdot (h_1, \dots, h_n, \tau) := (g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)}, \sigma \tau).$$

$$\text{Group action: } (x_1, \dots, x_n) \cdot (g_1, \dots, g_n, \sigma) := (x_{\sigma(1)} g_{\sigma(1)}, \dots, x_{\sigma(n)} g_{\sigma(n)}).$$

## Definition of Equivariant Power Operation

[May][Ganter]

$$P_n : E_G(X) \longrightarrow E_{G \wr \Sigma_n}(X^{\times n})$$

satisfying some axioms.

# Power Operation of Equivariant cohomology theories

## Power Operation of K-theory

[Atiyah]

$$P_n : K(X) \longrightarrow K_{\Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Power Operation of equivariant K-theory

[Atiyah]

$$P_n : K_G(X) \longrightarrow K_{G \wr \Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Wreath product $G \wr \Sigma_n$

$$(g_1, \dots, g_n, \sigma) \cdot (h_1, \dots, h_n, \tau) := (g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)}, \sigma \tau).$$

$$\text{Group action: } (x_1, \dots, x_n) \cdot (g_1, \dots, g_n, \sigma) := (x_{\sigma(1)} g_{\sigma(1)}, \dots, x_{\sigma(n)} g_{\sigma(n)}).$$

## Definition of Equivariant Power Operation

[May][Ganter]

$$P_n : E_G(X) \longrightarrow E_{G \wr \Sigma_n}(X^{\times n})$$

satisfying some axioms.

# Power Operation of Equivariant cohomology theories

## Power Operation of K-theory

[Atiyah]

$$P_n : K(X) \longrightarrow K_{\Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Power Operation of equivariant K-theory

[Atiyah]

$$P_n : K_G(X) \longrightarrow K_{G \wr \Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Wreath product $G \wr \Sigma_n$

$$(g_1, \dots, g_n, \sigma) \cdot (h_1, \dots, h_n, \tau) := (g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)}, \sigma \tau).$$

$$\text{Group action: } (x_1, \dots, x_n) \cdot (g_1, \dots, g_n, \sigma) := (x_{\sigma(1)} g_{\sigma(1)}, \dots, x_{\sigma(n)} g_{\sigma(n)}).$$

## Definition of Equivariant Power Operation

[May][Ganter]

$$P_n : E_G(X) \longrightarrow E_{G \wr \Sigma_n}(X^{\times n})$$

satisfying some axioms.

# Power Operation of Equivariant cohomology theories

## Power Operation of K-theory

[Atiyah]

$$P_n : K(X) \longrightarrow K_{\Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Power Operation of equivariant K-theory

[Atiyah]

$$P_n : K_G(X) \longrightarrow K_{G \wr \Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Wreath product $G \wr \Sigma_n$

$$(g_1, \dots, g_n, \sigma) \cdot (h_1, \dots, h_n, \tau) := (g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)}, \sigma \tau).$$

$$\text{Group action: } (x_1, \dots, x_n) \cdot (g_1, \dots, g_n, \sigma) := (x_{\sigma(1)} g_{\sigma(1)}, \dots, x_{\sigma(n)} g_{\sigma(n)}).$$

## Definition of Equivariant Power Operation

[May][Ganter]

$$P_n : E_G(X) \longrightarrow E_{G \wr \Sigma_n}(X^{\times n})$$

satisfying some axioms.

# Power Operation of Equivariant cohomology theories

## Power Operation of K-theory

[Atiyah]

$$P_n : K(X) \longrightarrow K_{\Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Power Operation of equivariant K-theory

[Atiyah]

$$P_n : K_G(X) \longrightarrow K_{G \wr \Sigma_n}(X^{\times n}), \quad V \mapsto V^{\boxtimes n}$$

## Wreath product $G \wr \Sigma_n$

$$(g_1, \dots, g_n, \sigma) \cdot (h_1, \dots, h_n, \tau) := (g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)}, \sigma \tau).$$

$$\text{Group action: } (x_1, \dots, x_n) \cdot (g_1, \dots, g_n, \sigma) := (x_{\sigma(1)} g_{\sigma(1)}, \dots, x_{\sigma(n)} g_{\sigma(n)}).$$

## Definition of Equivariant Power Operation

[May][Ganter]

$$P_n : E_G(X) \longrightarrow E_{G \wr \Sigma_n}(X^{\times n})$$

satisfying some axioms.

# Power Operation

Quasi-elliptic cohomology has power operations, which gives it the structure of an " $H_\infty$ -ring theory" [Ganter 06].

Atiyah's Power Operation

[Ganter]

$V$ : a vector bundle over  $\Lambda(X // G)$ .

$P_n(V) := V^{\widehat{\otimes}_{\mathbb{Z}[q^\pm]} }^n$  defines an operation

$$P_n : QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n})$$

$$\mathbb{P}_n = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} \mathbb{P}_{(\underline{g}, \sigma)} :$$

$$QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n}) = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)}).$$

$$\mathbb{P}_{(\underline{g}, \sigma)} : QEll_G(X) \xrightarrow{U^*} K_{orb}(\Lambda_{(\underline{g}, \sigma)}(X)) \xrightarrow{(\ )_k^\wedge} K_{orb}(\Lambda_{(\underline{g}, \sigma)}^{var}(X))$$

$$\xrightarrow{\boxtimes} K_{orb}(d_{(\underline{g}, \sigma)}(X)) \xrightarrow{f_{(\underline{g}, \sigma)}^*} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)})$$



# Power Operation

Quasi-elliptic cohomology has power operations, which gives it the structure of an " $H_\infty$ -ring theory" [Ganter 06].

## Atiyah's Power Operation

[Ganter]

$V$ : a vector bundle over  $\Lambda(X//G)$ .

$P_n(V) := V^{\hat{\otimes}_{\mathbb{Z}[q^\pm]} n}$  defines an operation

$$P_n : QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n})$$

$$\mathbb{P}_n = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} \mathbb{P}_{(\underline{g}, \sigma)} :$$

$$QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n}) = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)}).$$

$$\mathbb{P}_{(\underline{g}, \sigma)} : QEll_G(X) \xrightarrow{U^*} K_{orb}(\Lambda_{(\underline{g}, \sigma)}(X)) \xrightarrow{(\ )_k^\Lambda} K_{orb}(\Lambda_{(\underline{g}, \sigma)}^{var}(X))$$

$$\xrightarrow{\boxtimes} K_{orb}(d_{(\underline{g}, \sigma)}(X)) \xrightarrow{f_{(\underline{g}, \sigma)}^*} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)})$$

# Power Operation

Quasi-elliptic cohomology has power operations, which gives it the structure of an " $H_\infty$ -ring theory" [Ganter 06].

## Atiyah's Power Operation

[Ganter]

$V$ : a vector bundle over  $\Lambda(X // G)$ .

$P_n(V) := V^{\hat{\otimes}_{\mathbb{Z}[q^\pm]} n}$  defines an operation

$$P_n : QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n})$$

$$\mathbb{P}_n = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} \mathbb{P}_{(\underline{g}, \sigma)} :$$

$$QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n}) = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)}).$$

$$\mathbb{P}_{(\underline{g}, \sigma)} : QEll_G(X) \xrightarrow{U^*} K_{orb}(\Lambda_{(\underline{g}, \sigma)}(X)) \xrightarrow{(\ )_k^\wedge} K_{orb}(\Lambda_{(\underline{g}, \sigma)}^{var}(X))$$

$$\xrightarrow{\boxtimes} K_{orb}(d_{(\underline{g}, \sigma)}(X)) \xrightarrow{f_{(\underline{g}, \sigma)}^*} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)})$$

# Power Operation

Quasi-elliptic cohomology has power operations, which gives it the structure of an " $H_\infty$ -ring theory" [Ganter 06].

## Atiyah's Power Operation

[Ganter]

$V$ : a vector bundle over  $\Lambda(X // G)$ .

$P_n(V) := V^{\hat{\otimes}_{\mathbb{Z}[q^\pm]} }^n$  defines an operation

$$P_n : QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n})$$

$$\mathbb{P}_n = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} \mathbb{P}_{(\underline{g}, \sigma)} :$$


$$QEll_G(X) \longrightarrow QEll_{G \wr \Sigma_n}(X^{\times n}) = \prod_{(\underline{g}, \sigma) \in (G \wr \Sigma_n)_{conj}^{tors}} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)}).$$

$$\begin{aligned} \mathbb{P}_{(\underline{g}, \sigma)} : QEll_G(X) &\xrightarrow{U^*} K_{orb}(\Lambda_{(\underline{g}, \sigma)}(X)) \xrightarrow{(\ )_k^\wedge} K_{orb}(\Lambda_{(\underline{g}, \sigma)}^{var}(X)) \\ &\xrightarrow{\boxtimes} K_{orb}(d_{(\underline{g}, \sigma)}(X)) \xrightarrow{f_{(\underline{g}, \sigma)}^*} K_{\Lambda_{G \wr \Sigma_n}(\underline{g}, \sigma)}((X^{\times n})^{(\underline{g}, \sigma)}) \end{aligned}$$

# the functor $f_{(\underline{g}, \sigma)}$ : the KEY isomorphism

$\mathcal{L}_{(\underline{g}, \sigma)}(X^{\times n})$  and  $\prod_k \prod_{(i_1, \dots, i_k)} {}^k \mathcal{L}_{g_{i_k} \dots g_{i_1}} X$  are  $\Lambda_{G|\Sigma_n}(\underline{g}, \sigma)$ -equivariant homeomorphic.

Example  $(\mathcal{L}_{(g_1, \dots, g_5, (135)(24))}(X^{\times 5})$  and  ${}_3 \mathcal{L}_{g_5 g_3 g_1}(X) \times {}_2 \mathcal{L}_{g_4 g_2}(X)$ )




This picture is from [Ganter].

$f_{(\underline{g}, \sigma)}$  is the restriction of this homeomorphism to the constant loops.

# the functor $f_{(\underline{g}, \sigma)}$ : the KEY isomorphism

$\mathcal{L}_{(\underline{g}, \sigma)}(X^{\times n})$  and  $\prod_k \prod_{(i_1, \dots, i_k)} {}^k \mathcal{L}_{g_{i_k} \dots g_{i_1}} X$  are  $\Lambda_{G|\Sigma_n}(\underline{g}, \sigma)$ -equivariant homeomorphic.

Example  $(\mathcal{L}_{(g_1, \dots, g_5, (135)(24))}(X^{\times 5})$  and  ${}_3 \mathcal{L}_{g_5 g_3 g_1}(X) \times {}_2 \mathcal{L}_{g_4 g_2}(X)$ )




This picture is from [Ganter].

$f_{(\underline{g}, \sigma)}$  is the restriction of this homeomorphism to the constant loops.

# the functor $f_{(\underline{g}, \sigma)}$ : the KEY isomorphism

$\mathcal{L}_{(\underline{g}, \sigma)}(X^{\times n})$  and  $\prod_k \prod_{(i_1, \dots, i_k)} {}^k \mathcal{L}_{g_{i_k} \dots g_{i_1}} X$  are  $\Lambda_{G|\Sigma_n}(\underline{g}, \sigma)$ -equivariant homeomorphic.

Example  $(\mathcal{L}_{(g_1, \dots, g_5, (135)(24))}(X^{\times 5})$  and  ${}_3 \mathcal{L}_{g_5 g_3 g_1}(X) \times {}_2 \mathcal{L}_{g_4 g_2}(X)$ )



This picture is from [Ganter].

$f_{(\underline{g}, \sigma)}$  is the restriction of this homeomorphism to the constant loops.

$$C_G(g, g') := \{x \in G \mid gx = xg'\}.$$

$$\Lambda_G^k(g, g') := C_G(g, g') \times \mathbb{R} / (x, t) \sim (gx, t - k).$$

Note

$$\Lambda_G^1(g, g) = \Lambda_G(g).$$

A groupoid equivalent to  $\Lambda(X//G)$

- objects  $\coprod_{g \in G^{\text{tors}}} X^g$ ;
- morphisms  $\coprod_{g, g' \in G^{\text{tors}}} \Lambda_G^1(g, g') \times X^g$

Explicitly,  $(\alpha, x)$  maps  $x \in X^g$  to  $\alpha \cdot x \in X^{g'}$ .

We use the same symbol  $\Lambda(X//G)$  to denote it.

$$C_G(g, g') := \{x \in G \mid gx = xg'\}.$$

$$\Lambda_G^k(g, g') := C_G(g, g') \times \mathbb{R} / (x, t) \sim (gx, t - k).$$

Note

$$\Lambda_G^1(g, g) = \Lambda_G(g).$$

A groupoid equivalent to  $\Lambda(X//G)$

- objects  $\coprod_{g \in G^{\text{tors}}} X^g$ ;
- morphisms  $\coprod_{g, g' \in G^{\text{tors}}} \Lambda_G^1(g, g') \times X^g$

Explicitly,  $(\alpha, x)$  maps  $x \in X^g$  to  $\alpha \cdot x \in X^{g'}$ .

We use the same symbol  $\Lambda(X//G)$  to denote it.



$$C_G(g, g') := \{x \in G \mid gx = xg'\}.$$

$$\Lambda_G^k(g, g') := C_G(g, g') \times \mathbb{R} / (x, t) \sim (gx, t - k).$$

Note

$$\Lambda_G^1(g, g) = \Lambda_G(g).$$

A groupoid equivalent to  $\Lambda(X // G)$

- objects  $\coprod_{g \in G^{tors}} X^g$ ;
- morphisms  $\coprod_{g, g' \in G^{tors}} \Lambda_G^1(g, g') \times X^g$

Explicitly,  $(\alpha, x)$  maps  $x \in X^g$  to  $\alpha \cdot x \in X^{g'}$ .

We use the same symbol  $\Lambda(X // G)$  to denote it.

# the Functor $U$ : Draw necessary information

For  $(\underline{g}, \sigma) \in G \wr \Sigma_n$ , let  $\Lambda_{(\underline{g}, \sigma)}(X)$  denote the groupoid with

- objects: points in  $\coprod_k \coprod_{(i_1, \dots, i_k)} X^{g_{i_k} \cdots g_{i_1}}$   
where  $(i_1, \dots, i_k)$  goes over all the  $k$ -cycles of  $\sigma$ ;
- morphisms:  $\coprod_k \coprod_{(i_1, \dots, i_k), (j_1, \dots, j_k)} \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ ,  
Explicitly,  $(\alpha, x)$  maps  $x$  to  $\alpha \cdot x$ .

Let

$$U : \Lambda_{(\underline{g}, \sigma)}(X) \longrightarrow \Lambda(X // G)$$

denote the functor sending

- an object  $x \in X^{g_{i_k} \cdots g_{i_1}} \implies x \in X^{g_{i_k} \cdots g_{i_1}}$ ;
- a morphism  $(\alpha, x) \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}} \implies (\alpha, x) \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ .

Note:  $\Lambda_{(\underline{g}, \sigma)}(X)$  may not be a subgroupoid of  $\Lambda(X // G)$ .

# the Functor $U$ : Draw necessary information

For  $(\underline{g}, \sigma) \in G \wr \Sigma_n$ , let  $\Lambda_{(\underline{g}, \sigma)}(X)$  denote the groupoid with

- objects: points in  $\coprod_k \coprod_{(i_1, \dots, i_k)} X^{g_{i_k} \cdots g_{i_1}}$   
where  $(i_1, \dots, i_k)$  goes over all the  $k$ -cycles of  $\sigma$ ;
- morphisms:  $\coprod_k \coprod_{(i_1, \dots, i_k), (j_1, \dots, j_k)} \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ ,  
Explicitly,  $(\alpha, x)$  maps  $x$  to  $\alpha \cdot x$ .

Let

$$U : \Lambda_{(\underline{g}, \sigma)}(X) \longrightarrow \Lambda(X // G)$$

denote the functor sending

- an object  $x \in X^{g_{i_k} \cdots g_{i_1}} \implies x \in X^{g_{i_k} \cdots g_{i_1}}$ ;
- a morphism  $(\alpha, x) \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}} \implies (\alpha, x) \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ .

Note:  $\Lambda_{(\underline{g}, \sigma)}(X)$  may not be a subgroupoid of  $\Lambda(X // G)$ .

# the Functor $U$ : Draw necessary information

For  $(\underline{g}, \sigma) \in G \wr \Sigma_n$ , let  $\Lambda_{(\underline{g}, \sigma)}(X)$  denote the groupoid with

- objects: points in  $\coprod_k \coprod_{(i_1, \dots, i_k)} X^{g_{i_k} \cdots g_{i_1}}$   
where  $(i_1, \dots, i_k)$  goes over all the  $k$ -cycles of  $\sigma$ ;
- morphisms:  $\coprod_k \coprod_{(i_1, \dots, i_k), (j_1, \dots, j_k)} \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ ,  
Explicitly,  $(\alpha, x)$  maps  $x$  to  $\alpha \cdot x$ .

Let

$$U : \Lambda_{(\underline{g}, \sigma)}(X) \longrightarrow \Lambda(X // G)$$

denote the functor sending

- an object  $x \in X^{g_{i_k} \cdots g_{i_1}} \implies x \in X^{g_{i_k} \cdots g_{i_1}}$ ;
- a morphism  $(\alpha, x) \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}} \implies (\alpha, x) \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ .

Note:  $\Lambda_{(\underline{g}, \sigma)}(X)$  may not be a subgroupoid of  $\Lambda(X // G)$ .

Let  $\Lambda_{(\underline{g}, \sigma)}^{var}(X)$  be the groupoid with

- the same objects as  $\Lambda_{(\underline{g}, \sigma)}(X)$
- morphisms:  $\coprod_k \coprod_{(i_1, \dots, i_k), (j_1, \dots, j_k)} \Lambda_G^k(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ ,  
where  $(i_1, \dots, i_k)$  and  $(j_1, \dots, j_k)$  go over all the  $k$ -cycles of  $\sigma$ .

The functor

$$(\ )_k^\wedge : \Lambda_{(\underline{g}, \sigma)}^{var}(X) \longrightarrow \Lambda_{(\underline{g}, \sigma)}(X)$$

- identity on objects
- sends each  $[g, t] \in \Lambda_G^k(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1})$  to  $[g, \frac{t}{k}] \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1})$ .

Let  $\Lambda_{(\underline{g}, \sigma)}^{var}(X)$  be the groupoid with

- the same objects as  $\Lambda_{(\underline{g}, \sigma)}(X)$
- morphisms:  $\coprod_k \coprod_{(i_1, \dots, i_k), (j_1, \dots, j_k)} \Lambda_G^k(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1}) \times X^{g_{i_k} \cdots g_{i_1}}$ ,  
where  $(i_1, \dots, i_k)$  and  $(j_1, \dots, j_k)$  go over all the  $k$ -cycles of  $\sigma$ .

The functor

$$(\ )_k^\wedge : \Lambda_{(\underline{g}, \sigma)}^{var}(X) \longrightarrow \Lambda_{(\underline{g}, \sigma)}(X)$$

- identity on objects
- sends each  $[g, t] \in \Lambda_G^k(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1})$  to  $[g, \frac{t}{k}] \in \Lambda_G(g_{i_k} \cdots g_{i_1}, g_{j_k} \cdots g_{j_1})$ .

## Example ( $G = e$ )

$QEII_G^*(X) = K_{\mathbb{T}}^*(X)$ . For each  $\sigma \in \Sigma_n$ ,  $\mathbb{P}_{(\underline{1}, \sigma)}(x) = \boxtimes_k \boxtimes_{(i_1, \dots, i_k)} (x)_k$ .

When  $n = 2$ ,

$$QEII_{\Sigma_2}(X \times X) \cong K(X \times X)[q^{\pm}][1, s]/(s^2 - 1) \times K(X)[q^{\pm}][y]/(y^2 - q)$$

$$\mathbb{P}_2(x) = (\mathbb{P}_{(\underline{1}, (1)(1))}(x), \mathbb{P}_{(\underline{1}, (12))}(x)) = (x \boxtimes x, (x)_2).$$

When  $n = 3$ ,  $\mathbb{P}_3(x) = (\mathbb{P}_{(\underline{1}, (1)(1)(1))}(x), \mathbb{P}_{(\underline{1}, (12)(1))}(x), \mathbb{P}_{(\underline{1}, (123))}(x)) = (x \boxtimes x \boxtimes x, (x)_2 \boxtimes x, (x)_3).$

## A Ring Homomorphism

$$\begin{aligned} \bar{P}_N : QEII_G(X) &\xrightarrow{\mathbb{P}_N} QEII_{G \wr \Sigma_N}(X^{\times N}) \xrightarrow{res} QEII_{G \times \Sigma_N}(X^{\times N}) \xrightarrow{diag^*} \\ QEII_{G \times \Sigma_N}(X) &\cong QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(pt) \longrightarrow \\ QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(pt) &/ \mathcal{I}_{tr}^{QEII} \end{aligned}$$

- analogous to the Adams operations of equivariant K-theories.
- but different and new.

## Example ( $G = e$ )

$QEII_G^*(X) = K_{\mathbb{T}}^*(X)$ . For each  $\sigma \in \Sigma_n$ ,  $\mathbb{P}_{(\underline{1}, \sigma)}(x) = \boxtimes_k \boxtimes_{(i_1, \dots, i_k)} (x)_k$ .  
When  $n = 2$ ,

$$QEII_{\Sigma_2}(X \times X) \cong K(X \times X)[q^{\pm}][1, s]/(s^2 - 1) \times K(X)[q^{\pm}][y]/(y^2 - q)$$

$$\mathbb{P}_2(x) = (\mathbb{P}_{(\underline{1}, (1)(1))}(x), \mathbb{P}_{(\underline{1}, (12))}(x)) = (x \boxtimes x, (x)_2).$$

When  $n = 3$ ,  $\mathbb{P}_3(x) = (\mathbb{P}_{(\underline{1}, (1)(1)(1))}(x), \mathbb{P}_{(\underline{1}, (12)(1))}(x), \mathbb{P}_{(\underline{1}, (123))}(x)) = (x \boxtimes x \boxtimes x, (x)_2 \boxtimes x, (x)_3).$

## A Ring Homomorphism

$$\begin{aligned} \bar{P}_N : QEII_G(X) &\xrightarrow{\mathbb{P}_N} QEII_{G \wr \Sigma_N}(X^{\times N}) \xrightarrow{res} QEII_{G \times \Sigma_N}(X^{\times N}) \xrightarrow{diag^*} \\ QEII_{G \times \Sigma_N}(X) &\cong QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(pt) \longrightarrow \\ QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(pt) &/ \mathcal{I}_{tr}^{QEII} \end{aligned}$$

- analogous to the Adams operations of equivariant K-theories.
- but different and new.



## Example ( $G = e$ )

$QEll_G^*(X) = K_{\mathbb{T}}^*(X)$ . For each  $\sigma \in \Sigma_n$ ,  $\mathbb{P}_{(\underline{1}, \sigma)}(x) = \boxtimes_k \boxtimes_{(i_1, \dots, i_k)} (x)_k$ .  
When  $n = 2$ ,

$$QEll_{\Sigma_2}(X \times X) \cong K(X \times X)[q^{\pm}][1, s]/(s^2 - 1) \times K(X)[q^{\pm}][y]/(y^2 - q)$$

$$\mathbb{P}_2(x) = (\mathbb{P}_{(\underline{1}, (1)(1))}(x), \mathbb{P}_{(\underline{1}, (12))}(x)) = (x \boxtimes x, (x)_2).$$

When  $n = 3$ ,  $\mathbb{P}_3(x) = (\mathbb{P}_{(\underline{1}, (1)(1)(1))}(x), \mathbb{P}_{(\underline{1}, (12)(1))}(x), \mathbb{P}_{(\underline{1}, (123))}(x)) = (x \boxtimes x \boxtimes x, (x)_2 \boxtimes x, (x)_3).$

## A Ring Homomorphism

$$\begin{aligned} \bar{P}_N : QEll_G(X) &\xrightarrow{\mathbb{P}_N} QEll_{G \wr \Sigma_N}(X^{\times N}) \xrightarrow{res} QEll_{G \times \Sigma_N}(X^{\times N}) \xrightarrow{diag^*} \\ &QEll_{G \times \Sigma_N}(X) \cong QEll_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEll_{\Sigma_N}(pt) \longrightarrow \\ &QEll_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEll_{\Sigma_N}(pt) / \mathcal{I}_{tr}^{QEll} \end{aligned}$$

- analogous to the Adams operations of equivariant K-theories.
- but different and new.

## Example ( $G = e$ )

$QEII_G^*(X) = K_{\mathbb{T}}^*(X)$ . For each  $\sigma \in \Sigma_n$ ,  $\mathbb{P}_{(\underline{1}, \sigma)}(x) = \boxtimes_k \boxtimes_{(i_1, \dots, i_k)} (x)_k$ .  
When  $n = 2$ ,

$$QEII_{\Sigma_2}(X \times X) \cong K(X \times X)[q^{\pm}][1, s]/(s^2 - 1) \times K(X)[q^{\pm}][y]/(y^2 - q)$$

$$\mathbb{P}_2(x) = (\mathbb{P}_{(\underline{1}, (1)(1))}(x), \mathbb{P}_{(\underline{1}, (12))}(x)) = (x \boxtimes x, (x)_2).$$

When  $n = 3$ ,  $\mathbb{P}_3(x) = (\mathbb{P}_{(\underline{1}, (1)(1)(1))}(x), \mathbb{P}_{(\underline{1}, (12)(1))}(x), \mathbb{P}_{(\underline{1}, (123))}(x)) = (x \boxtimes x \boxtimes x, (x)_2 \boxtimes x, (x)_3).$

## A Ring Homomorphism

$$\begin{aligned} \bar{P}_N : QEII_G(X) &\xrightarrow{\mathbb{P}_N} QEII_{G \wr \Sigma_N}(X^{\times N}) \xrightarrow{res} QEII_{G \times \Sigma_N}(X^{\times N}) \xrightarrow{diag^*} \\ QEII_{G \times \Sigma_N}(X) &\cong QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(\text{pt}) \longrightarrow \\ QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(\text{pt}) &/ \mathcal{I}_{tr}^{QEII} \end{aligned}$$

- analogous to the Adams operations of equivariant K-theories.
- but different and new.

## Example ( $G = e$ )

$QEII_G^*(X) = K_{\mathbb{T}}^*(X)$ . For each  $\sigma \in \Sigma_n$ ,  $\mathbb{P}_{(\underline{1}, \sigma)}(x) = \boxtimes_k \boxtimes_{(i_1, \dots, i_k)} (x)_k$ .  
When  $n = 2$ ,

$$QEII_{\Sigma_2}(X \times X) \cong K(X \times X)[q^{\pm}][1, s]/(s^2 - 1) \times K(X)[q^{\pm}][y]/(y^2 - q)$$

$$\mathbb{P}_2(x) = (\mathbb{P}_{(\underline{1}, (1)(1))}(x), \mathbb{P}_{(\underline{1}, (12))}(x)) = (x \boxtimes x, (x)_2).$$

When  $n = 3$ ,  $\mathbb{P}_3(x) = (\mathbb{P}_{(\underline{1}, (1)(1)(1))}(x), \mathbb{P}_{(\underline{1}, (12)(1))}(x), \mathbb{P}_{(\underline{1}, (123))}(x)) = (x \boxtimes x \boxtimes x, (x)_2 \boxtimes x, (x)_3).$

## A Ring Homomorphism

$$\begin{aligned} \bar{P}_N : QEII_G(X) &\xrightarrow{\mathbb{P}_N} QEII_{G \wr \Sigma_N}(X^{\times N}) \xrightarrow{res} QEII_{G \times \Sigma_N}(X^{\times N}) \xrightarrow{diag^*} \\ QEII_{G \times \Sigma_N}(X) &\cong QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(pt) \longrightarrow \\ QEII_G(X) \otimes_{\mathbb{Z}[q^{\pm}]} QEII_{\Sigma_N}(pt) &/ \mathcal{I}_{tr}^{QEII} \end{aligned}$$

- analogous to the Adams operations of equivariant K-theories.
- but different and new.

# Finite Subgroups of Elliptic Curve

## Question

Can the classification of the finite subgroups of an elliptic curve be given by the associated elliptic cohomology theory?

## Morava E-theory

[Strickland]

The Morava  $E$ -theory of the symmetric group  $\Sigma_n$  modulo a certain transfer ideal classifies the power subgroups of rank  $n$  of the formal group  $\mathbb{G}_E$ .

## Generalized Morava $E$ -theories

[Schlank and Stapleton]

They generalized Strickland's result to generalized Morava  $E$ -theories  $E_G(\mathcal{L}^h(-))$  using Stapleton's transchromatic character theory.

## Question

What about generalized elliptic cohomology theories?

# Finite Subgroups of Elliptic Curve

## Question

Can the classification of the finite subgroups of an elliptic curve be given by the associated elliptic cohomology theory?

## Morava E-theory

[Strickland]

The Morava  $E$ -theory of the symmetric group  $\Sigma_n$  modulo a certain transfer ideal classifies the power subgroups of rank  $n$  of the formal group  $\mathbb{G}_E$ .

## Generalized Morava $E$ -theories

[Schlank and Stapleton]

They generalized Strickland's result to generalized Morava  $E$ -theories  $E_G(\mathcal{L}^h(-))$  using Stapleton's transchromatic character theory.

## Question

What about generalized elliptic cohomology theories?

# Finite Subgroups of Elliptic Curve

## Question

Can the classification of the finite subgroups of an elliptic curve be given by the associated elliptic cohomology theory?

## Morava E-theory

[Strickland]

The Morava  $E$ -theory of the symmetric group  $\Sigma_n$  modulo a certain transfer ideal classifies the power subgroups of rank  $n$  of the formal group  $\mathbb{G}_E$ .

## Generalized Morava $E$ -theories

[Schlank and Stapleton]

They generalized Strickland's result to generalized Morava  $E$ -theories  $E_G(\mathcal{L}^h(-))$  using Stapleton's transchromatic character theory.

## Question

What about generalized elliptic cohomology theories?

# Finite Subgroups of Elliptic Curve

## Question

Can the classification of the finite subgroups of an elliptic curve be given by the associated elliptic cohomology theory?

## Morava E-theory

[Strickland]

The Morava  $E$ -theory of the symmetric group  $\Sigma_n$  modulo a certain transfer ideal classifies the power subgroups of rank  $n$  of the formal group  $\mathbb{G}_E$ .

## Generalized Morava $E$ -theories

[Schlank and Stapleton]

They generalized Strickland's result to generalized Morava  $E$ -theories  $E_G(\mathcal{L}^h(-))$  using Stapleton's transchromatic character theory.

## Question

What about generalized elliptic cohomology theories?

$$E_q : y^2 + xy = x^3 + a_4x + a_6$$

$$a_4 = -5 \sum_{n \geq 1} n^3 q^n / (1 - q^n) \quad a_6 = -\frac{1}{12} \sum_{n \geq 1} (7n^5 + 5n^3) q^n / (1 - q^n).$$

$N$ -division points

$$Tate(q)[N] \cong \prod_{k=0}^{N-1} \text{Spec}(\mathbb{Z}((q))[x]/(x^N - q^k)).$$

$$\text{Spec}(QEll_{\mathbb{Z}/N}(pt)) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong Tate(q)[N].$$

Proposition

*The finite subgroups of the Tate curve are the kernels of isogenies.  
Its finite subgroups of order  $N$  can be classified by*

$$\mathcal{O}_{Sub_N} = \prod_{N=de} \mathbb{Z}((q))[q_s'^{\pm}] / \langle q^d - q_s'^e \rangle.$$



$$E_q : y^2 + xy = x^3 + a_4x + a_6$$

$$a_4 = -5 \sum_{n \geq 1} n^3 q^n / (1 - q^n) \quad a_6 = -\frac{1}{12} \sum_{n \geq 1} (7n^5 + 5n^3) q^n / (1 - q^n).$$

### $N$ -division points

$$Tate(q)[N] \cong \prod_{k=0}^{N-1} \text{Spec}(\mathbb{Z}((q))[x]/(x^N - q^k)).$$

$$\text{Spec}(QEll_{\mathbb{Z}/N}(pt)) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong Tate(q)[N].$$

### Proposition

*The finite subgroups of the Tate curve are the kernels of isogenies.  
Its finite subgroups of order  $N$  can be classified by*

$$\mathcal{O}_{Sub_N} = \prod_{N=de} \mathbb{Z}((q))[q_s'^{\pm}] / \langle q^d - q_s'^e \rangle.$$

$$E_q : y^2 + xy = x^3 + a_4x + a_6$$

$$a_4 = -5 \sum_{n \geq 1} n^3 q^n / (1 - q^n) \quad a_6 = -\frac{1}{12} \sum_{n \geq 1} (7n^5 + 5n^3) q^n / (1 - q^n).$$

### $N$ -division points

$$Tate(q)[N] \cong \prod_{k=0}^{N-1} \text{Spec}(\mathbb{Z}((q))[x]/(x^N - q^k)).$$

$$\text{Spec}(QEll_{\mathbb{Z}/N}(pt)) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong Tate(q)[N].$$

### Proposition

*The finite subgroups of the Tate curve are the kernels of isogenies.  
Its finite subgroups of order  $N$  can be classified by*

$$\mathcal{O}_{Sub_N} = \prod_{N=de} \mathbb{Z}((q))[q_s'^{\pm}] / \langle q^d - q_s'^e \rangle.$$

$$E_q : y^2 + xy = x^3 + a_4x + a_6$$

$$a_4 = -5 \sum_{n \geq 1} n^3 q^n / (1 - q^n) \quad a_6 = -\frac{1}{12} \sum_{n \geq 1} (7n^5 + 5n^3) q^n / (1 - q^n).$$

### $N$ -division points

$$Tate(q)[N] \cong \coprod_{k=0}^{N-1} \text{Spec}(\mathbb{Z}((q))[x]/(x^N - q^k)).$$

$$\text{Spec}(QEll_{\mathbb{Z}/N}(pt)) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong Tate(q)[N].$$

### Proposition

*The finite subgroups of the Tate curve are the kernels of isogenies.  
Its finite subgroups of order  $N$  can be classified by*

$$\mathcal{O}_{Sub_N} = \prod_{N=de} \mathbb{Z}((q))[q_s'^{\pm}] / \langle q^d - q_s'^e \rangle.$$

$$E_q : y^2 + xy = x^3 + a_4x + a_6$$

$$a_4 = -5 \sum_{n \geq 1} n^3 q^n / (1 - q^n) \quad a_6 = -\frac{1}{12} \sum_{n \geq 1} (7n^5 + 5n^3) q^n / (1 - q^n).$$

### $N$ -division points

$$Tate(q)[N] \cong \prod_{k=0}^{N-1} \text{Spec}(\mathbb{Z}((q))[x]/(x^N - q^k)).$$

$$\text{Spec}(QE_{\mathbb{Z}/N}(pt)) \otimes_{\mathbb{Z}[q^{\pm}]} \mathbb{Z}((q)) \cong Tate(q)[N].$$

### Proposition

*The finite subgroups of the Tate curve are the kernels of isogenies.  
Its finite subgroups of order  $N$  can be classified by*

$$\mathcal{O}_{Sub_N} = \prod_{N=de} \mathbb{Z}((q))[q_s'^{\pm}] / \langle q^d - q_s'^e \rangle.$$

$$I_{tr}^{Tate} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[I_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : K_{Tate}(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow K_{Tate}(\text{pt} // \Sigma_N)]$$

$$\mathcal{I}_{tr}^{QEII} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[\mathcal{I}_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : QEII(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow QEII(\text{pt} // \Sigma_N)]$$

Example ( $N = 2$  and  $X = \text{pt}$ )

$$\mathcal{I}_{tr}^{QEII} = \text{Ind}_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}^{\Lambda_{\Sigma_2}(1)} K_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}(\text{pt}).$$

$$\begin{aligned} QEII(\text{pt} // \Sigma_2) / \mathcal{I}_{tr}^{QEII} &= K_{\Lambda_{\Sigma_2}(1)}(\text{pt}) / \mathcal{I}_{tr}^{QEII} \times K_{\Lambda_{\Sigma_2}(12)}(\text{pt}) \\ &\cong \mathbb{Z}[q^{\pm}][q'] / (q' - q^2) \times \mathbb{Z}[q^{\pm}][q''] / (q''^2 - q). \end{aligned}$$

Note  $\mathbb{P}_2(q) = (q^2, q^{\frac{1}{2}}) = (q', q'')$ .

$$I_{tr}^{Tate} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[I_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : K_{Tate}(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow K_{Tate}(\text{pt} // \Sigma_N)]$$

$$\mathcal{I}_{tr}^{QEII} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[\mathcal{I}_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : QEII(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow QEII(\text{pt} // \Sigma_N)]$$

Example ( $N = 2$  and  $X = \text{pt}$ )

$$\mathcal{I}_{tr}^{QEII} = \text{Ind}_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}^{\Lambda_{\Sigma_2}(1)} K_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}(\text{pt}).$$

$$\begin{aligned} QEII(\text{pt} // \Sigma_2) / \mathcal{I}_{tr}^{QEII} &= K_{\Lambda_{\Sigma_2}(1)}(\text{pt}) / \mathcal{I}_{tr}^{QEII} \times K_{\Lambda_{\Sigma_2}(12)}(\text{pt}) \\ &\cong \mathbb{Z}[q^{\pm}][q'] / (q' - q^2) \times \mathbb{Z}[q^{\pm}][q''] / (q''^2 - q). \end{aligned}$$

Note  $\mathbb{P}_2(q) = (q^2, q^{\frac{1}{2}}) = (q', q'')$ .

$$I_{tr}^{Tate} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[I_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : K_{Tate}(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow K_{Tate}(\text{pt} // \Sigma_N)]$$

$$\mathcal{I}_{tr}^{QEII} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[\mathcal{I}_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : QEII(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow QEII(\text{pt} // \Sigma_N)]$$

Example ( $N = 2$  and  $X = \text{pt}$ )

$$\mathcal{I}_{tr}^{QEII} = \text{Ind}_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}^{\Lambda_{\Sigma_2}(1)} K_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}(\text{pt}).$$

$$\begin{aligned} QEII(\text{pt} // \Sigma_2) / \mathcal{I}_{tr}^{QEII} &= K_{\Lambda_{\Sigma_2}(1)}(\text{pt}) / \mathcal{I}_{tr}^{QEII} \times K_{\Lambda_{\Sigma_2}(12)}(\text{pt}) \\ &\cong \mathbb{Z}[q^{\pm}][q'] / (q' - q^2) \times \mathbb{Z}[q^{\pm}][q''] / (q''^2 - q). \end{aligned}$$

Note  $\mathbb{P}_2(q) = (q^2, q^{\frac{1}{2}}) = (q', q'')$ .

$$I_{tr}^{Tate} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[I_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : K_{Tate}(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow K_{Tate}(\text{pt} // \Sigma_N)]$$

$$\mathcal{I}_{tr}^{QEII} := \sum_{\substack{i+j=N, \\ N>j>0}} \text{Image}[\mathcal{I}_{\Sigma_i \times \Sigma_j}^{\Sigma_N} : QEII(\text{pt} // \Sigma_i \times \Sigma_j) \longrightarrow QEII(\text{pt} // \Sigma_N)]$$

Example ( $N = 2$  and  $X = \text{pt}$ )

$$\mathcal{I}_{tr}^{QEII} = \text{Ind}_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}^{\Lambda_{\Sigma_2}(1)} K_{\Lambda_{\Sigma_1 \times \Sigma_1}(1)}(\text{pt}).$$

$$\begin{aligned} QEII(\text{pt} // \Sigma_2) / \mathcal{I}_{tr}^{QEII} &= K_{\Lambda_{\Sigma_2}(1)}(\text{pt}) / \mathcal{I}_{tr}^{QEII} \times K_{\Lambda_{\Sigma_2}(12)}(\text{pt}) \\ &\cong \mathbb{Z}[q^{\pm}][q'] / (q' - q^2) \times \mathbb{Z}[q^{\pm}][q''] / (q''^2 - q). \end{aligned}$$

Note  $\mathbb{P}_2(q) = (q^2, q^{\frac{1}{2}}) = (q', q'')$ .



# Classification of Finite Subgroups of Tate Curve

## Theorem (Huan)

$$QEII(pt//\Sigma_N)/\mathcal{I}_{tr}^{QEII} \cong \prod_{N=de} \mathbb{Z}[q^{\pm}][q'^{\pm}]/\langle q^d - q'^e \rangle,$$

where  $\mathcal{I}_{tr}^{QEII}$  is the transfer ideal and  $q'$  is the image of  $q$  under the power operation  $\mathbb{P}_N$ . The product goes over all the ordered pairs of positive integers  $(d, e)$  such that  $N = de$ .

## Theorem (Huan)

*The Tate K-theory of symmetric groups modulo a certain transfer ideal classifies finite subgroups of the Tate curve.*

$$K_{Tate}(pt//\Sigma_N)/I_{tr}^{Tate} \cong \prod_{N=de} \mathbb{Z}((q))[q'_s{}^{\pm}]/\langle q^d - q'_s{}^e \rangle,$$

where  $I_{tr}^{Tate}$  is the transfer ideal and  $q'_s$  is the image of  $q$  under the stringy power operation  $P_N^{string}$ , the product goes over all the ordered pairs of positive integers  $(d, e)$  such that  $N = de$ .

# Classification of Finite Subgroups of Tate Curve

## Theorem (Huan)

$$QEll(pt//\Sigma_N)/\mathcal{I}_{tr}^{QEll} \cong \prod_{N=de} \mathbb{Z}[q^{\pm}][q'^{\pm}]/\langle q^d - q'^e \rangle,$$

where  $\mathcal{I}_{tr}^{QEll}$  is the transfer ideal and  $q'$  is the image of  $q$  under the power operation  $\mathbb{P}_N$ . The product goes over all the ordered pairs of positive integers  $(d, e)$  such that  $N = de$ .

## Theorem (Huan)

The Tate  $K$ -theory of symmetric groups modulo a certain transfer ideal classifies finite subgroups of the Tate curve.

$$K_{Tate}(pt//\Sigma_N)/I_{tr}^{Tate} \cong \prod_{N=de} \mathbb{Z}((q))[q'_s{}^{\pm}]/\langle q^d - q'_s{}^e \rangle,$$

where  $I_{tr}^{Tate}$  is the transfer ideal and  $q'_s$  is the image of  $q$  under the stringy power operation  $P_N^{string}$ , the product goes over all the ordered pairs of positive integers  $(d, e)$  such that  $N = de$ .

- 1967 Daniel G. Quillen "Homotopical algebra": form the foundation of homotopy theory.
- 1997 Vladimir Voevodsky "The Milnor Conjecture": apply homotopical thinking to non-homotopical situation.

Motivating example: Quillen model structure on topological spaces

Represent the standard homotopy theory of CW-complexes.

- weak homotopy equivalence
- Serre fibration
- retract of relative cell complex

Morphisms

Quillen adjunction:  $(L \dashv R) : \mathcal{C} \overset{R}{\underset{L}{\rightleftarrows}} \mathcal{D}.$

Quillen equivalence:  $Ho(\mathcal{C}) \overset{\mathbb{R}}{\underset{\mathbb{L}}{\rightleftarrows}} Ho(\mathcal{D}).$

- 1967 Daniel G. Quillen "Homotopical algebra": form the foundation of homotopy theory.
- 1997 Vladimir Voevodsky "The Milnor Conjecture": apply homotopical thinking to non-homotopical situation.

Definition: A model structure on a category  $\mathcal{C}$

Three distinguished classes of morphisms:

- Weak Equivalence
- Fibration
- Cofibration

satisfying the axioms: • Retracts • 2 of 3 • Lifting • Factorization

Morphisms

Quillen adjunction:  $(L \dashv R) : \mathcal{C} \xrightleftharpoons[L]{R} \mathcal{D}$ .

Quillen equivalence:  $Ho(\mathcal{C}) \xrightleftharpoons[\mathbb{L}]{\mathbb{R}} Ho(\mathcal{D})$ .

- 1967 Daniel G. Quillen "Homotopical algebra": form the foundation of homotopy theory.
- 1997 Vladimir Voevodsky "The Milnor Conjecture": apply homotopical thinking to non-homotopical situation.

Definition: A model structure on a category  $\mathcal{C}$

Three distinguished classes of morphisms:

- Weak Equivalence
- Fibration
- Cofibration

satisfying the axioms: • Retracts • 2 of 3 • Lifting • Factorization

Morphisms

Quillen adjunction:  $(L \dashv R) : \mathcal{C} \xrightleftharpoons[L]{R} \mathcal{D}.$

Quillen equivalence:  $Ho(\mathcal{C}) \xrightleftharpoons[\mathbb{L}]{\mathbb{R}} Ho(\mathcal{D}).$

- 1967 Daniel G. Quillen "Homotopical algebra": form the foundation of homotopy theory.
- 1997 Vladimir Voevodsky "The Milnor Conjecture": apply homotopical thinking to non-homotopical situation.

**Definition:** A model structure on a category  $\mathcal{C}$

Three distinguished classes of morphisms:

- Weak Equivalence
- Fibration
- Cofibration

satisfying the axioms: • Retracts • 2 of 3 • Lifting • Factorization

## Morphisms

Quillen adjunction:  $(L \dashv R) : \mathcal{C} \begin{smallmatrix} \xleftarrow{R} \\ \xrightarrow{L} \end{smallmatrix} \mathcal{D}.$

Quillen equivalence:  $Ho(\mathcal{C}) \begin{smallmatrix} \xleftarrow{R} \\ \xrightarrow{L} \end{smallmatrix} Ho(\mathcal{D}).$

- 1967 Daniel G. Quillen "Homotopical algebra": form the foundation of homotopy theory.
- 1997 Vladimir Voevodsky "The Milnor Conjecture": apply homotopical thinking to non-homotopical situation.

**Definition:** A model structure on a category  $\mathcal{C}$

Three distinguished classes of morphisms:

- Weak Equivalence
- Fibration
- Cofibration

satisfying the axioms: • Retracts • 2 of 3 • Lifting • Factorization

## Morphisms

Quillen adjunction:  $(L \dashv R) : \mathcal{C} \xrightleftharpoons[L]{R} \mathcal{D}$ .

Quillen equivalence:  $Ho(\mathcal{C}) \xrightleftharpoons[\mathbb{L}]{\mathbb{R}} Ho(\mathcal{D})$ .

- 1967 Daniel G. Quillen "Homotopical algebra": form the foundation of homotopy theory.
- 1997 Vladimir Voevodsky "The Milnor Conjecture": apply homotopical thinking to non-homotopical situation.

**Definition:** A model structure on a category  $\mathcal{C}$

Three distinguished classes of morphisms:

- Weak Equivalence
- Fibration
- Cofibration

satisfying the axioms: • Retracts • 2 of 3 • Lifting • Factorization

## Morphisms


Quillen adjunction:  $(L \dashv R) : \mathcal{C} \xrightleftharpoons[L]{R} \mathcal{D}$ .

Quillen equivalence:  $Ho(\mathcal{C}) \xrightleftharpoons[\mathbb{L}]{\mathbb{R}} Ho(\mathcal{D})$ .



# Why do we care equivariant orthogonal spectra?

## Right Philosophy for the Stable Homotopy Theory (1990s)




**The best model:** orthogonal spectra.

## Models for Equivariant Stable Homotopy Theory

$$Ho(\text{Orthogonal } G\text{-spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(G\text{-Spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(S_G\text{-Modules})$$

# Why do we care equivariant orthogonal spectra?

## Right Philosophy for the Stable Homotopy Theory (1990s)



S-modules  $\xrightarrow[\text{MM}]{\text{Q.E.}}$  Orthogonal Spectra  $\xrightarrow[\text{Schwede}]{\text{Q.E.}}$  Symmetric Spectra


**The best model:** orthogonal spectra.

## Models for Equivariant Stable Homotopy Theory

$Ho(\text{Orthogonal } G\text{-spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(G\text{-Spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(S_G\text{-Modules})$

# Why do we care equivariant orthogonal spectra?

## Right Philosophy for the Stable Homotopy Theory (1990s)




**The best model:** orthogonal spectra.

## Models for Equivariant Stable Homotopy Theory

$$Ho(\text{Orthogonal } G\text{-spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(G\text{-Spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(S_G\text{-Modules})$$

# Why do we care equivariant orthogonal spectra?

## Right Philosophy for the Stable Homotopy Theory (1990s)




**The best model:** orthogonal spectra.

## Models for Equivariant Stable Homotopy Theory

$$Ho(\text{Orthogonal } G\text{-spectra}) \xrightarrow[MM]{Q.E.} Ho(G\text{-Spectra}) \xrightarrow[MM]{Q.E.} Ho(S_G\text{-Modules})$$

# Why do we care equivariant orthogonal spectra?

## Right Philosophy for the Stable Homotopy Theory (1990s)



**The best model:** orthogonal spectra.

## Models for Equivariant Stable Homotopy Theory

$$Ho(\text{Orthogonal } G\text{-spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(G\text{-Spectra}) \xrightarrow[\text{MM}]{\text{Q.E.}} Ho(S_G\text{-Modules})$$

$\mathcal{I}_G$ : the category of orthogonal representations of  $G$ .

$Top_G$ : the category of based  $G$ -spaces and continuous based maps.

$\mathcal{I}_G$ -space

A  $G$ -continuous functor  $X : \mathcal{I}_G \longrightarrow Top_G$ .

Orthogonal  $G$ -spectrum

An  $\mathcal{I}_G$ -space  $X$  with a natural transformation  $X(-) \wedge S^- \longrightarrow X(- \oplus -)$  such that the associativity and unitality diagrams commute.

Equivariant notion of a functor with smash product

An  $\mathcal{I}_G$ -FSP is an  $\mathcal{I}_G$ -space  $X$  with

- a unit  $\eta : S \longrightarrow X$ ;
- a natural product  $\mu : X \overline{\wedge} X \longrightarrow X \circ \oplus$

such that the evident unit, associativity and centrality of unit diagram commute.

$\mathcal{I}_G$ : the category of orthogonal representations of  $G$ .

$Top_G$ : the category of based  $G$ -spaces and continuous based maps.

$\mathcal{I}_G$ -space

A  $G$ -continuous functor  $X : \mathcal{I}_G \longrightarrow Top_G$ .

Orthogonal  $G$ -spectrum

An  $\mathcal{I}_G$ -space  $X$  with a natural transformation  $X(-) \wedge S^- \longrightarrow X(- \oplus -)$  such that the associativity and unitality diagrams commute.

Equivariant notion of a functor with smash product

An  $\mathcal{I}_G$ -FSP is an  $\mathcal{I}_G$ -space  $X$  with

- a unit  $\eta : S \longrightarrow X$ ;
- a natural product  $\mu : X \overline{\wedge} X \longrightarrow X \circ \oplus$

such that the evident unit, associativity and centrality of unit diagram commute.

$\mathcal{I}_G$ : the category of orthogonal representations of  $G$ .

$Top_G$ : the category of based  $G$ -spaces and continuous based maps.

$\mathcal{I}_G$ -space

A  $G$ -continuous functor  $X : \mathcal{I}_G \longrightarrow Top_G$ .

Orthogonal  $G$ -spectrum

An  $\mathcal{I}_G$ -space  $X$  with a natural transformation  $X(-) \wedge S^- \longrightarrow X(- \oplus -)$  such that the associativity and unitality diagrams commute.

Equivariant notion of a functor with smash product

An  $\mathcal{I}_G$ -FSP is an  $\mathcal{I}_G$ -space  $X$  with

- a unit  $\eta : S \longrightarrow X$ ;
- a natural product  $\mu : X \overline{\wedge} X \longrightarrow X \circ \oplus$

such that the evident unit, associativity and centrality of unit diagram commute.



$\mathcal{I}_G$ : the category of orthogonal representations of  $G$ .

$Top_G$ : the category of based  $G$ -spaces and continuous based maps.

$\mathcal{I}_G$ -space

A  $G$ -continuous functor  $X : \mathcal{I}_G \longrightarrow Top_G$ .

Orthogonal  $G$ -spectrum

An  $\mathcal{I}_G$ -space  $X$  with a natural transformation  $X(-) \wedge S^- \longrightarrow X(- \oplus -)$  such that the associativity and unitality diagrams commute.

Equivariant notion of a functor with smash product

An  $\mathcal{I}_G$ -FSP is an  $\mathcal{I}_G$ -space  $X$  with

- a unit  $\eta : S \longrightarrow X$ ;
- a natural product  $\mu : X \overline{\wedge} X \longrightarrow X \circ \oplus$

such that the evident unit, associativity and centrality of unit diagram commute.

**Observation:** It has been noticed since the beginning of equivariant homotopy theory that certain theories naturally exist not just for a particular group, but in a uniform way for all groups in a specific class.

⇒ **global homotopy theory**

**Prominent examples:** equivariant stable homotopy, equivariant K-theory, equivariant bordism.

- objects: inner product real spaces;
- morphism set  $L(V, W)$ : the linear isometric embeddings.

An orthogonal space is a continuous functor  $Y : \mathbb{L} \longrightarrow \mathcal{T}$  to the category of topological spaces.

**Observation:** It has been noticed since the beginning of equivariant homotopy theory that certain theories naturally exist not just for a particular group, but in a uniform way for all groups in a specific class.

⇒ **global homotopy theory**

**Prominent examples:** equivariant stable homotopy, equivariant K-theory, equivariant bordism.

- objects: inner product real spaces;
- morphism set  $L(V, W)$ : the linear isometric embeddings.

An orthogonal space is a continuous functor  $Y : \mathbb{L} \longrightarrow \mathcal{T}$  to the category of topological spaces.

**Observation:** It has been noticed since the beginning of equivariant homotopy theory that certain theories naturally exist not just for a particular group, but in a uniform way for all groups in a specific class.

⇒ **global homotopy theory**

**Prominent examples:** equivariant stable homotopy, equivariant K-theory, equivariant bordism.

- objects: inner product real spaces;
- morphism set  $L(V, W)$ : the linear isometric embeddings.

An orthogonal space is a continuous functor  $Y : \mathbb{L} \longrightarrow \mathcal{T}$  to the category of topological spaces.

## Global Homotopy Theory

[Schwede][May]

**Observation:** It has been noticed since the beginning of equivariant homotopy theory that certain theories naturally exist not just for a particular group, but in a uniform way for all groups in a specific class.

⇒ **global homotopy theory**

**Prominent examples:** equivariant stable homotopy, equivariant K-theory, equivariant bordism.

## Orthogonal Space

[Schwede]

### The category $\mathbb{L}$

- objects: inner product real spaces;
- morphism set  $L(V, W)$ : the linear isometric embeddings.

An orthogonal space is a continuous functor  $Y : \mathbb{L} \rightarrow \mathcal{T}$  to the category of topological spaces.

**Observation:** It has been noticed since the beginning of equivariant homotopy theory that certain theories naturally exist not just for a particular group, but in a uniform way for all groups in a specific class.

⇒ **global homotopy theory**

**Prominent examples:** equivariant stable homotopy, equivariant K-theory, equivariant bordism.

- objects: inner product real spaces;
- morphism set  $L(V, W)$ : the linear isometric embeddings.

An orthogonal space is a continuous functor  $Y : \mathbb{L} \longrightarrow \mathcal{T}$  to the category of topological spaces.

# Orthogonal $G$ -spectrum of QEll

Goerss-Hopkins-Miller theorem constructs many examples of  $E_\infty$ -rings which represent elliptic cohomology theories, including Tate K-theory.

## Question

Can we construct  $E_\infty - G$ -spectrum which represents equivariant elliptic cohomology theory (e.g.  $G$ -equivariant Tate K-theory)?

## Orthogonal $G$ -spectrum of quasi-elliptic cohomology

[Huan]

We construct a commutative  $\mathcal{I}_G$ -FSP  $(E(G, -), \eta, \mu)$ . For each faithful  $G$ -representation  $V$ ,  $E(G, V)$  weakly represents  $QEII_G^V(-)$  in the sense

$$\pi_k(E(G, V)) = QEII_G^V(S^k), \text{ for each } k.$$

Can  $E(G, -)$  arise from an orthogonal spectrum?

No.

For a trivial  $G$ -representation  $V$ , the  $G$ -action on  $E(G, V)$  is not trivial.

# Orthogonal $G$ -spectrum of QEll

Goerss-Hopkins-Miller theorem constructs many examples of  $E_\infty$ -rings which represent elliptic cohomology theories, including Tate K-theory.

## Question

Can we construct  $E_\infty - G$ -spectrum which represents equivariant elliptic cohomology theory (e.g.  $G$ -equivariant Tate K-theory)?

## Orthogonal $G$ -spectrum of quasi-elliptic cohomology

[Huan]

We construct a commutative  $\mathcal{I}_G$ -FSP  $(E(G, -), \eta, \mu)$ . For each faithful  $G$ -representation  $V$ ,  $E(G, V)$  weakly represents  $QEII_G^V(-)$  in the sense

$$\pi_k(E(G, V)) = QEII_G^V(S^k), \text{ for each } k.$$

Can  $E(G, -)$  arise from an orthogonal spectrum?

No.

For a trivial  $G$ -representation  $V$ , the  $G$ -action on  $E(G, V)$  is not trivial.



# Orthogonal $G$ -spectrum of QEll

Goerss-Hopkins-Miller theorem constructs many examples of  $E_\infty$ -rings which represent elliptic cohomology theories, including Tate K-theory.

## Question

Can we construct  $E_\infty - G$ -spectrum which represents equivariant elliptic cohomology theory (e.g.  $G$ -equivariant Tate K-theory)?

## Orthogonal $G$ -spectrum of quasi-elliptic cohomology

[Huan]

We construct a commutative  $\mathcal{I}_G$ -FSP  $(E(G, -), \eta, \mu)$ . For each faithful  $G$ -representation  $V$ ,  $E(G, V)$  weakly represents  $QEII_G^V(-)$  in the sense

$$\pi_k(E(G, V)) = QEII_G^V(S^k), \text{ for each } k.$$

Can  $E(G, -)$  arise from an orthogonal spectrum?

No.

For a trivial  $G$ -representation  $V$ , the  $G$ -action on  $E(G, V)$  is not trivial.

# Orthogonal $G$ -spectrum of QEll

Goerss-Hopkins-Miller theorem constructs many examples of  $E_\infty$ -rings which represent elliptic cohomology theories, including Tate K-theory.

## Question

Can we construct  $E_\infty - G$ -spectrum which represents equivariant elliptic cohomology theory (e.g.  $G$ -equivariant Tate K-theory)?

## Orthogonal $G$ -spectrum of quasi-elliptic cohomology

[Huan]

We construct a commutative  $\mathcal{I}_G$ -FSP  $(E(G, -), \eta, \mu)$ . For each faithful  $G$ -representation  $V$ ,  $E(G, V)$  weakly represents  $QEII_G^V(-)$  in the sense

$$\pi_k(E(G, V)) = QEII_G^V(S^k), \text{ for each } k.$$

Can  $E(G, -)$  arise from an orthogonal spectrum?

No.

For a trivial  $G$ -representation  $V$ , the  $G$ -action on  $E(G, V)$  is not trivial.

# Orthogonal $G$ -spectrum of QEll

Goerss-Hopkins-Miller theorem constructs many examples of  $E_\infty$ -rings which represent elliptic cohomology theories, including Tate K-theory.

## Question

Can we construct  $E_\infty - G$ -spectrum which represents equivariant elliptic cohomology theory (e.g.  $G$ -equivariant Tate K-theory)?

## Orthogonal $G$ -spectrum of quasi-elliptic cohomology

[Huan]

We construct a commutative  $\mathcal{I}_G$ -FSP  $(E(G, -), \eta, \mu)$ . For each faithful  $G$ -representation  $V$ ,  $E(G, V)$  weakly represents  $QEII_G^V(-)$  in the sense

$$\pi_k(E(G, V)) = QEII_G^V(S^k), \text{ for each } k.$$

Can  $E(G, -)$  arise from an orthogonal spectrum?

No.

For a trivial  $G$ -representation  $V$ , the  $G$ -action on  $E(G, V)$  is not trivial.

# Sketch of the Construction

- $V : G$ –representation
- $Sym(V) := \bigoplus_{n \geq 0} Sym^n(V)$ : the total symmetric power.
- $S(G, V)_g := Sym(V) \setminus Sym(V)^g$ .
- $KU$ : the global complex  $K$ –spectrum;
- $(V)_g$ : a specific  $\Lambda_G(g)$ –representation.

$$F_g(G, V) := \mathrm{Map}_{\mathbb{R}}(S^{(V)_g}, KU((V)_g \oplus V^g))$$

$$E_g(G, V) := \{t_1 a + t_2 b \in F_g(G, V) * S(G, V)_g \mid \|b\| \leq t_2\} / \{t_1 c_0 + t_2 b\}.$$

$$E(G, V) := \prod_{g \in G_{conj}^{tors}} \mathrm{Map}_{C_G(g)}(G, E_g(G, V))$$

# Sketch of the Construction

- $V : G$ –representation
- $Sym(V) := \bigoplus_{n \geq 0} Sym^n(V)$ : the total symmetric power.
- $S(G, V)_g := Sym(V) \setminus Sym(V)^g$ .
- $KU$ : the global complex  $K$ –spectrum;
- $(V)_g$ : a specific  $\Lambda_G(g)$ –representation.

$$F_g(G, V) := \operatorname{Map}_{\mathbb{R}}(S^{(V)_g}, KU((V)_g \oplus V^g))$$

$$E_g(G, V) := \{t_1 a + t_2 b \in F_g(G, V) * S(G, V)_g \mid \|b\| \leq t_2\} / \{t_1 c_0 + t_2 b\}.$$

$$E(G, V) := \prod_{g \in G_{conj}^{tors}} \operatorname{Map}_{C_G(g)}(G, E_g(G, V))$$

# Sketch of the Construction

- $V$  :  $G$ –representation
- $Sym(V) := \bigoplus_{n \geq 0} Sym^n(V)$ : the total symmetric power.
- $S(G, V)_g := Sym(V) \setminus Sym(V)^g$ .
- $KU$ : the global complex  $K$ –spectrum;
- $(V)_g$ : a specific  $\Lambda_G(g)$ –representation.

$$F_g(G, V) := \mathrm{Map}_{\mathbb{R}}(S^{(V)_g}, KU((V)_g \oplus V^g))$$

$$E_g(G, V) := \{t_1 a + t_2 b \in F_g(G, V) * S(G, V)_g \mid \|b\| \leq t_2\} / \{t_1 c_0 + t_2 b\}.$$

$$E(G, V) := \prod_{g \in G_{conj}^{tors}} \mathrm{Map}_{C_G(g)}(G, E_g(G, V))$$

# Sketch of the Construction

- $V : G$ –representation
- $Sym(V) := \bigoplus_{n \geq 0} Sym^n(V)$ : the total symmetric power.
- $S(G, V)_g := Sym(V) \setminus Sym(V)^g$ .
- $KU$ : the global complex  $K$ –spectrum;
- $(V)_g$ : a specific  $\Lambda_G(g)$ –representation.

$$F_g(G, V) := \operatorname{Map}_{\mathbb{R}}(S^{(V)_g}, KU((V)_g \oplus V^g))$$

$$E_g(G, V) := \{t_1 a + t_2 b \in F_g(G, V) * S(G, V)_g \mid \|b\| \leq t_2\} / \{t_1 c_0 + t_2 b\}.$$

$$E(G, V) := \prod_{g \in G_{conj}^{tors}} \operatorname{Map}_{C_G(g)}(G, E_g(G, V))$$

# a new Global Homotopy Theory

A category  $D_0$  larger than  $\mathbb{L}$

- objects:  $(G, V, \rho)$  with  $V$  an inner product vector space,  $G$  a compact group and  $\rho$  a faithful group representations  $\rho : G \longrightarrow O(V)$ ,
- morphism:  $\phi = (\phi_1, \phi_2) : (G, V, \rho) \longrightarrow (H, W, \tau)$ 
  - $\phi_2 : V \longrightarrow W$  a linear isometric embedding
  - $\phi_1 : \tau^{-1}(O(\phi_2(V))) \longrightarrow G$  group homomorphism

$$\begin{array}{ccc} G & \xrightarrow{\rho} & O(V) \\ \phi_1 \uparrow & & \downarrow \phi_{2*} \\ \tau^{-1}(O(\phi_2(V))) & \xrightarrow{\tau} & O(W) \end{array}$$

The category of  $D_0$ -spaces

A  $D_0$ -space is a continuous functor  $X : D_0 \longrightarrow \mathcal{T}$ . A morphism of  $D_0$ -spaces is a natural transformation.



# a new Global Homotopy Theory

A category  $D_0$  larger than  $\mathbb{L}$

- objects:  $(G, V, \rho)$  with  $V$  an inner product vector space,  $G$  a compact group and  $\rho$  a faithful group representations  $\rho : G \longrightarrow O(V)$ ,
- morphism:  $\phi = (\phi_1, \phi_2) : (G, V, \rho) \longrightarrow (H, W, \tau)$ 
  - $\phi_2 : V \longrightarrow W$  a linear isometric embedding
  - $\phi_1 : \tau^{-1}(O(\phi_2(V))) \longrightarrow G$  group homomorphism

$$\begin{array}{ccc} G & \xrightarrow{\rho} & O(V) \\ \phi_1 \uparrow & & \downarrow \phi_{2*} \\ \tau^{-1}(O(\phi_2(V))) & \xrightarrow{\tau} & O(W) \end{array}$$

The category of  $D_0$ -spaces

A  $D_0$ -space is a continuous functor  $X : D_0 \longrightarrow \mathcal{T}$ . A morphism of  $D_0$ -spaces is a natural transformation.

## A category $D_0$ larger than $\mathbb{L}$

- objects:  $(G, V, \rho)$  with  $V$  an inner product vector space,  $G$  a compact group and  $\rho$  a faithful group representations  $\rho : G \longrightarrow O(V)$ ,
- morphism:  $\phi = (\phi_1, \phi_2) : (G, V, \rho) \longrightarrow (H, W, \tau)$ 
  - $\phi_2 : V \longrightarrow W$  a linear isometric embedding
  - $\phi_1 : \tau^{-1}(O(\phi_2(V))) \longrightarrow G$  group homomorphism

$$\begin{array}{ccc} G & \xrightarrow{\rho} & O(V) \\ \phi_1 \uparrow & & \downarrow \phi_{2*} \\ \tau^{-1}(O(\phi_2(V))) & \xrightarrow{\tau} & O(W) \end{array}$$

## The category of $D_0$ -spaces

A  $D_0$ -space is a continuous functor  $X : D_0 \longrightarrow \mathcal{T}$ . A morphism of  $D_0$ -spaces is a natural transformation.

## Almost Global Homotopy Theory

[Huan]

- an extension of global homotopy theory;
- classifies those theories that are almost "global";
- the restriction maps are equivariant weak equivalence.

We can define global quasi-elliptic cohomology.

[Huan]

Combining the orthogonal  $G$ -spectra  $\{E(G, -)\}$ , we get an ultra-commutative global ring spectrum in the new theory.

We formulate several model structures and are formulating the one below.

## Conjecture

There is a global model structure on the almost global spaces that is Quillen equivalent to the global model structure on the orthogonal spaces formulated by Schwede in Global Homotopy Theory.

# a new Global Homotopy Theory

## Almost Global Homotopy Theory

[Huan]

- an extension of global homotopy theory;
- classifies those theories that are almost "global";
- the restriction maps are equivariant weak equivalence.

We can define global quasi-elliptic cohomology.

[Huan]

Combining the orthogonal  $G$ -spectra  $\{E(G, -)\}$ , we get an ultra-commutative global ring spectrum in the new theory.

We formulate several model structures and are formulating the one below.

## Conjecture

There is a global model structure on the almost global spaces that is Quillen equivalent to the global model structure on the orthogonal spaces formulated by Schwede in Global Homotopy Theory.

## Almost Global Homotopy Theory

[Huan]

- an extension of global homotopy theory;
- classifies those theories that are almost "global";
- the restriction maps are equivariant weak equivalence.

We can define global quasi-elliptic cohomology.

[Huan]

Combining the orthogonal  $G$ -spectra  $\{E(G, -)\}$ , we get an ultra-commutative global ring spectrum in the new theory.

We formulate several model structures and are formulating the one below.

### Conjecture

There is a global model structure on the almost global spaces that is Quillen equivalent to the global model structure on the orthogonal spaces formulated by Schwede in Global Homotopy Theory.

## Almost Global Homotopy Theory

[Huan]

- an extension of global homotopy theory;
- classifies those theories that are almost "global";
- the restriction maps are equivariant weak equivalence.

We can define global quasi-elliptic cohomology.

[Huan]

Combining the orthogonal  $G$ -spectra  $\{E(G, -)\}$ , we get an ultra-commutative global ring spectrum in the new theory.

We formulate several model structures and are formulating the one below.

## Conjecture

There is a global model structure on the almost global spaces that is Quillen equivalent to the global model structure on the orthogonal spaces formulated by Schwede in Global Homotopy Theory.

*Thank you.*

<http://gagp.sysu.edu.cn/zhenhuan/Zhen-PKU-2017-Slides.pdf>

- Ando, "Isogenies of formal group laws and power operations in the cohomology theories  $E_n$ ", Duke J., 1995
- Ando, Hopkins, Strickland, "The sigma orientation is an  $H_\infty$  map", Amer. J. 2004; arXiv:math/0204053
- Adem, Leida, Ruan, "Orbifolds and stringy topology", Cambridge Tracts in Mathematics, 171. 2007.
- Atiyah, "K-theory", second edition, 1989. Notes by D.W.Anderson.
- Atiyah, "Power operations in K-theory", Quart. J. Math. Oxford Ser. (2) 17 1966.
- Atiyah, "Equivariant K-theory and completion", J. Differential Geometry 3 1969.
- Berger, Moerdijk, "On an extension of the notion of Reedy category", Mathematische Zeitschrift, December 2011.
- Bröcker, Dieck, "Representation of Compact Lie Groups", Springer GTM 98 1985.
- Bruner, May, McClure, Steinberger, " $H_\infty$  ring spectra and their applications", volume 1176 of Lecture Notes in Mathematics, 1986.
- Ganter, "Orbifold genera, product formulas, and power operations", Adv. Math, 2006; arXiv:math/0407021.
- Ganter, "Stringy power operations in Tate  $K$ -theory", Homology, Homotopy, Appl., 2013; arXiv:math/0701565
- Ganter, "Power operations in orbifold Tate  $K$ -theory"; arXiv:1301.2754
- Hirschhorn, "Model categories and their localizations", Mathematical Surveys and Monographs, 99. American Mathematical Society, 2003.
- Hopkins, Kuhn, Ravenel, "Generalized group characters and complex oriented cohomology theories", J. Am. Math. Soc. 13 (2000).
- James, Kerber, "Representation theory of the symmetric group", Encyclopedia of mathematics and its applications v. 16, 1981.
- Katz, Mazur, "Arithmetic moduli of elliptic curves", Annals of Mathematics Studies, vol. 108, 1985.
- Mandell, May, Schwede, Shipley, "Model categories of diagram spectra", Proc. London Math. Soc. 82(2001).
- May, "Equivariant homotopy and cohomology theory", CBMS Regional Conference Series in Mathematics, vol. 91, 1996.
- Rezk, "Quasi-elliptic cohomology", 2011.
- Schommer-Pries, "Central extensions of smooth 2-groups and a finite-dimensional string 2-group", Geom. Topol. 15 (2011).
- Schwede, "Global Homotopy Theory", global.pdf.
- Tomer M. Schlank, Nathaniel Stapleton, "A transchromatic proof of Strickland's theorem", available at arXiv:1404.0717.
- Strickland, "Morava E-theory of symmetric groups", Topology 37 (1998), no. 4.