

# Manual Phase Picking

- Large earthquakes are complete in the public catalogs, while may be missed by matched filter because of poor waveform coherence (e.g. Shelly 2019)
- The S arrivals of a large earthquake can be hard to pick, due to the contamination of P tails
- Thus, it may require manual phase picking to complement the microseismic catalog, and to test the location uncertainty with different combination of phase picks.

# Manual Phase Picking

- Input
  - fctlg: catalog file, just rough location would be enough
  - fsta: station file for predicting P & S arrival times
  - data\_dir: directory of continuous waveform
- Output
  - fpha\_hyp: manually picked and located phase file

Input	Operation	Output	<i>Notes</i>
fctlg & fsta	<i>ctlg2pha.py</i>	fpha_org	predicted phase arrival
fpha_org	<i>cut_events.py</i>	events/[event_name] /[net.sta.chn]	
	SAC <i>ppk</i> P/S/N in t0/1/2/3		only use <i>wh</i>
events & fpha_org	<i>head2pha.py</i>	fpha_man	only phase lines change
fpha_man	<i>cut_events.py</i>	events/[event_name] /[net.sta.chn]	may repeat the manual picking process
fpha_man	event location	fpha_hyp	use Hypo-Interface-Py

# References

- Douglas, A., Bowers, D., & Young, J. B. (1997). On the onset of P seismograms. *Geophysical Journal International*, 129(3), 681-690. <https://doi.org/10.1111/j.1365-246X.1997.tb04503.x>
- Diehl, T., Kissling, E., Bormann, P. (2012): Tutorial for consistent phase picking at local to regional distances. - In: Bormann, P. (Ed.), New Manual of Seismological Observatory Practice 2 (NMSOP-2), Potsdam : Deutsches GeoForschungsZentrum GFZ, 1-21. [https://doi.org/10.2312/GFZ.NMSOP-2\\_IS\\_11.4](https://doi.org/10.2312/GFZ.NMSOP-2_IS_11.4)
- Lomax, A. (2020). Absolute location of 2019 Ridgecrest seismicity reveals a shallow M w 7.1 hypocenter, migrating and pulsing M w 7.1 foreshocks, and duplex M w 6.4 ruptures. *Bulletin of the Seismological Society of America*, 110(4), 1845-1858. <https://doi.org/10.1785/0120200006>