

Canonical Quantization Notes

YILUN LIN

October 2025

Principle

The central idea of canonical quantization is to construct a noncommutative operator algebra that deforms the classical algebra of observables.

1 Procedure of Canonical Quantization

1. Specify the dynamics of the classical system.
 - (a) Construct the Lagrangian (density) $\mathcal{L}[\phi_\nu(\mathbf{x}), \partial_\mu \phi_\nu(\mathbf{x})]$ of the classical field.
 - (b) The Lagrangian must include a kinetic term and a potential term.
 - (c) The kinetic term describes the motion and propagation of the field. It must explicitly depend on the field's derivatives, ensuring that the field has proper propagating degrees of freedom. Crucially, the kinetic term must be a Lorentz scalar, which means that the structure must stay the same under Lorentz transformation. Actually, every term in the Lagrangian must be a Lorentz scalar.
 - (d) The potential term is composed of a mass term and a self-interaction term. The potential term must not include derivatives. The

quadratic term in the fields is the mass term, and the terms that are cubic or higher order in the fields describe self-interaction. Also, each term should be a Lorentz scalar.

- (e) The global symmetry of the theory should be considered while constructing the Lagrangean. For example, because quantum electrodynamics possesses $U(1)$ invariance, the Lagrangean of it must be invariant under $U(1)$ global transformation, or even introduce the corresponding gauge field A_μ to achieve $U(1)$ gauge symmetry if needed. This is because that the gauge symmetry is optional in classical field theory, but it is required in quantum field theory.
- (f) For example, the Lagrangean of the simplest field -- Klein-Gordon Field ($spin = 0$ particle field) -- is $\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}m^2\phi^2$.

Apply the Euler-Lagrange equation:

$$\begin{aligned}\frac{\partial\mathcal{L}}{\partial\phi} - \partial_\mu\left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\right] &= 0 \\ \longrightarrow (\square + m^2)\phi &= 0.\end{aligned}\tag{1.1}$$

2. Determine the conjugate momenta and calculate the Hamiltonian.

- (a) The canonical momentum is defined as

$$\pi_\mu[\phi_\nu(\mathbf{x}), \partial_\alpha\phi_\nu(\mathbf{x})] = \frac{\partial\mathcal{L}}{\partial[\partial_0\phi_\mu(\mathbf{x})]}.\tag{1.2}$$

- (b) Derive the Hamiltonian of the classical field through Legendre transformation, that is,

$$\mathcal{H}[\phi_\nu(\mathbf{x}), \partial_\alpha\phi_\nu(\mathbf{x})] = \pi_\mu[\phi_\nu(\mathbf{x}), \partial_\alpha\phi_\nu(\mathbf{x})]\partial_0\phi^\mu(\mathbf{x}) - \mathcal{L}[\phi_\nu(\mathbf{x}), \partial_\alpha\phi_\nu(\mathbf{x})].\tag{1.3}$$

- (c) But now the explicit parameters of \mathcal{H} are fields and their (both time and spatial) derivatives. In order to undergo canonical quantization, the explicit parameters of \mathcal{H} must be $[\phi_\nu, \pi_\nu, \nabla\phi_\nu]$. That is, time derivatives of the field should be replaced with corresponding canonical momenta. From (1.2), we can obtain the relations $\partial_0\phi_\nu = \dot{\phi}_\nu [\phi_\mu, \pi_\mu, \nabla\phi_\mu]$, with the time derivatives $\partial_0\phi_\mu$ replaced by the same amount of independent canonical momenta π_μ .

3. Analyze constraints in the dynamics of the classical system.

- (a) In order to determine whether there are constraints in the system, we need to observe the Hessian matrix:

$$H_{\mu\nu} := \frac{\partial^2 \mathcal{L}}{\partial (\partial_0\phi^\mu) \partial (\partial_0\phi^\nu)}. \quad (1.4)$$

If the Hessian matrix has full rank, then there is no constraint in the system. Otherwise, some constraints may exist.

- (b) Primary constraints: If the Hessian matrix is not invertible, some time derivatives cannot be solved, using relations (1.2), as a unique result in terms of canonical momenta. Those constraints fall into two types:

$$F(\phi^\mu, \pi_\mu, \mathbf{x}) = 0, \quad (1.5)$$

$$G(\phi^\mu, \partial_0\phi^\mu, \mathbf{x}) = 0. \quad (1.6)$$

The former type (1.5) is constraints on canonical momenta, while the later type (1.6) is constraints on time derivatives. Both two types lead to the fact that some time derivatives cannot be ex-

pressed in terms of canonical momenta. This is a pretty simple and obvious problem about linear equations in linear algebra.

4.