



Maternally expressed gene 3 (MEG3): A tumor suppressor long non coding RNA

Soudeh Ghafouri-Fard^a, Mohammad Taheri^{b,c,*}

^a Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^b Urology and Nephrology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^c Urogenital Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran



ARTICLE INFO

Keywords:

MEG3

Maternally expressed 3

Cancer

Expression

Single nucleotide polymorphism

ABSTRACT

Maternally expressed gene 3 (MEG3) is a long non-coding RNA (lncRNA) located on chromosome 14q32.3. Direct sequencing experiments have shown monoallelic expression of this lncRNA. Several studies have shown down-regulation of this lncRNA in human cancers. In some cases, hypermethylation of the promoter region has been suggested as the underlying mechanism. Functional studies have shown that this lncRNA controls expression of several tumor suppressor genes and oncogenes among them are p53, RB, MYC and TGF- β . Through regulation of Wnt- β -catenin pathway, it also affects epithelial-mesenchymal transition. *In vitro* studies have demonstrated contribution of *MEG3* in defining response to chemotherapeutic agents such as paclitaxel, cisplatin and oxaliplatin. Certain polymorphisms within *MEG3* are implicated in cancer risk (rs7158663, rs4081134 and rs11160608) or therapeutic response of cancer patients (rs10132552). Taken together, this lncRNA is regarded as a putative cancer biomarker and treatment target. In the current review, several aspects of the participation of *MEG3* in carcinogenesis are discussed.

1. Introduction

Human *maternally expressed gene 3 (MEG3)* and its mouse homologue (*Meg3*) have been identified through a subtraction-hybridization strategy using androgenetic and normally fertilized embryos and subsequent EST database searching [1]. This gene has been mapped on distal chromosome 12 and chromosome 14q in mouse and human respectively [1]. Alternative RNA splicing has resulted in production of numerous transcript isoforms from this genetic locus. The principally expressed isoform includes exons 1–4 and 8–10 [2]. Sequence analyses failed to identify significant open reading frame in *MEG3*, so this gene has been further classified as a long non-coding RNA (lncRNA). *MEG3* is a chromatin-interacting lncRNAs that interacts with the polycomb repressive complex 2 (PRC2). This lncRNA has similar targets with EZH2 among them are genes involved in the transforming growth factor- β (TGF- β) pathway. *MEG3* recognizes its targets through RNA–DNA triplex establishment [3]. The first evidences for contribution of *MEG3* in human cancers have been obtained from pituitary non-functioning adenomas where expression of this gene is totally lost [2]. Subsequently, expression of *MEG3* was shown to be decreased in several cancer types. Allelic loss of *MEG3* has been detected in high grade meningiomas but not low grade ones [4]. Promoter hypermethylation is

another underlying cause of *MEG3* down-regulation in non-functioning adenoma [5], meningioma [4] and neuroblastoma [6]. A number of miRNAs have been shown to regulate expression of *MEG3* at post-transcriptional level. In turn, *MEG3* acts as competing endogenous RNA for a number of miRNAs [7].

MEG3 regulates expression p53 tumor suppressor gene and induces p53-dependent transcription through different mechanisms. First, it decreases MDM2 levels and subsequently inhibits ubiquitin-mediated degradation of p53. This speculation has been verified by the observed down-regulation of MDM2 after forced over-expression of *MEG3* [8]. *MEG3* can also directly bind with DNA binding domain of p53 and numerous p53 target genes. Co-transfection of a p53-responsive reporter vector with *MEG3* expression vector has shown that *MEG3* augments the transcriptional activity of p53. Construction of deletion mutants of *MEG3* has confirmed that all three domains of *MEG3* are needed to stimulate the transcriptional activity of p53. As *MEG3* up-regulation has enhanced protein levels of p53 without affecting its transcript levels, it probably increases the stability of the p53 protein. Besides, *MEG3* and p53 direct interaction is mediated by M3 domain of *MEG3* and DNA binding domain of p53 [9]. Notably, *MEG3* can suppress cell proliferation even in the absence of p53 [8]. Expression of this lncRNA has been up-regulated in endothelial cells following p53

* Corresponding author at: Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

E-mail address: mohammad.taheri@sbmu.ac.ir (M. Taheri).




Fig. 1. (A) *MEG3* interacts with PRC2 and its cofactor JARID2 to regulate expression of genes related to TGF- β signaling through trimethylation of histone H3 at lysine 27 (H3K27me3). Expression of CDH1 and MDM2 is inhibited through this mechanism [3,8]. MDM2 has been regarded as one of the downstream targets of TGF- β [13]. CDH1 expression is inhibited by TGF- β -SNAI2 signaling in the process of epithelial-mesenchymal transition [14]. (B) *MEG3* directly binds to p53 and increases its stability. Growth differentiation factor 15 (GDF15) is among p53 targets whose expression is increased by *MEG3* [8,9].

activation [10]. Fig. 1 shows the molecular mechanism of *MEG3* interaction with p53 and its effects on TGF- β signaling.

MEG3 has interactions with the other important tumor suppressor namely retinoblastoma protein (Rb). *MEG3* expression is modulated by this tumor suppressor in association with cell proliferation. Genetic deletion of Rb family members in mice has resulted in remarkable silencing of *MEG3* expression. Conversely, activation of pRb enhanced the expression of *MEG3* [11]. *MEG3* effects on Rb can be exerted directly by modulation of Rb phosphorylation and indirectly by induction of p16^{INK4A} which enhances Rb expression [12].

Moreover, *MEG3* has distinct interactions with the PRC2 complex. Such interaction contributes in regulation of expression of transforming growth factor- β (TGF- β) pathway genes. An altered chromatin oligo affinity precipitation (ChOP) strategy has shown direct interactions between a number of TGF- β pathway genes and *MEG3*. Further experiments have shown the role of GA-rich sequences in facilitation of *MEG3* RNA transport to these target sequences [3].

1.1. Expression of *MEG3* in human cancers

Consistent with the proposed tumor suppressor role for *MEG3*, several studies have reported down-regulation of this lncRNA in tumoral tissues compared with non-tumoral tissues of the same origin (Table 1). *MEG3* over-expression has halted cell proliferation and induced apoptosis in several cancer cell lines including lung cancer [15], squamous cell carcinoma [16], gastrointestinal cancer [17] and so on. Contrary to the bulk of evidence, Wang et al. have shown over-expression of *MEG3* in osteosarcoma cell lines when compared to a human osteoblast cell line. Knock down of this lncRNA inhibited cells growth and metastasis and induced apoptosis in an osteosarcoma cell line through modulating the expression of miR-127 [18]. However, other studies in osteosarcoma cell lines and clinical samples did not confirm the results of the mentioned study [19–21]. A recent pilot study has shown higher levels of *MEG3* in plasma samples obtained from colorectal cancer patients compared with non-cancerous controls [22].

1.2. Diagnostic/ prognostic value of *MEG3*

A pilot study has shown that a three-lncRNA panel including *91H*, *PVT-1* and *MEG3* can differentiate colorectal cancer patients from non-cancerous individuals with area under curve (AUC) value of 0.877,

sensitivity of 82.76% and specificity of 78.57%. Notably, this panel was more sensitive (but not more specific) in the recognition of early-stage patients than the combination of two traditionally used biomarkers namely carcinoembryonic antigen (CEA) and CA19-9 [22]. Several studies have shown association between expression level of this lncRNA and patients' survival. A systematic literature search has shown association between *MEG3* down-regulation and poor patients' outcome in the pooled analysis [106]. Moreover, hypermethylation of *MEG3* in plasma has been linked with poor outcome in cervical cancer patients [107]. Table 2 summarizes studies reporting diagnostic/prognostic value of *MEG3* in cancers.

1.3. *MEG3* polymorphisms and cancer

Cao et al. have genotyped six tag single nucleotide polymorphisms (SNPs) of *MEG3* in a population of Chinese colorectal cancer patients and healthy subjects. They reported that *MEG3* rs7158663 AA genotype, but not the heterozygote genotype, confers risk of colorectal cancer in this population. Although the increased risk was correlated with lower age of disease onset and family history of cancer, it was not associated with tumor site or stage [114]. Zhou et al. have genotyped rs7158663 and rs4081134 in neuroblastoma children and controls. Although they found no association between mentioned SNPs and neuroblastoma risk, they reported associations between rs4081134 AG/AA genotypes and early disease onset and high clinical stage. They suggested *MEG3* as a weak-effect risk locus for this kind of cancer [115]. Yang et al. have genotyped these two SNPs in a population of Chinese patients with lung cancer and healthy subjects. They reported associations between rs4081134 AA genotype and risk of lung cancer in total participants as well as several subgroups analyses. However, they did not find association between rs7158663 SNP and lung cancer [116]. Recently, Hou et al. have assessed association between selected *MEG3* SNPs and risk of oral squamous cell carcinoma. They demonstrated significant associations between rs11160608 CC genotype and cancer risk especially among drinkers [117]. Finally, assessments in a population of nasopharyngeal cancer patients have shown association between *MEG3* rs10132552 CC genotype and higher risk of grade 3–4 anemia following chemoradiotherapy (CRT). Besides, the rs10132552 CT genotype was associated with better response to CRT [118]. Table 3 summarizes the data on the association between *MEG3* polymorphisms and cancer risk or therapeutic response.

Table 1
Summary of studies regarding the role of *MEG3* in human cancers.

Cancer type	Sub-type	Expression pattern	Numbers of clinical samples (tissues, serum, etc.)	Assessed cell line	Targets/Regulators	Signaling Pathways	Function	Ref.
Lung cancer	NSCLC	-	-	A549, A549/DDP	p53, β -catenin, survivin	WNT/ β -catenin signaling	\downarrow cisplatin resistance, \uparrow apoptosis	[23]
				A549, SK-MES-1	Rb, p107, DNMT1,	RB pathway	\downarrow Proliferation, \uparrow apoptosis	[15]
				A549, LC-2/ad	TGF- β , CDH1, EZH2, miR-200a, miR-200c, E-cadherin, ZEB1, ZEB2, JARID2	EMT	\downarrow Proliferation, \uparrow apoptosis	[24]
				A549	p53		\uparrow Response to paclitaxel, \downarrow proliferation, \uparrow apoptosis	[25]
Nasopharyngeal cancer		Down	46 resected tumor tissues (23 cisplatin-sensitive and 23 cisplatin-resistant)	A549, H1299,	miR-21-5p, SOX7		\downarrow Proliferation, \uparrow apoptosis, \uparrow response to cisplatin	[26]
				A549, A549/DDP	p53, Bcl-xl		\downarrow cisplatin resistance, \uparrow apoptosis, \downarrow proliferation	[27]
				A549, SPC-A1, NCI-H1650, NCI-H358, NCI-H1299, NCI-H1975, SK-MES-1, 16HBE	p53		\downarrow Proliferation, \uparrow apoptosis, \downarrow invasion	[28]
				A549	Skp2, p27, miR-3163		\downarrow cell growth	[15]
				H1299,	MYC		\downarrow Proliferation, \uparrow apoptosis, \downarrow migration	[29]
				BEAS-2B, A549, HCC823	MicroRNA-7-5p, Bcl-2, Bax, BRCA1		\uparrow Apoptosis	[30]
Breast cancer		-	39 primary cancer samples and 22 noncancerous nasopharyngeal epithelia	G666-1, HK-1, NP69, NP361, NP460, xeno-666, xeno-2117, xeno-1915, xeno-99186, C15, C17	p53		\downarrow Proliferation, \downarrow colony formation	[31]
				BT-549, MDA-MB-231, HF,	PRC2, EZH2	TGF- β signaling pathway		[32]
				MDA-MB-231, MCF7,	GRP78, IRE1, PERK, ATF6, CHOP, caspase-3, NF- κ B, p53	NF- κ B and p53 signaling	\downarrow tumor growth, \uparrow apoptosis	[33]
				MDA-MB-231, MCF-7, MCF-10A, HMEC-1,	SDF-1, VEGFA, VEGFR, bFGF, TGF- β 1, Angiogenin, MMP-9	AKT pathways	\downarrow cell growth, \downarrow invasion, \downarrow angiogenesis, \downarrow proliferation	[34]
Tongue cancer Oral cancer	SCC	Down	90 paired tissues	MDA-MB-231, MCF-7, SKBR3, MCF-10A,	miR-421, E-cadherin,	EMT	\downarrow Proliferation, \downarrow invasion	[34]
				MCF-7, MDA-MB-231, MDA-MB-453, T47D, MCF-10A	miR-21,	PI3K/Akt pathway	\downarrow tumor growth, \downarrow cell proliferation, \downarrow glycolysis, \uparrow apoptosis	[35]
				MCF-7, MDA-MB-231, SKBR3, HEK 293 T, MCF10A	SP3, DNMT1, miR-506, SP1,		\downarrow Migration, \downarrow invasion	[36]
				HOK, SCC-15, CAL27	miR-26a, DNMT3B			[37]
				SCC15, Cal27,	p53, MDM2	WNT/ β -catenin signaling pathway	\downarrow Proliferation, \uparrow apoptosis, \downarrow migration, \downarrow metastasis	[16]
				KYSE-410, HEEC, ECA-109, TE-1, TE-13	GRP78, IRE1, PERK, ATF6, CHOP, caspase-3		\downarrow Proliferation, \uparrow apoptosis, \downarrow metastasis	[38]
Esophageal cancer	SCC	Down	96 paired tissues	EC109,	miR-9, FOXO1, E-cadherin	ER stress pathway	\downarrow Proliferation, \uparrow apoptosis	[39]
				TE1, TE13, Eca109, YES2, T.Tn, HEEpic			\downarrow Proliferation, \downarrow invasion	[40]

(continued on next page)

Table 1 (continued)

Cancer type	Sub-type	Expression pattern	Numbers of clinical samples (tissues, serum, etc.)	Assessed cell line	Targets/Regulators	Signaling Pathways	Function	Ref.
Gastric cancer		Down	72 paired tissues	SGC7901, AGS, MGC803, MKN45, BGC823, GES-1	p53		↓ Proliferation, ↑ apoptosis	[17]
			30 gastric cancer tissues and 30 normal tissues	AGS, NCI-N87, SGC-7901, MKN-45, TMK-1, GES-1	miR-21	EMT	↓ Invasion, ↓ migration, ↓ cell mobility, ↓ tumor growth, ↓ metastasis	[42]
			82 paired tissues	MKN74, MKN45, SGC7901, AGS, GES-1	MIR-21	MEG3/miR-21 axis	↓ lymph node metastasis	[42]
			30 paired tissues				↓ Proliferation, ↓ metastasis	[43]
			31 paired tissues	SGC7901, BGC823, MKN45, HGC27, GES-1		p53 signaling pathway	↓ Proliferation, ↓ metastasis	[44]
			50 paired tissues	MGC-803, HGC-27, MKN-45, SGC-7901, BGC-823, AGS	miR-181 family, Bcl-2		↓ Proliferation, ↓ migration, ↓ invasion, ↑ apoptosis	[45]
Colon cancer		Down	75 tumor samples and 75 normal tissues	SGC-7901, MKN45, HEK-293 T, GES-1	miR-141, E2F3		↓ Proliferation, ↓ cell cycle progression, ↑ apoptosis, ↓ cell growth	[46]
			134 paired tissues	MKN45, SGC7901, BGC823, AGS	miR-770		↓ Proliferation, ↓ invasion	[47]
			52 gastric cancer samples	SGC-7901, BGC-823 GES-1	miR-148a, DNMT-1		↓ Proliferation	[48]
			50 paired tissues	HCT-116, DLD-1			↓ Proliferation	[49]
			62 paired tissues	HT29, SW480			↓ Proliferation	[50]
			80 oxaliplatin-responding tissues and 80 non-responding tissues + serum samples from 70 patients showing response to oxaliplatin and 70 showing no response				↓ Chemoresistance to oxaliplatin, ↑ apoptosis	[40]
			61 paired tissues			Vitamin D Signaling	↓ Proliferation, ↓ tumor growth, ↓ migration, ↓ metastasis	[51]
			84 paired tissues	RKO, SW1116, HT29, HCT116, LoVo, SW620, SW480, 293 T, NCM460, SW480, HT29, HCT116, HT29/OXA, HCT116/OXA	Clusterin miR-141, PDCD4		↓ oxaliplatin resistance	[52]
			84 paired tissues + Serum samples 34 healthy controls and CRC patients	Hs 698.T, Hs 255.T, SNU-C1,	SPHK1, TGF-β1		↓ Proliferation, ↑ apoptosis	[53]
			Plasma samples of 58 cancer patients and 56 healthy individuals					[22]
Pancreas cancer	PNET	-	-	MIN6-4 N, Vec-4 N, M27-4 N, HEK293, CCL-153, EC-304, BON1, QGP-1,	c-MET, EZH2, HGF, Cyclin D1, CDK1, Cyclin D1, CDK4, caspase-3, caspase-9, Bcl-2, miR-183, BR13 p53	p38/ERK/AKT and Wnt/β-Catenin	↓ Proliferation	[54]
			-	PANC-1, SW1990			↓ Metastasis, ↓ cell viability, ↑ apoptosis, ↓ migration, ↓ invasion, ↓ cells growth	[55]
			30 paired tissues	PANC-1	PI3K, AKT, Bcl-2, Bax, Cyclin D1, p53, MMP-2, MMP-9	PI3K/AKT/Bcl-2/Bax/Cyclin D1/p53 and MMP-9	↓ Proliferation, ↑ chemo-response to fenofibrate	[56]
			30 paired tissues	PANC-1	PI3K, AKT, Bcl-2, Bax, Cyclin D1, p53, MMP-2, MMP-9	PI3K/AKT/Bcl-2/Bax/Cyclin D1/p53 and MMP-9	↓ Proliferation, ↑ apoptosis, ↓ invasion	[47]
		Down	25 paired tissues	SW 1990, COLO357, MIA PaCa-2, T3M4, AsPC-1, BxPC-3, CAPAN-1 PANC-1, hTERT-HPNE	Snail	EMT	↓ Proliferation, ↓ migration, ↓ invasion, ↓ chemoresistance	[57]

(continued on next page)

Table 1 (continued)

Cancer type	Sub-type	Expression pattern	Numbers of clinical samples (tissues, serum, etc.)	Assessed cell line	Targets/Regulators	Signaling Pathways	Function	Ref.
Liver cancer		-		HepG2, Huh-7	miR-29a, miR-185, DNMT1, 3A, 3B, p53, caspase-3, MDM2, cyclin D1, ILF3		↑Chemo-response to DNMT1, 3A and 3B	[58]
				HepG2, Huh-7, HEK293	miR-29a, miR-185, DNMT1, 3A, 3B, p53, caspase-3, MDM2, cyclin D1, ILF3		↓Cell growth, ↑apoptosis	[59]
				HepG2	miR-29a, miR-185, DNMT1, 3A, 3B, p53, caspase-3, MDM2, cyclin D1, ILF3, PERK, ATF6, CHOP, caspase-3, Bay11-7082	ER stress pathway	↓Proliferation, ↑apoptosis	[59]
				Hep3B	miR-29a, miR-185, DNMT1, 3A, 3B, p53, caspase-3, MDM2, cyclin D1, ILF3, PERK, ATF6, CHOP, caspase-3, Bay11-7082		↓Cell growth	[51]
				SMMC-7721, Huh-7, MHCC97H, HCC LM3, L02	miR-29a, miR-185, DNMT1, 3A, 3B, p53, caspase-3, MDM2, cyclin D1, ILF3, PERK, ATF6, CHOP, caspase-3, Bay11-7082	EMT	Interaction with Arsenic trioxide	[60]
Gallbladder cancer		Down	72 paired tissues	HCC Huh-7, SMMC-7721, HepG2, 7402, L02	miR-664, ADH4, NF-κB		↓Proliferation, ↓colony formation	[61]
				MHCC97H, 97 L, Hep3B, L02	p53, UHRF1, DNMT1		↓Proliferation, ↑apoptosis	[62]
				HepG2, SK-Hep-1, HEK293, HCT116,	p53		↓Growth, ↓proliferation, ↑apoptosis	[63]
				HepG2, Hep3B, MHCC97H, SMMC-7721, L02	miR-26a, DNMT3B		↓Proliferation, ↓ migration, ↓ invasion, ↓metastasis	[64]
				THLE-3, SMMC-7721, Hep3B, BEL-7402, LM3	AP1G1	PI3K/AKT Pathway	↓Proliferation, ↓invasion	[65]
Cervical cancer		Down	84 paired tissues 50 paired tissues	GBC-SD, QBC939	p53, cyclin D1, caspase 3	EMT	↓Proliferation, ↑apoptosis	[66]
				NOZ, GBC-SD, SGC-996, EH-G1, OCUG-1, H69, Hela	EZH2, LATS2		↓Proliferation, ↑apoptosis, ↓ invasion, ↓tumorigenesis	[67]
				Hela, CaSki	PI3K, AKT, MMP-2, MMP-9, Bcl-2, Bax, P21		↓Proliferation, ↑ apoptosis, ↓ invasion	[68]
				Hela, C-33A, HCC94	miR-21-5p, p53, caspase 3		↓Proliferation, ↑apoptosis	[69]
				Hela, CaSki	Rac1	Rac1 pathway, EMT	↓Proliferation, ↑apoptosis, ↓ tumor growth	[70]
Endometrial cancer		Down	72 paired tissues 20 normal cervical epithelium tissue samples, 20 cervical intraepithelial neoplasia tissue samples, 20 cervical squamous cell carcinoma tissue samples	HEC-1A, KLE	Notch1, Hes1	Notch signaling pathway	↓Proliferation, ↓ tumor growth	[71]
				Ishikawa, HEC-1B	PI3K, m-TOR, P70S6K, VEGFA, BCL-XL		↓Cell viability, ↓migration, ↓ invasion, ↓tumor growth	[72]
				SKOV-3, CAOV-3, OVCAR-3, HO-8910, A2780	PTEN		↓Proliferation, ↑apoptosis, ↓ invasion, ↓migration	[73]
				OVCAR-3, SKOV3, A2780, A2780cp	miR-214		↑Chemo-response to curcumin	[74]
				OVCAR3, SKOV3, HP8910, ES-2 PDS	p53		↓Proliferation, apoptosis	[75]
Ovarian cancer		-	20 normal ovaries and 20 EOCs tissues		p53		↓Tumor growth	[76]
				14 clinically nonfunctioning tumors and 3 normal tissues	GADD45γ			[77]
				38 human pituitary tumor tissues and 88 normal tissues	DLK1			[78]
				44 human pituitary adenomas tissues and 10 normal tissues	Bcl-2, procaspase-9			[79]
				29 paired tissues	MIR-7, RASL11B			[80]
Pituitary cancer		Down	72 paired tissues	786-O, SN12, 293 T				[81]
				HK-2, A-498, 786-O				[82]
								[83]
								[84]
								[85]
Kidney cancer		Down	72 paired tissues					[86]
								[87]
								[88]
								[89]
								[90]

(continued on next page)

Table 1 (continued)

Cancer type	Sub-type	Expression pattern	Numbers of clinical samples (tissues, serum, etc.)	Assessed cell line	Targets/Regulators	Signaling Pathways	Function	Ref.
Bladder cancer		Down	31 paired tissues	T24			↓Autophagy, ↑apoptosis, ↓proliferation	[84]
			21 paired tissues	T24, 5637	p53, MMP2, MMP9, caspase-3, bax		↓Migration, ↓invasion, ↑chemo-response to cisplatin	[85]
			45 paired tissues	5637, RT4, RT-112, T-24, SV-HUC-1	Mir-96, TPM1, Bax, Bcl-2, Cyclin D1		↓Proliferation, ↑apoptosis, ↓tumor growth	[75]
			27 paired tissues	UMUC3, T24T	c-Myc, c-Jun, PHLPP2, MAPK, miR-27a		↓Invasion	[86]
Prostate cancer		Down	21 paired tissues	PC3, DU145, HepG2	Bcl-2, Bax, caspase 3, Cyclin D1		↓Proliferation, ↑apoptosis, ↓cell viability, ↓tumorigenesis	[87]
			85 paired tissues	PC-3, DU145, VCaP, 22RV1, WPMY-1,	miR-9-5p, QKI-5		↓Proliferation, ↓migration, ↓invasion, ↑apoptosis, ↓ tumor growth	[88]
Osteosarcoma		–	–	MG63, HOS	EWSAT1		↓Proliferation, ↓metastasis, ↓ migration, ↓invasion, ↓cell growth	[89]
		Up		MG63, 143B, G292, hFOB1.19, OS-732, SaOS,	miR-127, ZEB1	JNK and Wnt signaling pathways	↑cell viability, ↑ migration, ↑ invasion, ↓apoptosis, ↑metastasis, ↑cell growth	[18]
		Down	–	MG63, U2OS, hFOB 1.19	Notch1, Hes1, TGF-β, N-cadherin, E-cadherin		↓Proliferation, ↓migration, ↓cell growth, ↓metastasis	[19]
Retinoblastoma Melanoma		Down	64 paired tissues	hFOB 1.19, U-2 OS,	miR-664a		↓Migration	[20]
		Down	63 paired tissues	Weri-Rb1, Y79		Wnt/β-catenin	↓Proliferation, ↑ apoptosis	[21]
		Down	42 paired tissues	A375, WM35, SK-MEL-5, SK-MEL-2, HEMA-1P	CYLD, miR-499-5p, N-Cadherin, Cyclin D1, E-cadherin		↓Proliferation, ↑apoptosis, ↓metastasis, ↓invasion, ↓ migration, ↓ tumorigenesis	[90]
Thyroid cancer	PTC	Down	16 metastatic tissues versus matched group of non-metastatic tissues	TPC-1, HTH83, 293 T	Rac1		↓Cell migration, ↓ invasion	[91]
			20 paired tissues	FTC-133, TPC-1, 293 T,	miR-182		↑Apoptosis, ↓proliferation, ↑ radiosensitivity of ¹³¹ I, ↓cell viability	[92]
Glioma		–	–	U87			↑Apoptosis, ↑autophagy, ↓ cisplatin resistance	[83]
				U251	Sirt7, Bax, caspase-3/-9, Beclin-1, p62, LC3-II/LC3-I	PI3K/AKT and mTOR signaling pathways	↓Proliferation, ↓migration, ↑ apoptosis, ↑autophagy, ↓tumor formation	[93]
		Down	17 paired tissues	U251, U87	p53		↓Proliferation, ↑apoptosis	[94]
			71 glioma tumor tissues and 12 normal brain tissues	U251, U87, A172	DNNMT1, p53,		↓Proliferation, ↑apoptosis, ↓ colony formation	[95]
			40 paired tissues	NHAs, U87, U251	miR-19a, PTEN,		↓Migration, ↓invasion, ↓ proliferation, ↑Apoptosis, ↓ tumorigenesis	[96]
			30 paired tissues	U-251, M059J	miR-93, caspase-3, caspase-9	PI3K/AKT pathway	↓Cell growth, ↓proliferation, ↑ apoptosis,	[97]
			54 surgically resected glioma tissues and 10 non-tumor brain samples	U87, U251, NHA		Wnt/β-catenin signaling	↓Proliferation, ↓cell growth	[76]
			79 paired tissues	U251, NHA,			↓Proliferation, ↑apoptosis, autophagy	[98]
								[99]

(continued on next page)

Table 1 (continued)

Cancer type	Sub-type	Expression pattern	Numbers of clinical samples (tissues, serum, etc.)	Assessed cell line	Targets/Regulators	Signaling Pathways	Function	Ref.
Leukemia	CML	Down	68 patients with CML (imatinib-sensitive and imatinib-resistant)	K562,	miR-21, MRP1, MDR1, ABCG2		↓Proliferation, ↑apoptosis, ↑chemo-response to imatinib	[100]
			BM samples of 40 CML patients and 10 healthy donors	KCL22, K562	mi-R21, Bax, bcl-2, MMP-2, MMP-9, LSD1, MDM2, EZH2, PTEN, DNMT1, MYC, AKT, KLF4, SFRP2		↓Proliferation, ↑apoptosis	[101]
AML	AML		BM samples of 60 CML patients and 10 healthy donors	KCL22, K562, DNMT1, DNMT3B, MBD2, MECP2, HDAC1	miR-147, DNMT1, JAK2, TYK2, STAT3, HDAC1, DNMT3B, MBD2, MECP2	JAK/STAT Pathway	↑Chemo-response to Chidamide, ↓Proliferation, ↑apoptosis	[37]
			BM of 42 samples from AML patients and CD34+ cells derived from 15 potential donors for allogeneic bone marrow transplantation		p53, WT1, TET2		↓Tumor growth	[102]
Lymphoma	T-LBL	-	Blood samples from 57 patients with AML and 57 healthy volunteers	THP-1, HL-60, CCL-240, CRL-1582	miR-184		↓Proliferation, ↓invasion, ↓metastasis	[103]
			-	Jurkat, SUP-T1	p-glycoprotein	EMT, PI3K/mTOR signaling pathway	↓Cell growth, ↓invasion, ↓migration, ↑sensitivity to chemotherapeutic agents	[104]
Multiple myeloma		Down	50 human T-LBL tissues and 38 adjacent normal tissues	CCRF-CEM, Jurkat, SUP-T1, H9	miR-214, AIFM2		↓cell growth, ↑apoptosis, ↓proliferation	[105]
			-	ARP-1, LP-1	miR-181, HOXA11		↓tumor growth	[103]

Abbreviations: NSCLC: non-small cell lung cancer, LAD: lung adenocarcinoma, PNET: primitive neuroectodermal tumor, EOC: epithelial ovarian cancer, T-LBL: T-cell acute lymphoblastic lymphoma, AML: acute myeloid leukemia, CML: chronic myeloid leukemia, ccRCC: clear cell renal cell carcinoma, SCC: squamous cell carcinoma, CRC: colorectal cancer, BM: bone marrow, ↓: decrease, ↑: increase, paired samples: tumor and adjacent normal tissues of a same patient.

Table 2
Summary of selected studies reporting diagnostic/prognostic value of *MEG3* expression in cancers.

Sample origin	Sample number	Area under curve	Sensitivity	Specificity	Kaplan-Meier analysis	Univariate cox regression	Multivariate cox regression	Ref.
Cervical cancer	72 paired tissues	For discrimination between tumors < 4 cm and tumors ≥ 4 cm: 0.705 As a biomarker for lymph node metastasis: 0.716	For discrimination between tumors < 4 cm and tumors ≥ 4 cm: 56.1%. As a biomarker for lymph node metastasis: 70.5%	For discrimination between tumors < 4 cm and tumors ≥ 4 cm: 80.6% As a biomarker for lymph node metastasis: 67.9%	<i>MEG3</i> expression level was associated with RFS and OS.	<i>MEG3</i> expression was prognostic factor for RFS.	<i>MEG3</i> expression was independent prognostic marker for RFS.	[71]
Colorectal cancer	Serum samples from 316 patients	0.784	72.86%	61.43%	Low <i>MEG3</i> expression was associated with poor OS and RFS.	<i>MEG3</i> expression level was independent prognostic factor for OS of patients receiving oxaliplatin.	<i>MEG3</i> expression level was independent prognostic factor for OS of patients receiving oxaliplatin.	[108]
Oral squamous cell carcinoma	371 colorectal cancer patient	–	–	–	Patients with higher <i>MEG3</i> expression have better DFS.	<i>MEG3</i> is an independent prognostic factor for the DFS.	<i>MEG3</i> is an independent prognostic factor for the DFS.	[51]
Epithelial ovarian carcinoma	45 paired tissues	0.84	–	–	–	–	–	[109]
	95 ovarian cancer samples, 8 normal samples, 17 benign tissues	benign tumor vs. ovarian cancer: 0.72 normal vs. ovarian cancer: 0.76	–	–	–	–	–	[110]
Breast cancer	40 paired tissues	0.67	–	–	–	–	–	[111]
	207 paired tissues	–	–	–	Patients with low <i>MEG3</i> expression had poor OS rate and PFS.	–	<i>MEG3</i> expression was an independent poor prognostic factor for 5-year OS and 5-year PFS.	[112]
Osteosarcoma	64 paired tissues	–	–	–	Patients with low lncRNA <i>MEG3</i> expression had a shorter OS.	<i>MEG3</i> expression was associated with OS.	<i>MEG3</i> expression was independent prognostic factor.	[21]
Glioma	79 paired tissues	–	–	–	The OS rates in patients with low <i>MEG3</i> expression were lower than patients with high <i>MEG3</i> expression.	Low level of <i>MEG3</i> expression was the leading variable for prognosis.	Low level of <i>MEG3</i> was correlated with poor prognosis.	[113]
Gallbladder cancer	50 paired tissues	–	–	–	Patients with low <i>MEG3</i> levels had a shorter survival than those with high levels.	Low <i>MEG3</i> was prognostic factor.	Low <i>MEG3</i> expression was an independent prognostic factor.	[67]

Abbreviations: OS: overall survival, RFS: recurrence-free survival, PFS: progression-free survival, DFS: disease-free survival.

Table 3
MEG3 polymorphisms and their association with cancer risk or therapeutic response.

Cancer type	Number of cases	MEG3 variant associated with		Ref.
		Cancer risk/phenotype modifier	Therapeutic response	
Colon cancer	518 cases and 527 control	The rs7158663 AA genotype, but not the heterozygote genotype, confers risk of cancer.		[114]
Nasopharyngeal cancer	505 newly diagnosed cases			[118]
Neuroblastoma	392 neuroblastoma children and 783 controls	The rs4081134 AG/AA genotypes are associated with early disease onset and high clinical stage.	The rs10132552 CT genotype was associated with better response to chemoradiotherapy.	[115]
Lung cancer	526 lung cancer patients and 526 healthy controls	The rs4081134 AA genotype was associated with risk of lung cancer.		[116]
Oral squamous cell carcinoma	444 patients and 984 cancer-free controls	The rs11160608 CC genotype was associated with cancer risk especially among drinkers.		[117]

2. Discussion

MEG3 is a tumor suppressor lncRNA which modulates expression of p53 and some of its targets [2]. Although the bulk of evidence indicates remarkable role of *MEG3* in up-regulation of p53 transcription and induction of accumulation of p53 protein [2], a single study in endothelial cells has shown that *MEG3* knock down triggers DNA damage, induces p53 signaling and up-regulates the expression of p53 target genes [10]. However, the same study has indicated that *MEG3* expression is up-regulated in endothelial cells following p53 activation [10]. Further studies are needed to resolve this discrepancy.

MEG3 also contributes in the regulation of several cancer-related pathways such as TGF- β , PI3K/AKT, Notch, mTOR and Wnt/ β -catenin pathways. *MEG3* expression is activated by Wilms' tumor 1 (WT1) whose dysregulation contributes in carcinogenesis process [102]. Preliminary results have indicated the possibility of application of serum levels or methylation status of this lncRNA in cancer diagnosis. Notably, some SNPs within this gene have been associated with risk of diverse cancers. In the prevailing concept, *MEG3* down-regulation is associated with the increased risk of cancer. However, the luciferase assay has shown higher activity of a cancer-associated allele (rs11160608 C allele) in a single study [117]. Consequently, future studies are needed to clarify the mechanism by which this risk allele contributes in cancer risk.

Notably, expression levels of this lncRNA correlates with response of cancer cells to chemotherapeutic agents. In non-small cell lung cancer cells, *MEG3* increases cisplatin sensitivity through modulating miR-21-5p/SOX7 axis [26]. Assessment of clinical samples has shown associations between decreased level of *MEG3* and poor responses to cisplatin-based chemotherapy. Low levels of this lncRNA in clinical samples were accompanied by low p53 protein levels and high Bcl-xl protein levels [27].

Taken together, the results of *in vitro* studies as well as clinical assays indicate that *MEG3* has a pivotal role in the abrogation of carcinogenesis. Consequently, strategies for induction of its expression might be effective in treatment of cancer. As epigenetic mechanisms are implicated in the down-regulation of its expression in tumor tissues, epigenetic-therapies are putative methods for induction of *MEG3* expression and reprogramming of cancer cells.

Acknowledgment

This study was financially supported by Shahid Beheshti University of Medical Sciences.

References

- [1] N. Miyoshi, H. Wagatsuma, S. Wakana, T. Shiroishi, M. Nomura, K. Aisaka, et al., Identification of an imprinted gene, *Meg3/Gtl2* and its human homologue *MEG3*, first mapped on mouse distal chromosome 12 and human chromosome 14q, *Genes Cells* 5 (3) (2000) 211–220.
- [2] Y. Zhou, X. Zhang, A. Klibanski, *MEG3* noncoding RNA: a tumor suppressor, *J. Mol. Endocrinol.* 48 (3) (2012) R45–53.
- [3] T. Mondal, S. Subhash, R. Vaid, S. Enroth, S. Uday, B. Reinius, et al., *MEG3* long noncoding RNA regulates the TGF- β pathway genes through formation of RNA–DNA triplex structures, *Nat. Commun.* 6 (2015) 7743.
- [4] X. Zhang, R. Gejman, A. Mahta, Y. Zhong, K.A. Rice, Y. Zhou, et al., Maternally expressed gene 3, an imprinted noncoding RNA gene, is associated with meningioma pathogenesis and progression, *Cancer Res.* 70 (6) (2010) 2350–2358.
- [5] J. Zhao, D. Dahle, Y. Zhou, X. Zhang, A. Klibanski, Hypermethylation of the promoter region is associated with the loss of *MEG3* gene expression in human pituitary tumors, *J. Clin. Endocrinol. Metab.* 90 (4) (2005) 2179–2186.
- [6] D. Astuti, F. Latif, K. Wagner, D. Gentle, W. Cooper, D. Catchpoole, et al., Epigenetic alteration at the *DLK1-GTL2* imprinted domain in human neoplasia: analysis of neuroblastoma, pheochromocytoma and Wilms' tumour, *Br. J. Cancer* 92 (8) (2005) 1574.
- [7] M.T. Moradi, H. Fallahi, Z. Rahimi, Interaction of long noncoding RNA *MEG3* with miRNAs: a reciprocal regulation, *J. Cell. Biochem.* 120 (3) (2019) 3339–3352.
- [8] Y. Zhou, Y. Zhong, Y. Wang, X. Zhang, D.L. Batista, R. Gejman, et al., Activation of p53 by *MEG3* non-coding RNA, *J. Biol. Chem.* 282 (34) (2007) 24731–24742.
- [9] J.J. Zhu, S.S. Liu, F.Q. Ye, Y. Shen, Y. Tie, J. Zhu, et al., Long noncoding RNA

- MEG3 interacts with p53 protein and regulates partial p53 target genes in hepatoma cells, *PLoS One* 10 (10) (2015).
- [10] M.S. Shihabudeen Haider Ali, X. Cheng, M. Moran, S. Haemmig, M.J. Naldrett, S. Alvarez, et al., lncRNA MEG3 protects endothelial function by regulating the DNA damage response, *Nucleic Acids Res.* 47 (3) (2019) 1505–1522.
- [11] T.L. Kruer, S.M. Dougherty, L. Reynolds, E. Long, T. de Silva, W.W. Lockwood, et al., Expression of the lncRNA maternally expressed gene 3 (MEG3) contributes to the control of lung cancer cell proliferation by the Rb pathway, *PLoS One* 11 (11) (2016) e0166363.
- [12] L. Benetatos, G. Vartholomatos, E. Hatzimichael, MEG3 imprinted gene contribution in tumorigenesis, *Int. J. Cancer* 129 (4) (2011) 773–779.
- [13] S. Araki, J.A. Eitel, C.N. Batuello, K. Bijangi-Vishehsaraei, X.J. Xie, D. Danielpour, et al., TGF-beta1-induced expression of human Mdm2 correlates with late-stage metastatic breast cancer, *J. Clin. Invest.* 120 (1) (2010) 290–302.
- [14] H. Peinado, D. Olmeda, Cano A. Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat. Rev. Cancer* 7 (6) (2007) 415–428.
- [15] T.L. Kruer, S.M. Dougherty, L. Reynolds, E. Long, T. de Silva, W.W. Lockwood, et al., Expression of the lncRNA maternally expressed gene 3 (MEG3) contributes to the control of lung cancer cell proliferation by the Rb pathway, *PLoS One* 11 (11) (2016) e0166363.
- [16] L.F. Jia, S.B. Wei, Y.H. Gan, Y. Guo, K. Gong, K. Mitchelson, et al., Expression, regulation and roles of miR-26a and MEG3 in tongue squamous cell carcinoma, *Int. J. Cancer* 135 (10) (2014) 2282–2293.
- [17] M. Sun, R. Xia, F. Jin, T. Xu, Z. Liu, W. De, et al., Downregulated long noncoding RNA MEG3 is associated with poor prognosis and promotes cell proliferation in gastric cancer, *Tumour Biol.* 35 (2) (2014) 1065–1073.
- [18] Y. Wang, D. Kong, Knockdown of lncRNA MEG3 inhibits viability, migration, and invasion and promotes apoptosis by sponging miR-127 in osteosarcoma cell, *J. Cell. Biochem.* 119 (1) (2018) 669–679.
- [19] S. Zhang, L. Cai, B. Li, MEG3 long non-coding RNA prevents cell growth and metastasis of osteosarcoma, *Bratisl. Lek. Listy* 118 (10) (2017) 632–636.
- [20] Y. Sahin, Z. Altan, K. Arman, E. Bozgeyik, M.K. Ozer, A. Arslan, Inhibition of miR-664a interferes with the migration of osteosarcoma cells via modulation of MEG3, *Biochem. Biophys. Res. Commun.* 490 (3) (2017) 1100–1105.
- [21] Z.-Z. Tian, X.-J. Guo, Y.-M. Zhao, Y. Fang, Decreased expression of long non-coding RNA MEG3 acts as a potential predictor biomarker in progression and poor prognosis of osteosarcoma, *Int. J. Clin. Exp. Pathol.* 8 (11) (2015) 15138.
- [22] H. Liu, D. Ye, A. Chen, D. Tan, W. Zhang, W. Jiang, M. Wang, X. Zhang, A pilot study of new promising non-coding RNA diagnostic biomarkers for early-stage colorectal cancers, *Clin. Chem. Lab. Med. (CCLM)* 57 (7) (2019) 1073–1083.
- [23] Y. Xia, Z. He, B. Liu, P. Wang, Y. Chen, Downregulation of Meg3 enhances cisplatin resistance of lung cancer cells through activation of the WNT/ β -catenin signaling pathway, *Mol. Med. Rep.* 12 (3) (2015) 4530–4537.
- [24] M. Terashima, S. Tange, A. Ishimura, T. Suzuki, MEG3 long noncoding RNA contributes to the epigenetic regulation of epithelial-mesenchymal transition in lung cancer cell lines, *J. Biol. Chem.* 292 (1) (2017) 82–99.
- [25] J. Xu, C. Su, F. Zhao, J. Tao, D. Hu, A. Shi, et al., Paclitaxel promotes lung cancer cell apoptosis via MEG3-P53 pathway activation, *Biochem. Biophys. Res. Commun.* 504 (1) (2018) 123–128.
- [26] P. Wang, D. Chen, H. Ma, Y. Li, lncRNA MEG3 enhances cisplatin sensitivity in non-small cell lung cancer by regulating miR-21-5p/SOX7 axis, *Oncotarget* 10 (2017) 5137.
- [27] J. Liu, L. Wan, K. Lu, M. Sun, X. Pan, P. Zhang, et al., The long noncoding RNA MEG3 contributes to cisplatin resistance of human lung adenocarcinoma, *PLoS One* 10 (5) (2015) e0114586.
- [28] K.H. Lu, W. Li, X.H. Liu, M. Sun, M.L. Zhang, W.Q. Wu, et al., Long non-coding RNA MEG3 inhibits NSCLC cells proliferation and induces apoptosis by affecting p53 expression, *BMC Cancer* 13 (2013) 461.
- [29] L. Yan-Hua, L. Xiang-Lei, L. Hong, W. Jian-Jun, Long noncoding ribonucleic acids maternally expressed gene 3 inhibits lung cancer tumor progression through downregulation of MYC, *Indian J. Cancer* 52 (7) (2015) 190.
- [30] J. Wu, F. Meng, H.J. Li, High expression of lncRNA MEG3 participates in non-small cell lung cancer by regulating microRNA-7-5p, *Eur. Rev. Med. Pharmacol. Sci.* 22 (18) (2018) 5938–5945.
- [31] W.P. Chak, R.W. Lung, J.H. Tong, S.Y. Chan, S.W. Lun, S.W. Tsao, et al., Downregulation of long non-coding RNA MEG3 in nasopharyngeal carcinoma, *Mol. Carcinog.* 56 (3) (2017) 1041–1054.
- [32] T. Mondal, S. Subhash, R. Vaid, S. Enroth, S. Uday, B. Reinius, et al., MEG3 long noncoding RNA regulates the TGF- β pathway genes through formation of RNA-DNA triplex structures, *Nat. Commun.* 6 (2015) 7743.
- [33] Y. Zhang, J. Wu, H. Jing, G. Huang, Z. Sun, S. Xu, Long noncoding RNA MEG3 inhibits breast cancer growth via upregulating endoplasmic reticulum stress and activating NF- κ B and p53, *J. Cell. Biochem.* 120 (4) (2019) 6789–6797.
- [34] C.-y Zhang, M.-s Yu, X. Li, Z. Zhang, C.-r Han, B. Yan, Overexpression of long non-coding RNA MEG3 suppresses breast cancer cell proliferation, invasion, and angiogenesis through AKT pathway, *Tumour Biol.* 39 (6) (2017) 1010428317701311.
- [35] J. Zhang, S. Guo, B. Jia, Down-regulation of long non-coding RNA MEG3 serves as an unfavorable risk factor for survival of patients with breast cancer, *Eur. Rev. Med. Pharmacol. Sci.* 20 (24) (2016) 5143–5147.
- [36] M. Zhu, X. Wang, Y. Gu, F. Wang, L. Li, X. Qiu, MEG3 overexpression inhibits the tumorigenesis of breast cancer by downregulating miR-21 through the PI3K/Akt pathway, *Arch. Biochem. Biophys.* 661 (2019) 22–30.
- [37] Li Z-y, L. Yang, Liu X-j, Wang X-z, Pan Y-x, Luo J-m, The long noncoding RNA MEG3 and its target miR-147 regulate JAK/STAT pathway in advanced chronic myeloid leukemia, *EBioMedicine* 34 (2018) 61–75.
- [38] Z. Liu, C. Wu, N. Xie, P. Wang, Long non-coding RNA MEG3 inhibits the proliferation and metastasis of oral squamous cell carcinoma by regulating the WNT/ β -catenin signaling pathway, *Oncol. Lett.* 14 (4) (2017) 4053–4058.
- [39] D. Lv, R. Sun, Q. Yu, X. Zhang, The long non-coding RNA maternally expressed gene 3 activates p53 and is downregulated in esophageal squamous cell cancer, *Tumour Biol.* 37 (12) (2016) 16259–16267.
- [40] Z.-L. Huang, R.-P. Chen, X.-T. Zhou, H.-L. Zhan, M.-M. Hu, B. Liu, et al., Long non-coding RNA MEG3 induces cell apoptosis in esophageal cancer through endoplasmic reticulum stress, *Oncol. Rep.* 37 (5) (2017) 3093–3099.
- [41] Z. Dong, A. Zhang, S. Liu, F. Lu, Y. Guo, G. Zhang, et al., Aberrant methylation-mediated silencing of lncRNA MEG3 functions as a ceRNA in esophageal cancer, *Mol. Cancer Res.* 15 (7) (2017) 800–810.
- [42] G. Xu, L. Meng, D. Yuan, K. Li, Y. Zhang, C. Dang, et al., MEG3/miR-21 axis affects cell mobility by suppressing epithelial-mesenchymal transition in gastric cancer, *Oncol. Rep.* 40 (1) (2018) 39–48.
- [43] J. Dan, J. Wang, Y. Wang, M. Zhu, X. Yang, Z. Peng, et al., lncRNA-MEG3 inhibits proliferation and metastasis by regulating miRNA-21 in gastric cancer, *Biomed. Pharmacother.* 99 (2018) 931–938.
- [44] G. Wei, X. Wang, lncRNA MEG3 inhibit proliferation and metastasis of gastric cancer via p53 signaling pathway, *Eur. Rev. Med. Pharmacol. Sci.* 21 (17) (2017) 3850–3856.
- [45] W. Peng, S. Si, Q. Zhang, C. Li, F. Zhao, F. Wang, et al., Long non-coding RNA MEG3 functions as a competing endogenous RNA to regulate gastric cancer progression, *J. Exp. Clin. Cancer Res.* 34 (1) (2015) 79.
- [46] X. Zhou, G. Ji, X. Ke, H. Gu, W. Jin, G. Zhang, MiR-141 inhibits gastric cancer proliferation by interacting with long noncoding RNA MEG3 and down-regulating E2F3 expression, *Dig. Dis. Sci.* 60 (11) (2015) 3271–3282.
- [47] W. Guo, Z. Dong, S. Liu, Y. Qiao, G. Kuang, Y. Guo, et al., Promoter hypermethylation-mediated downregulation of miR-770 and its host gene MEG3, a long non-coding RNA, in the development of gastric cardia adenocarcinoma, *Mol. Carcinog.* 56 (8) (2017) 1924–1934.
- [48] J. Yan, X. Guo, J. Xia, T. Shan, C. Gu, Z. Liang, et al., MiR-148a regulates MEG3 in gastric cancer by targeting DNA methyltransferase 1, *Med. Oncol.* 31 (3) (2014) 879.
- [49] C. Li, G. Liang, W. Yao, J. Sui, X. Shen, Y. Zhang, et al., Differential expression profiles of long non-coding RNAs reveal potential biomarkers for identification of human gastric cancer, *Oncol. Rep.* 35 (3) (2016) 1529–1540.
- [50] Yin D-d, Liu Z-j, E. Zhang, R. Kong, Zhang Z-h, Guo R-h, Decreased expression of long noncoding RNA MEG3 affects cell proliferation and predicts a poor prognosis in patients with colorectal cancer, *Tumour Biol.* 36 (6) (2015) 4851–4859.
- [51] Y. Zhu, P. Chen, Y. Gao, N. Ta, Y. Zhang, J. Cai, et al., MEG3 activated by vitamin D inhibits colorectal cancer cells proliferation and migration via regulating clusterin, *EBioMedicine* 30 (2018) 148–157.
- [52] H. Wang, H. Li, L. Zhang, D. Yang, Overexpression of MEG3 sensitizes colorectal cancer cells to oxaliplatin through regulation of miR-141/PDCD4 axis, *Biomed. Pharmacother.* 106 (2018) 1607–1615.
- [53] X. Dong, J. Wang, T. Li, Y. Xu, Li S, Down regulation of lncRNA MEG3 promotes colorectal adenocarcinoma cell proliferation and inhibits the apoptosis by up-regulating TGF- β 1 and its downstream sphingosine kinase 1, *Eur. Rev. Med. Pharmacol. Sci.* 22 (23) (2018) 8265–8272.
- [54] S. Iyer, S.D. Modali, S.K. Agarwal, Long noncoding RNA MEG3 is an epigenetic determinant of oncogenic signaling in functional pancreatic neuroendocrine tumor cells, *Mol. Cell. Biol.* 37 (22) (2017) e00278–17.
- [55] Y.-Y. Zhang, H.-M. Feng, MEG3 suppresses human pancreatic neuroendocrine tumor cells growth and metastasis by down-regulation of MiR-183, *Cell. Physiol. Biochem.* 44 (1) (2017) 345–356.
- [56] D. Hu, C. Su, M. Jiang, Y. Shen, A. Shi, F. Zhao, et al., Fenofibrate inhibited pancreatic cancer cells proliferation via activation of p53 mediated by upregulation of lncRNA MEG3, *Biochem. Biophys. Res. Commun.* 471 (2) (2016) 290–295.
- [57] L. Ma, F. Wang, C. Du, Z. Zhang, H. Guo, X. Xie, et al., Long non-coding RNA MEG3 functions as a tumour suppressor and has prognostic predictive value in human pancreatic cancer, *Oncol. Rep.* 39 (3) (2018) 1132–1140.
- [58] M. Zamani, M. Sadeghizadeh, M. Behmanesh, F. Najafi, Dendrosomal curcumin increases expression of the long non-coding RNA gene MEG3 via up-regulation of epi-miRs in hepatocellular cancer, *Phytomedicine* 22 (10) (2015) 961–967.
- [59] R.P. Chen, Z.L. Huang, L.X. Liu, M.Q. Xiang, G.P. Li, J.L. Feng, et al., Involvement of endoplasmic reticulum stress and p53 in lncRNA MEG3-induced human hepatoma HepG2 cell apoptosis, *Oncol. Rep.* 36 (3) (2016) 1649–1657.
- [60] Z. Fan, J. He, T. Fu, W. Zhang, G. Yang, X. Qu, R. Liu, L. Lv, J. Wang, Arsenic trioxide inhibits EMT in hepatocellular carcinoma by promoting lncRNA MEG3 via PKM2, *Biochem. Biophys. Res. Commun.* 513 (4) (2019) 834–840.
- [61] J.H. He, Z.P. Han, J.M. Liu, J.B. Zhou, M.X. Zou, Y.B. Lv, et al., Overexpression of long non-coding RNA MEG3 inhibits proliferation of hepatocellular carcinoma Huh7 cells via negative modulation of miRNA-664, *J. Cell. Biochem.* 118 (11) (2017) 3713–3721.
- [62] H. Zhuo, J. Tang, Z. Lin, R. Jiang, X. Zhang, J. Ji, et al., The aberrant expression of MEG3 regulated by UHRF1 predicts the prognosis of hepatocellular carcinoma, *Mol. Carcinog.* 55 (2) (2016) 209–219.
- [63] J. Zhu, S. Liu, F. Ye, Y. Shen, Y. Tie, J. Zhu, et al., Long noncoding RNA MEG3 interacts with p53 protein and regulates partial p53 target genes in hepatoma cells, *PLoS One* 10 (10) (2015) e0139790.
- [64] Y. Li, M. Ren, Y. Zhao, X. Lu, M. Wang, J. Hu, et al., MicroRNA-26a inhibits proliferation and metastasis of human hepatocellular carcinoma by regulating DNMT3B-MEG3 axis, *Oncol. Rep.* 37 (6) (2017) 3527–3535.
- [65] Y. Sun, F. Cao, L. Qu, Z. Wang, Liu XJ, MEG3 promotes liver cancer by activating

- PI3K/AKT pathway through regulating AP1G1, *Eur. Rev. Med. Pharmacol. Sci.* 23 (4) (2019) 1459–1467.
- [66] B. Liu, E.-D. Shen, M.-M. Liao, Y.-B. Hu, K. Wu, P. Yang, et al., Expression and mechanisms of long non-coding RNA genes MEG3 and ANRIL in gallbladder cancer, *Tumour Biol.* 37 (7) (2016) 9875–9886.
- [67] L. Jin, Q. Cai, S. Wang, S. Wang, T. Mondal, J. Wang, et al., Long noncoding RNA MEG3 regulates LATS2 by promoting the ubiquitination of EZH2 and inhibits proliferation and invasion in gallbladder cancer, *Cell Death Dis.* 9 (10) (2018) 1017.
- [68] X. Wang, Z. Wang, J. Wang, Y. Wang, L. Liu, X. Xu, LncRNA MEG3 has anti-activity effects of cervical cancer, *Biomed. Pharmacother.* 94 (2017) 636–643.
- [69] R. Qin, Z. Chen, Y. Ding, J. Hao, J. Hu, F. Guo, Long non-coding RNA MEG3 inhibits the proliferation of cervical carcinoma cells through the induction of cell cycle arrest and apoptosis, *Neoplasia* 60 (5) (2013) 486–492.
- [70] J. Zhang, T. Yao, Y. Wang, J. Yu, Y. Liu, Z. Lin, Long noncoding RNA MEG3 is downregulated in cervical cancer and affects cell proliferation and apoptosis by regulating miR-21, *Cancer Biol. Ther.* 17 (1) (2016) 104–113.
- [71] J. Zhang, Z. Lin, Y. Gao, T. Yao, Downregulation of long noncoding RNA MEG3 is associated with poor prognosis and promoter hypermethylation in cervical cancer, *J. Exp. Clin. Cancer Res.* 36 (1) (2017) 5.
- [72] X. Chen, J. Qu, Long non-coding RNA MEG3 suppresses survival, migration, and invasion of cervical cancer, *Oncotarget* 11 (2018) 4999.
- [73] Q. Guo, Z. Qian, D. Yan, L. Li, L. Huang, LncRNA-MEG3 inhibits cell proliferation of endometrial carcinoma by repressing Notch signaling, *Biomed. Pharmacother.* 82 (2016) 589–594.
- [74] K.-X. Sun, D.-D. Wu, S. Chen, Y. Zhao, Z.-H. Zong, LncRNA MEG3 inhibit endometrial carcinoma tumorigenesis and progression through PI3K pathway, *Apoptosis* 22 (12) (2017) 1543–1552.
- [75] J. Wang, W. Xu, Y. He, Q. Xia, S. Liu, LncRNA MEG3 impacts proliferation, invasion, and migration of ovarian cancer cells through regulating PTEN, *Inflamm. Res.* 67 (11–12) (2018) 927–936.
- [76] J. Zhang, J. Liu, X. Xu, L. Li, Curcumin suppresses cisplatin resistance development partly via modulating extracellular vesicle-mediated transfer of MEG3 and miR-214 in ovarian cancer, *Cancer Chemother. Pharmacol.* 79 (3) (2017) 479–487.
- [77] X. Sheng, J. Li, L. Yang, Z. Chen, Q. Zhao, L. Tan, et al., Promoter hypermethylation influences the suppressive role of maternally expressed 3, a long non-coding RNA, in the development of epithelial ovarian cancer, *Cancer Chemother. Pharmacol.* 32 (1) (2014) 277–285.
- [78] P. Chunharojirith, Y. Nakayama, X. Jiang, R.E. Kery, J. Ma, S. Cristine, et al., Tumor suppression by MEG3 lncRNA in a human pituitary tumor derived cell line, *Mol. Cell. Endocrinol.* 416 (2015) 27–35.
- [79] J. Zhao, D. Dahle, Y. Zhou, X. Zhang, A. Klibanski, Hypermethylation of the promoter region is associated with the loss of MEG3 gene expression in human pituitary tumors, *J. Clin. Endocrinol. Metab.* 90 (4) (2005) 2179–2186.
- [80] L.C. Mezzomo, P.H. Gonzales, F.G. Pesce, N. Kretzmair Filho, N.P. Ferreira, M.C. Oliveira, et al., Expression of cell growth negative regulators MEG3 and GADD45 is lost in most sporadic human pituitary adenomas, *Pituitary* 15 (3) (2012) 420–427.
- [81] P. Cheunsuchon, Y. Zhou, X. Zhang, H. Lee, W. Chen, Y. Nakayama, et al., Silencing of the imprinted DLK1-MEG3 locus in human clinically nonfunctioning pituitary adenomas, *Am. J. Pathol.* 179 (4) (2011) 2120–2130.
- [82] M. Wang, T. Huang, G. Luo, C. Huang, Xiao X.-y, L. Wang, et al., Long non-coding RNA MEG3 induces renal cell carcinoma cells apoptosis by activating the mitochondrial pathway, *J. Huazhong Univ. Sci. Technol. Med. Sci.* 35 (4) (2015) 541–545.
- [83] Y. Liu, P. Yue, T. Zhou, F. Zhang, H. Wang, X. Chen, et al., LncRNA MEG3 enhances 131 I sensitivity in thyroid carcinoma via sponging miR-182, *Biomed. Pharmacother.* 105 (2018) 1232–1239.
- [84] L. Ying, Y. Huang, H. Chen, Y. Wang, L. Xia, Y. Chen, et al., Downregulated MEG3 activates autophagy and increases cell proliferation in bladder cancer, *Mol. Biosyst.* 9 (3) (2013) 407–411.
- [85] S. Feng, X. Zhang, H. Fan, Q. Sun, M. Zhang, Up-regulation of LncRNA MEG3 inhibits cell migration and invasion and enhances cisplatin chemosensitivity in bladder cancer cells, *Neoplasia* 65 (6) (2018) 925.
- [86] C. Huang, X. Liao, H. Jin, F. Xie, F. Zheng, J. Li, et al., MEG3, as a competing endogenous RNA, binds with miR-27a to promote PHLPP2 protein translation and impairs bladder cancer invasion, *Mol. Ther. Nucleic Acids* 16 (2019) 51–62.
- [87] G. Luo, M. Wang, X. Wu, D. Tao, X. Xiao, L. Wang, et al., Long non-coding RNA MEG3 inhibits cell proliferation and induces apoptosis in prostate cancer, *Cell. Physiol. Biochem.* 37 (6) (2015) 2209–2220.
- [88] M. Wu, Y. Huang, T. Chen, W. Wang, S. Yang, Z. Ye, et al., LncRNA MEG3 inhibits the progression of prostate cancer by modulating miR-9-5p/QKI-5 axis, *J. Cell. Mol. Med.* 23 (1) (2019) 29–38.
- [89] L. Sun, C. Yang, J. Xu, Y. Feng, L. Wang, T. Cui, Long noncoding RNA EWSAT1 promotes osteosarcoma cell growth and metastasis through suppression of MEG3 expression, *DNA Cell Biol.* 35 (12) (2016) 812–818.
- [90] Y. Gao, L. XJTB, Decreased expression of MEG3 contributes to retinoblastoma progression and affects retinoblastoma cell growth by regulating the activity of Wnt/ β -catenin pathway, *Tumour Biol.* 37 (2) (2016) 1461–1469.
- [91] J. Long, X. Pi, LncRNA-MEG3 suppresses the proliferation and invasion of melanoma by regulating CYLD expression mediated by sponging miR-499-5p, *Biomed Res. Int.* 2018 (2018).
- [92] C. Wang, G. Yan, Y. Zhang, X. Jia, P. Bu, Long non-coding RNA MEG3 suppresses migration and invasion of thyroid carcinoma by targeting of Rac1, *Neoplasia* 62 (4) (2015) 541–549.
- [93] B. Ma, Z. Gao, J. Lou, H. Zhang, Z. Yuan, Q. Wu, et al., Long non-coding RNA MEG3 contributes to cisplatin-induced apoptosis via inhibition of autophagy in human glioma cells, *Mol. Med. Rep.* 16 (3) (2017) 2946–2952.
- [94] D.H. Xu, G.N. Chi, C.H. Zhao, D.Y. Li, Long noncoding RNA MEG3 inhibits proliferation and migration but induces autophagy by regulation of Sirt7 and PI3K/AKT/mTOR pathway in glioma cells, *J. Cell. Biochem.* 120 (5) (2019) 7516–7526.
- [95] P. Wang, Z. Ren, P. Sun, Overexpression of the long non-coding RNA MEG3 impairs in vitro glioma cell proliferation, *J. Cell. Biochem.* 113 (6) (2012) 1868–1874.
- [96] J. Li, E.-B. Bian, X.-J. He, C.-C. Ma, G. Zong, H.-L. Wang, et al., Epigenetic repression of long non-coding RNA MEG3 mediated by DNMT1 represses the p53 pathway in gliomas, *Int. J. Oncol.* 48 (2) (2016) 723–733.
- [97] N. Qin, G.-F. Tong, L.-W. Sun, X.-L. Xu, Long noncoding RNA MEG3 suppresses glioma cell proliferation, migration, and invasion by acting as a competing endogenous RNA of miR-19a, *Oncol. Res.* 25 (9) (2017) 1471–1478.
- [98] X. Gong, M. Huang, Long non-coding RNA MEG3 promotes the proliferation of glioma cells through targeting Wnt/ β -catenin signal pathway, *Cancer Gene Ther.* 24 (9) (2017) 381.
- [99] H. Zhao, X. Wang, X. Feng, X. Li, L. Pan, J. Liu, et al., Long non-coding RNA MEG3 regulates proliferation, apoptosis, and autophagy and is associated with prognosis in glioma, *J. Neurooncol.* 140 (2) (2018) 281–288.
- [100] X. Zhou, P. Yuan, Q. Liu, Z. Liu, LncRNA MEG3 regulates imatinib resistance in chronic myeloid leukemia via suppressing microRNA-21, *Biomol. Ther. (Seoul)* 25 (5) (2017) 490.
- [101] Z. Li, L. Yang, X. Liu, Z. Nie, J. Luo, Long noncoding RNA MEG3 inhibits proliferation of chronic myeloid leukemia cells by sponging microRNA21, *Biomed. Pharmacother.* 104 (2018) 181–192.
- [102] Y. Lyu, J. Lou, Y. Yang, J. Feng, Y. Hao, S. Huang, et al., Dysfunction of the WT1-MEG3 signaling promotes AML leukemogenesis via p53-dependent and-independent pathways, *Leukemia* 31 (12) (2017) 2543.
- [103] X. Shen, H. Bai, H. Zhu, Q. Yan, Y. Yang, W. Yu, et al., Long non-coding RNA MEG3 functions as a competing endogenous RNA to regulate HOXA11 expression by sponging miR-181a in multiple myeloma, *Cell. Physiol. Biochem.* 49 (1) (2018) 87–100.
- [104] R. Deng, F.Y. Fan, H. Yi, F. Liu, G.C. He, H.P. Sun, et al., MEG3 affects the progression and chemoresistance of T-cell lymphoblastic lymphoma by suppressing epithelial-mesenchymal transition via the PI3K/mTOR pathway, *J. Cell. Biochem.* 120 (5) (2019) 8144–8153.
- [105] F.-Y. Fan, R. Deng, H. Yi, H.-P. Sun, Y. Zeng, G.-C. He, et al., The inhibitory effect of MEG3/miR-214/AIFM2 axis on the growth of T-cell lymphoblastic lymphoma, *Int. J. Oncol.* 51 (1) (2017) 316–326.
- [106] M.M. Binabaj, A. Bahrami, A. Bahreyni, M. Shafiee, F. Rahmani, M. Khazaei, et al., The prognostic value of long noncoding RNA MEG3 expression in the survival of patients with cancer: a meta-analysis, *J. Cell. Biochem.* 119 (11) (2018) 9583–9590.
- [107] J. Zhang, T. Yao, Z. Lin, Y. Gao, Aberrant methylation of MEG3 functions as a potential plasma-based biomarker for cervical cancer, *Sci. Rep.* 7 (1) (2017) 6271.
- [108] L.X. Li, J. Shang, Y.P. Zhang, S. Liu, Y.N. Peng, Z. Zhou, et al., MEG3 is a prognostic factor for CRC and promotes chemosensitivity by enhancing oxaliplatin-induced cell apoptosis, *Oncol. Rep.* 38 (3) (2017) 1383–1392.
- [109] L.L. Zhang, D. Hu, L.H. Zou, Low expression of lncRNA MEG3 promotes the progression of oral squamous cell carcinoma by targeting miR-21. European review for medical and pharmacological sciences, *Eur. Rev. Med. Pharmacol. Sci.* 22 (23) (2018) 8315–8323.
- [110] Y.L. Xiu, K.X. Sun, X. Chen, S. Chen, Y. Zhao, Q.G. Guo, et al., Upregulation of the lncRNA Meg3 induces autophagy to inhibit tumorigenesis and progression of epithelial ovarian carcinoma by regulating activity of ATG3, *Oncotarget* 8 (19) (2017) 31714–31725.
- [111] E. Soleimanpour, M. Hosseinpourfeizi, E. Babaei, V. Montazeri, The study of long noncoding RNA, MEG3, expression level and its association with clinicopathologic features in breast cancer, *JBUMS.* 19 (2017) 14–20.
- [112] J. Zhang, S. Guo, B. Jia, Down-regulation of long non-coding RNA MEG3 serves as an unfavorable risk factor for survival of patients with breast cancer, *Eur. Rev. Med. Pharmacol. Sci.* 20 (24) (2016) 5143–5147.
- [113] H. Zhao, X. Wang, X. Feng, X. Li, L. Pan, J. Liu, et al., Long non-coding RNA MEG3 regulates proliferation, apoptosis, and autophagy and is associated with prognosis in glioma, *J. Neurooncol.* 140 (2) (2018) 281–288.
- [114] X. Cao, S. Zhuang, Y. Hu, L. Xi, L. Deng, H. Sheng, et al., Associations between polymorphisms of long non-coding RNA MEG3 and risk of colorectal cancer in Chinese, *Oncotarget* 7 (14) (2016) 19054.
- [115] Z.-J. Zhuo, R. Zhang, J. Zhang, J. Zhu, T. Yang, Y. Zou, et al., Associations between lncRNA MEG3 polymorphisms and neuroblastoma risk in Chinese children, *Aging (Albany NY)* 10 (3) (2018) 481.
- [116] Z. Yang, H. Li, J. Li, X. Lv, M. Gao, Y. Bi, et al., Association between long non-coding RNA MEG3 polymorphisms and lung cancer susceptibility in Chinese northeast population, *DNA Cell Biol.* 37 (10) (2018) 812–820.
- [117] Y. Hou, B. Zhang, L. Miao, Y. Ji, Y. Yu, L. Zhu, et al., Association of long non-coding RNA MEG 3 polymorphisms with oral squamous cell carcinoma risk, *Oral Dis.* 25 (5) (2019) in press paper.
- [118] Y. Wang, Z. Guo, Y. Zhao, Y. Jin, L. An, B. Wu, et al., Genetic polymorphisms of lncRNA-p53 regulatory network genes are associated with concurrent chemoradiotherapy toxicities and efficacy in nasopharyngeal carcinoma patients, *Sci. Rep.* 7 (1) (2017) 8320.