

Medulloblastoma: From Myth to Molecular

Vijay Ramaswamy and Michael D. Taylor

Author affiliations and support information (if applicable) appear at the end of this article.

Published at jco.org on June 22, 2017.

Corresponding author: Michael D. Taylor, MD, PhD, 555 University Ave, Hospital for Sick Children, Toronto, ON M5G1X8; e-mail: mdtaylor@sickkids.ca.

© 2017 by American Society of Clinical Oncology

0732-183X/17/3521w-2355w/\$20.00

ABSTRACT

Current therapies for medulloblastoma were introduced primarily in the 1980s and consist of predominantly cytotoxic, nontargeted approaches. Mortality from medulloblastoma remains significant. In addition, many survivors suffer from severe treatment-related effects of radiation and cytotoxic chemotherapy. Further intensification of nonspecific therapy is unlikely to offer additional benefits, because survival rates have reached a plateau. Recent publications in medulloblastoma have revolved largely around the recognition that medulloblastoma per se does not exist, but rather, that there are a group of histologically similar but clinically and molecularly distinct entities that have been grouped under that rubric. Distinguishing the four molecular subgroups of medulloblastoma—wingless (WNT), sonic hedgehog (SHH), group 3, and group 4—in the daily treatment of patients, as well in the setting of clinical trials, is an important challenge in the near term for the pediatric neuro-oncology community. The preponderance of morbidity in treating patients with medulloblastoma is secondary to the treatment or prophylaxis of leptomeningeal metastases, and the cause of most deaths is leptomeningeal metastases. Recurrence of medulloblastoma is a nearly universally fatal event, with no significant salvage rate. The extent of spatial and temporal intratumoral heterogeneity as medulloblastoma metastasizes to leptomeninges and as it evolves in the face of radiation and cytotoxic chemotherapy is just beginning to be understood as a major barrier to therapeutic success. Pediatric neuro-oncology clinicians and scientists must now determine how best to incorporate rapid changes in our biologic understanding of medulloblastoma into the next generation of upfront clinical trials, with the goal of both improving survival for the highest-risk patients and improving quality of life for survivors.

J Clin Oncol 35:2355-2363. © 2017 by American Society of Clinical Oncology

INTRODUCTION

Medulloblastoma, the most common malignant brain tumor of childhood, was described initially by Cushing and Bailey in the 1920s as a small blue cell tumor of the cerebellum.¹ During the latter years of the 20th century, medulloblastoma and intracranial primitive neuroectodermal tumors were considered similar entities because of their comparable morphology. This controversy in nosology extended to the WHO classification and clinical trial stratification.^{2,3} Moreover, before the clear identification of atypical teratoid/rhabdoid tumors (ATRTs) as a distinct entity, with loss of *SMARCB1*-*INI1*, cerebellar ATRTs were commonly classified as medulloblastoma.^{4,5} Molecular profiling subsequently confirmed the original description that medulloblastoma comprises a distinct entity, separate from other intracranial embryonal tumors.^{6,7}

The revised 2016 WHO classification of CNS tumors defines medulloblastoma both

histologically and genetically.⁸ The histologic classification consists of desmoplastic-nodular, classic, large-cell-anaplastic, and medulloblastoma with extensive nodularity (MBEN). Most medulloblastomas have classic histology. MBENs are restricted to infants, but desmoplastic-nodular-enriched medulloblastoma may occur in infants and adults.⁸⁻¹⁰ The WHO genetic classification divides tumors into wingless (WNT)-activated, sonic hedgehog (SHH)-activated-TP53 wildtype, SHH-activated-TP53 mutant, and non-WNT-SHH, where group 3 and group 4 medulloblastomas are included as provisional entities when a diagnosis is possible. There is overlap between the histologic and genetic classifications; desmoplastic-nodular and MBEN tumors are almost exclusively SHH-activated tumors, and large-cell-anaplastic tumors are enriched for SHH-TP53-mutant, or high-risk group 3 tumors. The usefulness of identifying large-cell-anaplastic morphology in WNT and group 4 is unclear.¹¹ Overall, genetic classification has been

DOI: <https://doi.org/10.1200/JCO.2017.72.7842>

shown to be more reliable and robust, and it forms the basis for the next generation of clinical trials.^{10,11}

Current Risk Stratification and Treatment

Currently, medulloblastomas in children older than 3 to 5 years are stratified into average (or standard-risk in Europe) and high-risk groups. High-risk disease is defined by residual disease > 1.5 cm², metastatic dissemination, and, in some studies, large-cell–anaplastic histology.¹² Infants under the age of 3 to 5 years are treated with radiation-sparing protocols. Adults have been a historically neglected group and are treated mostly with radiation-only approaches.¹³

The origins of current therapy are complex and date back to the 1920s, when Cushing and Bailey observed that in the setting of complete surgical resection, relapse was inevitable.¹⁴ By 1953, 3-year survival had reached 50% when patients were treated with craniospinal irradiation of at least 30 Gy with the posterior fossa boosted to 50 Gy.¹⁵ By 1969, metastatic dissemination as defined by Chang staging (M0 to M4) could significantly predict poor outcome.¹⁶

Medulloblastoma metastasizes largely to the pial surface of the spinal cord and brain in the so-called leptomeninges. The need to provide prophylaxis against future leptomeningeal relapse or the treatment of frank magnetic resonance imaging–visible metastases is the justification for irradiation of the entire nervous system and not just the primary tumor bed. This irradiation of the entire CNS is the major source of severe neurocognitive morbidity in long-term medulloblastoma survivors. This prompted trials evaluating adjuvant chemotherapy, reductions in craniospinal irradiation for average-risk patients, and the elimination of radiation in infants under the age of 3 to 5 years (age cutoffs vary by cooperative group). Although residual disease was never evaluated formally, studies from the 1980s suggested that it was a marker of poorer outcome, when 1.5 cm² was the limit of detection on postoperative computerized tomography.¹²

Currently, high-risk patients older than 3 to 5 years are treated with craniospinal irradiation of 36 to 39 Gy, with a boost to 55 Gy to the posterior fossa–tumor bed, followed by cisplatin–cyclophosphamide–based chemotherapy. This results in 5-year survivals of 60% to 65% across most studies.^{17–22} Average-risk patients receive 23.4 Gy of craniospinal irradiation, with a boost to 55 Gy to the posterior fossa–tumor bed, followed by adjuvant chemotherapy. Average-risk patients achieve 5-year outcomes of approximately 80%, independent of the regimen used.^{23,24} Patients with residual disease have a more favorable outcome compared with those with metastatic disease. It is clear that unselected patients with metastatic disease treated with 23.4 Gy have a dismal outcome.²⁴

Infants are treated with radiation-sparing approaches because of the devastating long-term sequelae of radiation to the entire neuroaxis. Early studies focused on delaying radiation therapy until the age of 3 years, whereas subsequent studies of more intensive chemotherapy revealed that some nonmetastatic infants could be cured without radiation.^{25–30} Importantly, radiation avoidance in infants results in superior neurocognitive outcomes.³¹ Infants with nondesmoplastic metastatic disease (specifically group 3) continue to have dismal outcomes without radiation therapy.

Current risk stratification and therapies applied since the late 1980s pose tremendous challenges. Survival has been stagnant for

almost 30 years despite the intensification of therapy.³² Moreover, incremental advances in anesthesia, radiation, neurosurgery, and diagnostic imaging over the past three to four decades have affected patient outcomes as much as the actual treatments administered. Practices such as increased craniospinal dose in the setting of residual disease and anaplastic morphology are based on the exclusion of presumed high-risk groups from average-risk studies, rather than on an evidence-based rationale that an increased craniospinal dose improves outcome.^{11,33}

Molecular Classification of Medulloblastoma

Initial attempts to molecularly classify medulloblastoma using transcriptomics demonstrate that medulloblastomas cluster apart from ATRTs and primitive neuroectodermal tumors and that combining transcriptomics with clinical parameters stratifies patients more robustly than do clinical criteria alone.^{6,34} A group of favorable-risk patients expressing nuclear β -catenin (WNT subgroup) was identified as having survivals of close to 100%.^{17,35}

Transcriptional profiling of larger cohorts consistently identified four distinct molecular entities termed WNT, SHH, group 3, and group 4.^{36–41} These four subgroups likely arise from distinct cells of origin, and, unlike that of glioblastoma, medulloblastoma subgroup affiliation is stable at recurrence.^{42–44} The four subgroups have distinct clinical and molecular characteristics and form the basis of future preclinical modeling and clinical trial design (Fig 1).^{10,42,43,45–57}

WNT-ACTIVATED MEDULLOBLASTOMA

WNT tumors have activated WNT signaling. They rarely metastasize, have a longer prediagnostic interval, occur in older patients, and have a balanced sex ratio.^{40,41,53,58} WNT tumors frequently invade the lateral recess of the brainstem, probably because their cell of origin resides in the lower rhombic lip.^{42,54}

Highly recurrent, activating somatic mutations in exon 3 of β -catenin (*CTNNB1*) are found in 85% to 90% of WNT tumors, whereas monosomy 6 is seen in 70% to 80%.^{9,41,47} Less frequent events include somatic mutations in *TP53*, *SMARCA4*, and *DDX3X*.^{46–48,52,59} Unlike *TP53* mutation in SHH tumors, *TP53* mutation in WNT tumors is not prognostic.⁵⁹ Rare germline mutations in *APC*, consistent with Turcot syndrome, have been reported in WNT tumors lacking *CTNNB1* mutations, and emerging data suggest referral for genetic counseling in such cases.⁶⁰

Patients younger than 16 years with WNT tumors have a favorable outcome across multiple studies and consistently show 5-year survival rates of > 95%.^{10,11,56,57,61} Patients older than 16 years with WNT tumors have indeterminate outcomes.^{39,61,62} As such, rationale therapies for patients with WNT tumors involve de-escalation of therapy in current clinical trials ([ClinicalTrials.gov](https://clinicaltrials.gov) identifiers: NCT02066220, NCT01878617, and NCT02724579).

SHH-ACTIVATED MEDULLOBLASTOMA

SHH tumors are characterized transcriptionally and genetically by activation of the SHH pathway.^{36,37,40,41,58,63} Infrequently





	WNT	SHH	Group 3	Group 4
Age group				
Metastases at diagnosis	< 5%	20%	40%-50%	35%-40%
Sex ratio	1:1	1:1	2:1	2:1
Somatic nucleotide variants	<i>CTNNB1</i> (90%) <i>SMARCA4</i> (25%) <i>DDX3X</i> (50%) <i>TP53</i> (12%)	<i>PTCH1</i> (25%) <i>SUFU</i> (10%, infants) <i>SMO</i> (15%, adults) <i>TERT</i> (20%, adults) <i>IDH1</i> (< 5%, adults) <i>TP53</i> (15%, children) <i>MLL2</i> (12%)	<i>SMARCA4</i> (10%) <i>CTDNEP1</i> (5%) <i>MLL2</i> (4%)	<i>KDM6A</i> (10%) <i>MLL3</i> (5%)
Germline variants	<i>APC</i> (< 5%)	<i>PTCH1</i> (25% infants, 10% older) <i>SUFU</i> (20% infants) <i>TP53</i> (8% older)		
Focal copy number aberrations		<i>MYCN</i> (8%) <i>GLI2</i> (5%)	<i>MYC</i> (15%) <i>OTX2</i> (7%), <i>GFI1/1b</i> activation (25%)	<i>SNCAIP</i> (10%) <i>MYCN</i> (6%) <i>CDK6</i> (5%) <i>GFI1/1b</i> activation (5%)
Broad copy number aberrations	Monosomy 6 (85%)	9q (35%) 10q (22%) 17p (18%)	i17q (22%) 8 (29%) 10 (45%) 11 (30%) 16 (48%) 1q (23%) 7 (25%) 18 (20%)	i17q (70%) 8p (49%) 11p (28%) X (80% of females) 7q (40%) 18q (20%)
Pattern of relapse	Local and metastatic	Predominantly local	Metastatic	Metastatic
High risk		<i>TP53</i> mutation <i>MYCN</i> amp	Infants Metastases	Metastases
Low risk	Age < 16 years	Infants		Whole chr 11 loss

Fig 1. Molecular subgroups of medulloblastoma. Copy number gains are in red type and losses are in blue type. Amp, amplification; Chr, chromosome; i17q, isochromosome 17q; SHH, sonic hedgehog; WNT, wingless.

metastatic at diagnosis, these tumors have a short presymptomatic interval and occur across all ages. SHH tumors are almost exclusively located within the cerebellum rather than in the fourth ventricle, consistent with their cells of origin (granule cell progenitors).^{42,53-55}

Genetic events in SHH tumors are highly age dependent. Infants frequently harbor germline or somatic mutations in *PTCH1* or *SUFU*. Infants harboring germline *PTCH1* mutations present with Gorlin syndrome (nevroid basal-cell carcinoma syndrome).^{63,64} Children between the ages of 3 and 16 years have mutually exclusive somatic mutations in *PTCH1* or germline and/or-somatic *TP53* mutations, the latter of which frequently co-occur with amplifications of *GLI2* and *MYCN*, most likely secondary to underlying chromosome shattering (chromothripsis).^{63,65} *TP53* mutations are present in approximately 30% of childhood SHH, are frequently germline, and confer a poor prognosis.^{10,59,63} All pediatric SHH tumors warrant genetic counseling because of possible *PTCH1* or *SUFU* germline events in infants, and germline *TP53* (Li-Fraumeni syndrome) in children.¹¹ Adult SHH tumors are characterized by a higher mutational load, with recurrent somatic mutations in *PTCH1*, *SMO*, *TERT* promoter, and rarely *IDH1*.^{63,66,67} The majority of transgenic preclinical mouse models of medulloblastoma are SHH tumors, including a metastatic model.^{68,69}

Outcomes of SHH tumors are age specific. Infants have an excellent outcome, even with chemotherapy-only regimens.^{9,70} In childhood, *TP53*-mutant SHH tumors have a dismal prognosis

compared with that of *TP53*-wildtype tumors.^{10,59} SHH pathway antagonists such as *SMO* inhibitors are currently being evaluated in clinical trials; however, concerns exist regarding premature osseous fusion. Additionally, high-risk SHH patients are unlikely to benefit from their use because of activation of the SHH pathway downstream of *SMO*, limiting widespread clinical adoption.^{63,71} New preclinical therapeutic avenues for the highest-risk group of SHH patients focus on agents acting on downstream targets of the SHH pathway, such as bromodomain inhibitors, PI3K inhibitors, aurora kinase inhibitors, itraconazole, arsenic trioxide, and GLI antagonists.⁷²⁻⁷⁵

GROUP 3 MEDULLOBLASTOMA

Group 3 tumors are characterized transcriptionally by the activation of the GABAergic and photoreceptor pathways.^{37,40,41,58} Group 3 tumors occur in younger children, 40% to 50% are metastatic at diagnosis, have short prediagnostic intervals, and a male preponderance.^{53,58} Anatomically, they are located in the fourth ventricle up against the brainstem. Metastatic group 3 tumors frequently have small primaries.⁵⁵

Somatic nucleotide variants, which have been considered copy number-driven tumors because of the frequency of structural genomic variations, are rare in group 3.^{46-49,52} The most common cytogenetic aberration is amplification of *MYC*, (10% to 20% of tumors), which occurs frequently as a fusion with *PVT1* secondary to complex chromosomal rearrangements at the 8q24

locus.^{40,47,49,56} Focal copy number gains and losses on chromosomes 1 and 9 are also observed in 20% of group 3 tumors (and in 5% of group 4 tumors), which lead to activation of the GFI1 family of oncogenes through a mechanism termed enhancer hijacking.⁴⁵ Arm-level copy number gains and losses are frequent, the most common being isochromosome 17q (i17q) in 40% of group 3 tumors.⁵⁸ Transgenic preclinical models faithfully recapitulating group 3 tumors are lacking, although allograft models of MYC overexpression and p53 loss resemble group 3 tumors.⁷⁶⁻⁷⁸ Most medulloblastoma cell lines and patient-derived xenografts currently in widespread use are also MYC-amplified group 3 tumors.⁷⁹ This unwitting convergence of the research community on MYC-amplified group 3 tumors means that most research is focused on a small percentage of patients, albeit a population with a high mortality rate. This has resulted in a number of promising therapeutic avenues, including bromodomain inhibition, PI3K inhibition including combinations with histone deacetylase (HDAC) and mammalian target of rapamycin (mTOR), aurora kinase inhibitors, gemcitabine, and CDK4/6 inhibitors, being evaluated preclinically for MYC-amplified patients.⁸⁰⁻⁸⁴

Patients with group 3 tumors have a poor outcome overall, particularly nonirradiated infants, but likely benefit from myeloablative treatment regimens.⁷⁰ Although patients with irradiated nonmetastatic group 3 tumors have intermediate outcomes, patients with metastatic group 3 tumors have a poor outcome.^{9,10,21,70} Metastases and/or MYC amplification confer a poor prognosis for patients with group 3 tumors; however, the significance of MYC amplification in nonmetastatic group 3 tumors is indeterminate.^{9,11,61,85} i17q is also a possible marker of poor prognosis in group 3 tumors.⁸⁵

GROUP 4 MEDULLOBLASTOMA

Group 4 tumors are characterized transcriptionally by overrepresentation of neuronal and glutaminergic pathways. Group 4 tumors are the predominant subgroup in children 3 to 16 years of age and have long prediagnostic intervals, and 30% and 40% are metastatic at diagnosis.⁵³ They often present with a nonenhancing primary on initial magnetic resonance imaging.^{54,55}

Somatic nucleotide variants are rare, and group 4 medulloblastomas are also considered copy number-driven tumors. The most common are inactivating mutations of the histone demethylase *KDM6A*, which occur in 10% of cases.^{46-48,52} Tandem duplications of *SNCAIP* are identified in 10% of tumors and are mutually exclusive of amplifications of *CDK6* and *MYCN*, which are observed in 5% to 10% of tumors.^{41,47,49} Unlike SHH tumors, *MYCN* amplifications do not confer a poor prognosis in group 4 medulloblastoma.^{85,86} The most common cytogenetic aberration is i17q, seen in almost 80% of tumors, with less frequent aberrations in 8p, 7q, 11p, and 18q.^{38,40,41,58} No known preclinical models exist that faithfully recapitulate group 4 medulloblastoma other than a limited number of xenografts that slowly grow intracranially, which significantly limits the development of new therapies.

Group 4 tumors have an intermediate prognosis, although in some series, patients with nonmetastatic group 4 tumors can have an excellent prognosis with > 90% survival.¹⁰ Loss of chromosome 11 is a favorable prognostic marker that is currently being

evaluated prospectively.⁸⁵ Patients with group 4 tumors likely benefit from radiation therapy because infants with group 4 tumors have a poor prognosis; furthermore, adults also seem to be at higher risk.⁶² Patients with group 4 tumors have a more frequent need for CSF diversion, possibly relating to both metastatic disease and a longer prediagnostic interval.⁸⁷

Unanswered Controversies

Despite the agreement that medulloblastoma comprises four distinct molecular subgroups, it was acknowledged as early as in the original consensus that additional heterogeneity exists, specifically subtypes of subgroups.⁵⁸ Specifically, there exists age-dependent heterogeneity in SHH, MYC-activated groups in group 3, and copy number-driven group 4 groups; however, the true extent of intrasubgroup heterogeneity is unknown.^{41,88}

A second issue that still requires clarification is the boundary between groups 3 and 4. The WHO classification includes them as provisional entities under the umbrellas of non-WNT and non-SHH.⁸ Groups 3 and 4 share several copy number aberrations such as enrichment of i17q and the fact that they consistently cluster closer to one another than do WNT or SHH.^{10,58} The degree and nature of the overlap between groups 3 and 4 is an active area of exploration and continued investigation that must be resolved to accurately stratify patients in future molecularly informed clinical trials.¹¹

Subgroup Ascertainment

There are several current methods of subgroup determination, including immunohistochemistry, gene expression profiling, and genome-wide methylation profiling.^{11,89} Immunohistochemistry is the least robust of these methods and can identify patients with WNT using a nuclear localization of β -catenin and patients with SHH using GAB1 immunoreactivity.⁹⁰ At least two methods of subgrouping should be used, particularly when changes in clinical management are involved, because some non-WNT tumors have nuclear localization of β -catenin.^{9,11,89} To robustly identify all four subgroups, including groups 3 and 4, either gene expression or DNA methylation profiling is required.^{91,92}

RECURRENT MEDULLOBLASTOMA

Salvage rates for recurrent medulloblastoma are < 10% in previously irradiated children.^{93,94} Unsuccessful approaches tested previously include repeat surgery, re-irradiation, and multiple chemotherapy regimens including myeloablative chemotherapy. In rare instances, infants treated previously with only chemotherapy can be salvaged with 36 Gy of craniospinal irradiation at relapse, albeit with profound neurocognitive sequelae.

The temporal and spatial pattern of relapse is highly subgroup dependent. Group 4 tumors recur later than either SHH or group 3 tumors and survive significantly longer postrecurrence.⁴³ Independent of treatment, the anatomic location of relapse is subgroup dependent; SHH tumors recur predominantly locally; group 3 and 4 tumors recur almost exclusively with metastatic




Fig 2. Resistance to treatment in recurrent group 3 and 4 medulloblastoma. Circles represent individual clones. The clones populating the metastatic compartment at recurrence are highly disparate from the primary site and are more similar to each other. Note the absence of tumor in the primary site after radiation therapy.

dissemination, usually without any evidence of disease in the previously treated surgical bed.⁴³ Although rare, WNT tumors can recur in either the tumor bed or with metastatic dissemination.^{43,94} Isolated metastatic relapses are more common in group 4, whereas diffuse relapses are more common in group 3.

A major barrier to the treatment of recurrent medulloblastoma is temporal heterogeneity. Although subgroup affiliation on the basis of transcription or DNA methylation patterns remains stable, likely related to a conserved cell of origin, most somatic nucleotide variants and copy number events are not conserved between diagnosis and relapse.^{43,81,95} Evaluation of paired samples from diagnosis and relapse revealed drastic genetic divergence after therapy, with only a fraction of events shared between both samples, including the emergence of *TP53* mutations and *MYC*

amplifications at recurrence.^{81,95} This suggests that treatment results in a clonal squeeze caused by expansion of a treatment-resistant clone (Fig 2). Patients with group 3 or 4 medulloblastoma die almost exclusively as a result of metastatic disease, which suggests that a treatment-resistant clone in the metastatic compartment results in relapse.⁴³ These observations are consistent with a cross-species analysis of metastatic medulloblastoma where metastatic clones are an early event and are genetically divergent from the primary tumor (Fig 3).⁶⁹ This represents a tremendous treatment challenge in the context of personalized therapies and suggests that concurrent biopsy is warranted if targeted agents are to be used at recurrence, both to confirm the presence of the target and to exclude secondary radiation-induced glioblastoma, particularly in late relapses.^{11,42}




Fig 3. Clonal selection drives seeding of the metastatic compartment in medulloblastoma. Proposed progression of metastatic dissemination with tumor initiation from the cell of origin, followed by expansion of clones within the tumor. Metastatic clones are shed out of the primary site, and surviving clones then reseed along the leptomeningeal space, followed by growth of highly divergent clones in the leptomeningeal compartment. Metastatic clones are disparate from the primary tumor and are more similar to each other than to the primary.

Major barriers to the treatment of recurrent medulloblastoma include a lack of bona fide model systems and the continued practice in the field of studying the primary tumor, rather than searching for treatments specific to the metastatic compartment and/or targeting clones resistant to conventional treatments. Advances in single-cell genomic techniques and liquid biopsies from plasma and CSF, together with multiregional sampling of both primary and metastatic compartments, have the potential to fully capture the spectrum of intratumoral heterogeneity across the metastatic compartment. Repeat biopsies and sampling at the time of autopsy have the potential to significantly advance our knowledge of the biology of recurrent medulloblastoma. The next generation of preclinical modeling and clinical trials must take into account this clonal divergence. Additional clinical trials targeting genetic events present in the untreated primary tumor, but absent from the post-therapy metastases, are unlikely to confer benefit.

Quality of Life With Medulloblastoma

Neurocognitive adverse effects are common, and many survivors are incapable of living an independent life.⁹⁶ Hearing loss is frequent in patients treated with radiation and/or platinum agents, and growth failure is nearly ubiquitous in prepubertal patients.^{97,98} Secondary malignancies, including secondary glioblastoma and leukemias, approach 3% to 5%.^{99,100} Studies of radiation avoidance in infants have shown that long-term functional outcomes are significantly better than in older irradiated children.³¹ Although advances in proton radiation may improve functional outcomes, there will likely be a limit to this improvement in the youngest children, and the risk of secondary malignancies is unlikely to decrease.⁹⁷

A major challenge in determining long-term functional outcomes has been a lack of long-term quality-of-life parameters in clinical trials. Recent studies have shown that the dose of both craniospinal irradiation and boost volume have a tremendous effect on neurocognitive outcome, which worsens over time.^{101,102} Reducing the boost volume to the tumor bed, as prescribed in the closed SJMB03 and ACNS0331 studies, and introducing proton radiotherapy have seemed to significantly improve neurocognitive outcomes, without compromising survival.^{97,101} Biologic parameters also seem to influence long-term neurocognitive outcome; patients with WNT and group 4 medulloblastoma seem to benefit most from reductions in craniospinal irradiation.³³ Early interventions such as exercise, cognitive training, and possibly neuroprotectants, such as metformin (ClinicalTrials.gov identifier: NCT02040376), have the potential to improve neurocognitive function in survivors.^{103,104}

Future Trial Design

Because the current large cooperative studies are either closed or nearing accrual it is of utmost importance that the next generation of clinical trials incorporate advances in biology. Currently stagnant survival rates will continue to exist if we continue to evaluate nonspecific therapies across biologically discrete groups; smaller studies evaluating more personalized approaches hold more promise for improving survival.




Fig 4. Proposed future molecular risk stratification for both infant and noninfant medulloblastoma. SHH, sonic hedgehog; WNT, wingless.

Although current studies are evaluating therapy de-escalation for patients with WNT tumors, more comprehensive and precise application of biologic risk stratification is needed to improve both survival and functional outcomes. A recent consensus on childhood medulloblastoma, in which patients are risk stratified by subgroup into very low risk, standard risk, high risk, and very high risk, provides a putative framework for the next generation of clinical trials.¹¹ This risk stratification acknowledges *MYC*-amplified and/or metastatic group 3 and *TP53*-mutant SHH as being very high-risk diseases warranting novel upfront approaches. However, unless concepts from biologic studies, such as targeted therapies directed at metastatic disease in groups 3 and 4, are applied rationally, we are unlikely to see any discernable progress (Fig 4). As was the case with de-escalation trials in the 1980s, the adoption of molecular stratification confers an opportunity for bold studies that could potentially improve long-term quality of life and could diminish the risk of secondary malignancies while simultaneously maintaining or improving survival.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at jco.org.

AUTHOR CONTRIBUTIONS

Conception and design: All authors

Collection and assembly of data: All authors

Data analysis and interpretation: All authors

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

REFERENCES

- Ostrom QT, de Blank PM, Kruchko C, et al: Alex's Lemonade Stand Foundation infant and childhood primary brain and central nervous system tumors diagnosed in the United States in 2007-2011. *Neuro-oncol* 16:x1-x36, 2015
- Rorke LB, Gilles FH, Davis RL, et al: Revision of the World Health Organization classification of brain tumors for childhood brain tumors. *Cancer* 56:1869-1886, 1985
- Louis DN, Ohgaki H, Wiestler OD, et al: The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 114:97-109, 2007
- Rorke LB, Packer RJ, Biegel JA: Central nervous system atypical teratoid/rhabdoid tumors of infancy and childhood: Definition of an entity. *J Neurosurg* 85:56-65, 1996
- Biegel JA, Zhou JY, Rorke LB, et al: Germ-line and acquired mutations of INI1 in atypical teratoid and rhabdoid tumors. *Cancer Res* 59:74-79, 1999
- Pomeroy SL, Tamayo P, Gaasenbeek M, et al: Prediction of central nervous system embryonal tumour outcome based on gene expression. *Nature* 415:436-442, 2002
- Sturm D, Orr BA, Toprak UH, et al: New brain tumor entities emerge from molecular classification of CNS-PNETs. *Cell* 164:1060-1072, 2016
- Louis DN, Perry A, Reifenberger G, et al: The 2016 World Health Organization classification of tumors of the central nervous system: A summary. *Acta Neuropathol* 131:803-820, 2016
- Pietsch T, Schmidt R, Remke M, et al: Prognostic significance of clinical, histopathological, and molecular characteristics of medulloblastomas in the prospective HIT2000 multicenter clinical trial cohort. *Acta Neuropathol* 128:137-149, 2014
- Ramaswamy V, Remke M, Adamski J, et al: Medulloblastoma subgroup-specific outcomes in irradiated children: Who are the true high-risk patients? *Neuro-oncol* 18:291-297, 2016
- Ramaswamy V, Remke M, Bouffet E, et al: Risk stratification of childhood medulloblastoma in the molecular era: The current consensus. *Acta Neuropathol* 131:821-831, 2016
- Zeltzer PM, Boyett JM, Finlay JL, et al: Metastasis stage, adjuvant treatment, and residual tumor are prognostic factors for medulloblastoma in children: Conclusions from the Children's Cancer Group 921 randomized phase III study. *J Clin Oncol* 17:832-845, 1999
- Lassaledda A, Ramaswamy V: Medulloblastoma in adults: They're not just big kids. *Neuro-oncol* 18:895-897, 2016
- Bailey P, Cushing H: Medulloblastoma cerebelli: A common type of midcerebellar glioma of childhood. *Arch Neurol Psychiatry* 14:192-224, 1925
- Paterson E, Farr RF: Cerebellar medulloblastoma: Treatment by irradiation of the whole central nervous system. *Acta Radiol* 39:323-336, 1953
- Chang CH, Housepian EM, Herbert C Jr: An operative staging system and a megavoltage radiotherapeutic technique for cerebellar medulloblastomas. *Radiology* 93:1351-1359, 1969
- Gajjar A, Chintagumpala M, Ashley D, et al: Risk-adapted craniospinal radiotherapy followed by high-dose chemotherapy and stem-cell rescue in children with newly diagnosed medulloblastoma (St Jude Medulloblastoma-96): Long-term results from a prospective, multicentre trial. *Lancet Oncol* 7:813-820, 2006
- Tarbell NJ, Friedman H, Polkinghorn WR, et al: High-risk medulloblastoma: A pediatric oncology group randomized trial of chemotherapy before or after radiation therapy (POG 9031). *J Clin Oncol* 31:2936-2941, 2013
- Jakacki RI, Burger PC, Zhou T, et al: Outcome of children with metastatic medulloblastoma treated with carboplatin during craniospinal radiotherapy: A Children's Oncology Group phase I/II study. *J Clin Oncol* 30:2648-2653, 2012
- Gandola L, Massimino M, Cefalo G, et al: Hyperfractionated accelerated radiotherapy in the Milan strategy for metastatic medulloblastoma. *J Clin Oncol* 27:566-571, 2009
- von Bueren AO, Kortmann RD, von Hoff K, et al: Treatment of children and adolescents with metastatic medulloblastoma and prognostic relevance of clinical and biologic parameters. *J Clin Oncol* 34:4151-4160, 2016
- Esbenshade AJ, Kocak M, Hershon L, et al: A phase II feasibility study of oral etoposide given concurrently with radiotherapy followed by dose intensive adjuvant chemotherapy for children with newly diagnosed high-risk medulloblastoma (protocol POG 9631): A report from the Children's Oncology Group. *Pediatr Blood Cancer* 10.1002/pbc.26373 [epub ahead of print on December 21, 2016]
- Taylor RE, Bailey CC, Robinson K, et al: Results of a randomized study of preradiation chemotherapy versus radiotherapy alone for nonmetastatic medulloblastoma: The International Society of Paediatric Oncology/United Kingdom Children's Cancer Study Group PNET-3 Study. *J Clin Oncol* 21:1581-1591, 2003
- Packer RJ, Gajjar A, Vezina G, et al: Phase III study of craniospinal radiation therapy followed by adjuvant chemotherapy for newly diagnosed average-risk medulloblastoma. *J Clin Oncol* 24:4202-4208, 2006
- Rutkowski S, Bode U, Deinlein F, et al: Treatment of early childhood medulloblastoma by postoperative chemotherapy alone. *N Engl J Med* 352:978-986, 2005
- Grill J, Sainte-Rose C, Jouvett A, et al: Treatment of medulloblastoma with postoperative chemotherapy alone: An SFOP prospective trial in young children. *Lancet Oncol* 6:573-580, 2005
- Cohen BH, Geyer JR, Miller DC, et al: Pilot study of intensive chemotherapy with peripheral hematopoietic cell support for children less than 3 years of age with malignant brain tumors, the CCG-99703 phase I/II study. A report from the Children's Oncology Group. *Pediatr Neurol* 53:31-46, 2015
- Geyer JR, Spoto R, Jennings M, et al: Multiagent chemotherapy and deferred radiotherapy in infants with malignant brain tumors: A report from the Children's Cancer Group. *J Clin Oncol* 23:7621-7631, 2005
- Rutkowski S, Cohen B, Finlay J, et al: Medulloblastoma in young children. *Pediatr Blood Cancer* 54:635-637, 2010
- Duffner PK, Horowitz ME, Krischer JP, et al: Postoperative chemotherapy and delayed radiation in children less than three years of age with malignant brain tumors. *N Engl J Med* 328:1725-1731, 1993
- Lafay-Cousin L, Bouffet E, Hawkins C, et al: Impact of radiation avoidance on survival and neurocognitive outcome in infant medulloblastoma. *Curr Oncol* 16:21-28, 2009
- Johnston DL, Keene D, Kostova M, et al: Survival of children with medulloblastoma in Canada diagnosed between 1990 and 2009 inclusive. *J Neurooncol* 124:247-253, 2015
- Thompson EM, Hielscher T, Bouffet E, et al: Prognostic value of medulloblastoma extent of resection after accounting for molecular subgroup: A retrospective integrated clinical and molecular analysis. *Lancet Oncol* 17:484-495, 2016
- Fernandez-Teijeiro A, Betensky RA, Sturla LM, et al: Combining gene expression profiles and clinical parameters for risk stratification in medulloblastomas. *J Clin Oncol* 22:994-998, 2004
- Ellison DW, Onilude OE, Lindsey JC, et al: Beta-catenin status predicts a favorable outcome in childhood medulloblastoma: The United Kingdom Children's Cancer Study Group Brain Tumour Committee. *J Clin Oncol* 23:7951-7957, 2005
- Thompson MC, Fuller C, Hogg TL, et al: Genomics identifies medulloblastoma subgroups that are enriched for specific genetic alterations. *J Clin Oncol* 24:1924-1931, 2006
- Kool M, Koster J, Bunt J, et al: Integrated genomics identifies five medulloblastoma subtypes with distinct genetic profiles, pathway signatures and clinicopathological features. *PLoS One* 3:e3088, 2008
- Remke M, Hielscher T, Korshunov A, et al: FSTL5 is a marker of poor prognosis in non-WNT/non-SHH medulloblastoma. *J Clin Oncol* 29:3852-3861, 2011
- Remke M, Hielscher T, Northcott PA, et al: Adult medulloblastoma comprises three major molecular variants. *J Clin Oncol* 29:2717-2723, 2011
- Northcott PA, Korshunov A, Witt H, et al: Medulloblastoma comprises four distinct molecular variants. *J Clin Oncol* 29:1408-1414, 2011
- Cho Y-J, Tsherniak A, Tamayo P, et al: Integrative genomic analysis of medulloblastoma identifies a molecular subgroup that drives poor clinical outcome. *J Clin Oncol* 29:1424-1430, 2011
- Gibson P, Tong Y, Robinson G, et al: Subtypes of medulloblastoma have distinct developmental origins. *Nature* 468:1095-1099, 2010
- Ramaswamy V, Remke M, Bouffet E, et al: Recurrence patterns across medulloblastoma subgroups: An integrated clinical and molecular analysis. *Lancet Oncol* 14:1200-1207, 2013
- Phillips HS, Kharbanda S, Chen R, et al: Molecular subclasses of high-grade glioma predict prognosis, delineate a pattern of disease progression, and

resemble stages in neurogenesis. *Cancer Cell* 9:157-173, 2006

45. Northcott PA, Lee C, Zichner T, et al: Enhancer hijacking activates GF1 family oncogenes in medulloblastoma. *Nature* 511:428-434, 2014

46. Robinson G, Parker M, Kranenburg TA, et al: Novel mutations target distinct subgroups of medulloblastoma. *Nature* 488:43-48, 2012

47. Northcott PA, Jones DT, Kool M, et al: Medulloblastoma: The end of the beginning. *Nat Rev Cancer* 12:818-834, 2012

48. Pugh TJ, Weeraratne SD, Archer TC, et al: Medulloblastoma exome sequencing uncovers subtype-specific somatic mutations. *Nature* 488:106-110, 2012

49. Northcott PA, Shih DJ, Peacock J, et al: Subgroup-specific structural variation across 1,000 medulloblastoma genomes. *Nature* 488:49-56, 2012

50. Lin CY, Erkek S, Tong Y, et al: Active medulloblastoma enhancers reveal subgroup-specific cellular origins. *Nature* 530:57-62, 2016

51. Hovestadt V, Jones DT, Picelli S, et al: Decoding the regulatory landscape of medulloblastoma using DNA methylation sequencing. *Nature* 510:537-541, 2014

52. Jones DT, Jäger N, Kool M, et al: Dissecting the genomic complexity underlying medulloblastoma. *Nature* 488:100-105, 2012

53. Ramaswamy V, Remke M, Shih D, et al: Duration of the pre-diagnostic interval in medulloblastoma is subgroup dependent. *Pediatr Blood Cancer* 61:1190-1194, 2014

54. Perreault S, Ramaswamy V, Achrol AS, et al: MRI surrogates for molecular subgroups of medulloblastoma. *AJNR Am J Neuroradiol* 35:1263-1269, 2014

55. Raybaud C, Ramaswamy V, Taylor MD, et al: Posterior fossa tumors in children: Developmental anatomy and diagnostic imaging. *Childs Nerv Syst* 31:1661-1676, 2015

56. Kool M, Korshunov A, Remke M, et al: Molecular subgroups of medulloblastoma: An international meta-analysis of transcriptome, genetic aberrations, and clinical data of WNT, SHH, group 3, and group 4 medulloblastomas. *Acta Neuropathol* 123:473-484, 2012

57. Schwalbe EC, Williamson D, Lindsey JC, et al: DNA methylation profiling of medulloblastoma allows robust subclassification and improved outcome prediction using formalin-fixed biopsies. *Acta Neuropathol* 125:359-371, 2013

58. Taylor MD, Northcott PA, Korshunov A, et al: Molecular subgroups of medulloblastoma: The current consensus. *Acta Neuropathol* 123:465-472, 2012

59. Zhukova N, Ramaswamy V, Remke M, et al: Subgroup-specific prognostic implications of TP53 mutation in medulloblastoma. *J Clin Oncol* 31:2927-2935, 2013

60. Hamilton SR, Liu B, Parsons RE, et al: The molecular basis of Turcot's syndrome. *N Engl J Med* 332:839-847, 1995

61. Clifford SC, Lannering B, Schwalbe EC, et al: Biomarker-driven stratification of disease-risk in non-metastatic medulloblastoma: Results from the multicenter HIT-SIOP-PNET4 clinical trial. *Oncotarget* 6:38827-38839, 2015

62. Zhao F, Ohgaki H, Xu L, et al: Molecular subgroups of adult medulloblastoma: A long-term single-institution study. *Neuro-oncol* 18:982-990, 2016

63. Kool M, Jones DT, Jäger N, et al: Genome sequencing of SHH medulloblastoma predicts

genotype-related response to smoothened inhibition. *Cancer Cell* 25:393-405, 2014

64. Smith MJ, Beetz C, Williams SG, et al: Germline mutations in SUFU cause Gorlin syndrome-associated childhood medulloblastoma and redefine the risk associated with PTCH1 mutations. *J Clin Oncol* 32:4155-4161, 2014

65. Rausch T, Jones DT, Zapatka M, et al: Genome sequencing of pediatric medulloblastoma links catastrophic DNA rearrangements with TP53 mutations. *Cell* 148:59-71, 2012

66. Snuderl M, Triscott J, Northcott PA, et al: Deep sequencing identifies IDH1 R132S mutation in adult medulloblastoma. *J Clin Oncol* 33:e27-e31, 2015

67. Remke M, Ramaswamy V, Peacock J, et al: TERT promoter mutations are highly recurrent in SHH subgroup medulloblastoma. *Acta Neuropathol* 126:917-929, 2013

68. Goodrich LV, Milenković L, Higgins KM, et al: Altered neural cell fates and medulloblastoma in mouse patched mutants. *Science* 277:1109-1113, 1997

69. Wu X, Northcott PA, Dubuc A, et al: Clonal selection drives genetic divergence of metastatic medulloblastoma. *Nature* 482:529-533, 2012

70. Lafay-Cousin L, Smith A, Chi SN, et al: Clinical, pathological, and molecular characterization of infant medulloblastomas treated with sequential high-dose chemotherapy. *Pediatr Blood Cancer* 63:1527-1534, 2016

71. Rudin CM, Hann CL, Laterra J, et al: Treatment of medulloblastoma with hedgehog pathway inhibitor GDC-0449. *N Engl J Med* 361:1173-1178, 2009

72. Tang Y, Gholamin S, Schubert S, et al: Epigenetic targeting of Hedgehog pathway transcriptional output through BET bromodomain inhibition. *Nat Med* 20:732-740, 2014

73. Chung FH, Chiang YR, Tseng AL, et al: Functional Module Connectivity Map (FMCMap): A framework for searching repurposed drug compounds for systems treatment of cancer and an application to colorectal adenocarcinoma. *PLoS One* 9:e86299, 2014

74. Kim J, Aftab BT, Tang JY, et al: Itraconazole and arsenic trioxide inhibit Hedgehog pathway activation and tumor growth associated with acquired resistance to smoothened antagonists. *Cancer Cell* 23:23-34, 2013

75. Gustafson WC, Meyerowitz JG, Nekritz EA, et al: Drugging MYCN through an allosteric transition in Aurora kinase A. *Cancer Cell* 26:414-427, 2014

76. Pei Y, Moore CE, Wang J, et al: An animal model of MYC-driven medulloblastoma. *Cancer Cell* 21:155-167, 2012

77. Kawauchi D, Robinson G, Uziel T, et al: A mouse model of the most aggressive subgroup of human medulloblastoma. *Cancer Cell* 21:168-180, 2012

78. Pöschl J, Stark S, Neumann P, et al: Genomic and transcriptomic analyses match medulloblastoma mouse models to their human counterparts. *Acta Neuropathol* 128:123-136, 2014

79. Northcott PA, Nakahara Y, Wu X, et al: Multiple recurrent genetic events converge on control of histone lysine methylation in medulloblastoma. *Nat Genet* 41:465-472, 2009

80. Pei Y, Liu KW, Wang J, et al: HDAC and PI3K antagonists cooperate to inhibit growth of MYC-driven medulloblastoma. *Cancer Cell* 29:311-323, 2016

81. Hill RM, Kuijper S, Lindsey JC, et al: Combined MYC and P53 defects emerge at medulloblastoma

relapse and define rapidly progressive, therapeutically targetable disease. *Cancer Cell* 27:72-84, 2015

82. Hanaford AR, Archer TC, Price A, et al: DIS-CoVERing innovative therapies for rare tumors: Combining genetically accurate disease models with in silico analysis to identify novel therapeutic targets. *Clin Cancer Res* 22:3903-3914, 2016

83. Morfouace M, Shelat A, Jacus M, et al: Pemetrexed and gemcitabine as combination therapy for the treatment of group3 medulloblastoma. *Cancer Cell* 25:516-529, 2014

84. Bandopadhyay P, Bergthold G, Nguyen B, et al: BET bromodomain inhibition of MYC-amplified medulloblastoma. *Clin Cancer Res* 20:912-925, 2014

85. Shih DJ, Northcott PA, Remke M, et al: Cytogenetic prognostication within medulloblastoma subgroups. *J Clin Oncol* 32:886-896, 2014

86. Korshunov A, Remke M, Kool M, et al: Biological and clinical heterogeneity of MYCN-amplified medulloblastoma. *Acta Neuropathol* 123:515-527, 2012

87. Schneider C, Ramaswamy V, Kulkarni AV, et al: Clinical implications of medulloblastoma subgroups: Incidence of CSF diversion surgery. *J Neurosurg Pediatr* 15:236-242, 2015

88. Northcott PA, Hielscher T, Dubuc A, et al: Pediatric and adult sonic hedgehog medulloblastomas are clinically and molecularly distinct. *Acta Neuropathol* 122:231-240, 2011

89. Gottardo NG, Hansford JR, McGlade JP, et al: Medulloblastoma Down Under 2013: A report from the third annual meeting of the International Medulloblastoma Working Group. *Acta Neuropathol* 127:189-201, 2014

90. Ellison DW, Dalton J, Kocak M, et al: Medulloblastoma: Clinicopathological correlates of SHH, WNT, and non-SHH/WNT molecular subgroups. *Acta Neuropathol* 121:381-396, 2011

91. Hovestadt V, Remke M, Kool M, et al: Robust molecular subgrouping and copy-number profiling of medulloblastoma from small amounts of archival tumour material using high-density DNA methylation arrays. *Acta Neuropathol* 125:913-916, 2013

92. Northcott PA, Shih DJ, Remke M, et al: Rapid, reliable, and reproducible molecular sub-grouping of clinical medulloblastoma samples. *Acta Neuropathol* 123:615-626, 2012

93. Gajjar A, Pizer B: Role of high-dose chemotherapy for recurrent medulloblastoma and other CNS primitive neuroectodermal tumors. *Pediatr Blood Cancer* 54:649-651, 2010

94. Sabel M, Fleischhack G, Tippelt S, et al: Relapse patterns and outcome after relapse in standard risk medulloblastoma: A report from the HIT-SIOP-PNET4 study. *J Neurooncol* 129:515-524, 2016

95. Morrissy AS, Garzia L, Shih DJ, et al: Divergent clonal selection dominates medulloblastoma at recurrence. *Nature* 529:351-357, 2016

96. King AA, Seidel K, Di C, et al: Long-term neurologic health and psychosocial function of adult survivors of childhood medulloblastoma/PNET: A report from the Childhood Cancer Survivor Study. *Neuro-oncol* 19:689-698, 2016

97. Yock TI, Yeap BY, Ebb DH, et al: Long-term toxic effects of proton radiotherapy for paediatric medulloblastoma: A phase 2 single-arm study. *Lancet Oncol* 17:287-298, 2016

98. Lannering B, Rutkowski S, Doz F, et al: Hyperfractionated versus conventional radiotherapy followed by chemotherapy in standard-risk medulloblastoma: Results from the randomized multicenter HIT-SIOP PNET 4 trial. *J Clin Oncol* 30:3187-3193, 2012

99. Packer RJ, Zhou T, Holmes E, et al: Survival and secondary tumors in children with medulloblastoma receiving radiotherapy and adjuvant chemotherapy: Results of Children's Oncology Group trial A9961. *Neuro-oncol* 15:97-103, 2013

100. Morris B, Partap S, Yeom K, et al: Cerebrovascular disease in childhood cancer survivors: A Children's Oncology Group report. *Neurology* 73:1906-1913, 2009

101. Moxon-Emre I, Bouffet E, Taylor MD, et al: Impact of craniospinal dose, boost volume, and

neurologic complications on intellectual outcome in patients with medulloblastoma. *J Clin Oncol* 32: 1760-1768, 2014

102. Câmara-Costa H, Resch A, Kieffer V, et al: Neuropsychological outcome of children treated for standard risk medulloblastoma in the PNET4 European randomized controlled trial of hyperfractionated versus standard radiation therapy and maintenance chemotherapy. *Int J Radiat Oncol Biol Phys* 92: 978-985, 2015

103. Riggs L, Piscione J, Laughlin S, et al: Exercise training for neural recovery in a restricted sample of pediatric brain tumor survivors: A controlled clinical trial with crossover of training versus no training. *Neuro-oncol* 19:440-450, 2017

104. Conklin HM, Ashford JM, Clark KN, et al: Long-term efficacy of computerized cognitive training among survivors of childhood cancer: A single-blind randomized controlled trial. *J Pediatr Psychol* 42: 220-231, 2017

Affiliations

All authors: Hospital for Sick Children, Toronto, Ontario, Canada.



Be the First to Hear When New Clinical Cancer Research Is Published Online



By signing up for JCO's Early Release Notification, you will be alerted and have access to new articles posted online every Monday, weeks before they appear in print. All Early Release articles are searchable and citable, and are posted on jco.org in advance of print publication.

Stay informed—sign up today at JCO.org!



American Society of Clinical Oncology

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Medulloblastoma: From Myth to Molecular

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/site/ifc.

Vijay Ramaswamy

No relationship to disclose

Michael D. Taylor

No relationship to disclose