

# Semisimple 2-categories

Ying Hong Tham

UHH

2022

# Goal

$$\{\text{multifusion category}\} \xrightarrow[\simeq]{\mathcal{M}\text{od}_{s.s.}^{fin}(-)} \{\text{finite semisimple 2-cat}\}$$

- ▶ The 2-category  $\mathcal{M}\text{od}_{s.s.}^{fin}(C)$  of finite semisimple module categories over a multifusion category  $C$  is semisimple.
- ▶ For any finite semisimple 2-category  $\mathcal{C}$ , there exists a multifusion category  $C$  such that  $\mathcal{C} \simeq \mathcal{M}\text{od}_{s.s.}^{fin}(C)$

# Overview

- ▶ direct sum

# Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff

# Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff
- ▶ simple objects

# Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff
- ▶ simple objects
- ▶ 'Schur lemma'

# Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff
- ▶ simple objects
- ▶ 'Schur lemma'
- ▶ main results

# Conventions

In relation to a 2-category:

- ▶  $\mathcal{C}$  (caligraphic font): 2-category;
- ▶  $X, Y, F$  (upper case latin): object of 2-category, functor between 2-categories;
- ▶  $f, g$  (lower case latin): 1-morphism; we write  $\mathcal{C}(X, Y)$  for the category of morphisms from  $X$  to  $Y$ ;
- ▶  $\eta, \varepsilon, \delta$  (lower case greek): 2-morphism; for a 2-morphism  $\alpha : f \Rightarrow g : X \rightarrow Y$ , we may write  $\alpha \in \mathcal{C}(X, Y)(f, g)$  to indicate its sources and targets, or simply  $\alpha \in \text{Hom}(f, g)$  if the objects are clear



# Conventions

In relation to a 1-category:

- ▶  $C, A$  (upper case latin): category;
- ▶  $a, b, f, g$  (lower case latin): objects in category, functor between categories;
- ▶  $\alpha, \beta$  (lower case greek): morphism in category

# Conventions

We also compose morphisms from right to left: in a 2-category  $\mathcal{C}$ , for  $\alpha \in \mathcal{C}(X, Y)(f, f')$ ,  $\beta \in \mathcal{C}(Y, Z)(g, g')$ ,  $\gamma \in \mathcal{C}(X, Y)(f', f'')$ , we write:

for composition of 1-morphisms,

$$g \circ f, g \circ f', \dots : X \rightarrow Z$$

for horizontal composition of 2-morphisms,

$$\beta \circ \alpha : (g \circ f) \Rightarrow (g' \circ f') : X \rightarrow Z$$

for vertical composition of 2-morphisms,

$$\gamma \cdot \alpha : f \Rightarrow f'' : X \rightarrow Y$$

# Conventions

In general, if  $P$  is a property of a 1-category, we say that a 2-category  $\mathcal{C}$  is *locally  $P$*  if every hom-category  $\mathcal{C}(X, Y)$  satisfies  $P$ .

# Conventions

In general, if  $P$  is a property of a 1-category, we say that a 2-category  $\mathcal{C}$  is *locally  $P$*  if every hom-category  $\mathcal{C}(X, Y)$  satisfies  $P$ . By 2-category we always mean a weak 2-category that is furthermore locally additive over  $\mathbf{k}$ . By 2-functor (sometimes just functor for simplicity) between 2-categories will always be locally  $\mathbf{k}$ -linear.

# Review

# Additive 2-category, direct sum of objects

# Additive 2-category, direct sum of objects

## Definition (direct sum of objects in 2-category)

A *direct sum* of two objects  $A_1, A_2$  in  $\mathcal{C}$  is an object  $A_1 \boxplus A_2$  together with inclusion and projection 1-morphisms

$i_k : A_k \Rightarrow A_1 \boxplus A_2 : p_k$ , such that

- ▶  $p_k \circ i_k \simeq \text{id}_{A_k}$ ,
- ▶  $p_2 \circ i_1, p_1 \circ i_2$  are zero 1-morphisms,
- ▶  $\text{id}_{A_1 \boxplus A_2} \simeq i_1 \circ p_1 \oplus i_2 \circ p_2$

# Additive 2-category, direct sum of objects

## Definition (direct sum of objects in 2-category)

A *direct sum* of two objects  $A_1, A_2$  in  $\mathcal{C}$  is an object  $A_1 \boxplus A_2$  together with inclusion and projection 1-morphisms

$i_k : A_k \Rightarrow A_1 \boxplus A_2 : p_k$ , such that

- ▶  $p_k \circ i_k \simeq \text{id}_{A_k}$ ,
- ▶  $p_2 \circ i_1, p_1 \circ i_2$  are zero 1-morphisms,
- ▶  $\text{id}_{A_1 \boxplus A_2} \simeq i_1 \circ p_1 \oplus i_2 \circ p_2$

## Definition (zero object)

A *zero object* in  $\mathcal{C}$  is an object  $0$  with trivial endomorphism category  $\mathcal{C}(0, 0)$  (has one object  $\text{id}_0$  with only identity morphism  $\text{id}_{\text{id}_0}$ ).



# Additive 2-category, direct sum of objects

## Definition (direct sum of objects in 2-category)

A *direct sum* of two objects  $A_1, A_2$  in  $\mathcal{C}$  is an object  $A_1 \boxplus A_2$  together with inclusion and projection 1-morphisms

$i_k : A_k \Rightarrow A_1 \boxplus A_2 : p_k$ , such that

- ▶  $p_k \circ i_k \simeq \text{id}_{A_k}$ ,
- ▶  $p_2 \circ i_1, p_1 \circ i_2$  are zero 1-morphisms,
- ▶  $\text{id}_{A_1 \boxplus A_2} \simeq i_1 \circ p_1 \oplus i_2 \circ p_2$

## Definition (zero object)

A *zero object* in  $\mathcal{C}$  is an object  $0$  with trivial endomorphism category  $\mathcal{C}(0, 0)$  (has one object  $\text{id}_0$  with only identity morphism  $\text{id}_{\text{id}_0}$ ).

## Definition

2-category  $\mathcal{C}$  is *additive* if finite direct sums of objects exist, has a zero object, (and is locally additive).

# Additive 2-category, direct sum of objects

## Definition (subobject)

A 1-morphism  $i : X \rightarrow Y$  is *fully faithful* (or  $(X, i)$  is a *subobject* of  $Y$ ) if it induces fully faithful functors between hom-categories by post-composition, i.e. for all objects  $A$ ,  $i \circ - : \mathcal{C}(A, X) \rightarrow \mathcal{C}(A, Y)$  is fully faithful.

# Additive 2-category, direct sum of objects

## Definition (subobject)

A 1-morphism  $i : X \rightarrow Y$  is *fully faithful* (or  $(X, i)$  is a *subobject* of  $Y$ ) if it induces fully faithful functors between hom-categories by post-composition, i.e. for all objects  $A$ ,  $i \circ - : \mathcal{C}(A, X) \rightarrow \mathcal{C}(A, Y)$  is fully faithful.

## Proposition

$i_k : A_k \rightarrow A_1 \boxplus A_2$  is fully faithful.

# Additive 2-category, direct sum of objects

## Definition (Direct sum of 2-categories)

Given additive 2-categories  $\mathcal{C}_j$ ,  $j \in J$ , we may consider the direct sum 2-category  $\mathcal{C} := \boxplus_{j \in J} \mathcal{C}_j$ :

- ▶  $\text{Obj } \mathcal{C} = \prod_{j \in J} \text{Obj } \mathcal{C}_j$

# Additive 2-category, direct sum of objects

## Definition (Direct sum of 2-categories)

Given additive 2-categories  $\mathcal{C}_j$ ,  $j \in J$ , we may consider the direct sum 2-category  $\mathcal{C} := \boxplus_{j \in J} \mathcal{C}_j$ :

- ▶  $\text{Obj } \mathcal{C} = \prod_{j \in J} \text{Obj } \mathcal{C}_j$
- ▶  $X_i \in \mathcal{C}_i$  becomes  $(\dots, 0, X_i, 0, \dots) \in \mathcal{C}$ .

# Additive 2-category, direct sum of objects

## Definition (Direct sum of 2-categories)

Given additive 2-categories  $\mathcal{C}_j$ ,  $j \in J$ , we may consider the direct sum 2-category  $\mathcal{C} := \boxplus_{j \in J} \mathcal{C}_j$ :

- ▶  $\text{Obj } \mathcal{C} = \prod_{j \in J} \text{Obj } \mathcal{C}_j$
- ▶  $X_i \in \mathcal{C}_i$  becomes  $(\dots, 0, X_i, 0, \dots) \in \mathcal{C}$ .
- ▶  $\mathcal{C}((X_i)_{i \in J}, (Y_j)_{j \in J}) = \bigoplus \mathcal{C}_i(X_i, Y_i)$

# Idempotent completeness, separable monads, splittings

# Idempotent completeness, separable monads, splittings

## Definition

A *separable algebra*  $(a, \mu, \eta)$  in a tensor category  $\mathcal{C}$  is an algebra that admits an  $a$ - $a$ -bimodule section  ${}_a a_a \rightarrow {}_a a \otimes a_a$  to  $\mu$ .



# Idempotent completeness, separable monads, splittings

## Definition

A *separable algebra*  $(a, \mu, \eta)$  in a tensor category  $\mathcal{C}$  is an algebra that admits an  $a$ - $a$ -bimodule section  ${}_a a_a \rightarrow {}_a a \otimes a_a$  to  $\mu$ .

## Definition

Let  $(t, \mu, \eta)$  be a monad on an object  $X$  in a 2-category  $\mathcal{C}$ . We say  $t$  is *separable* if there is a  $t$ - $t$ -bimodule section  $t \Rightarrow t \circ t$  to  $\mu$ .

# Idempotent completeness, separable monads, splittings

## Definition

A *separable algebra*  $(a, \mu, \eta)$  in a tensor category  $\mathcal{C}$  is an algebra that admits an  $a$ - $a$ -bimodule section  ${}_a a_a \rightarrow {}_a a \otimes a_a$  to  $\mu$ .

## Definition

Let  $(t, \mu, \eta)$  be a monad on an object  $X$  in a 2-category  $\mathcal{C}$ . We say  $t$  is *separable* if there is a  $t$ - $t$ -bimodule section  $t \Rightarrow t \circ t$  to  $\mu$ .

In other words, a separable monad over  $X$  is a separable algebra in  $\mathcal{C}(X, X)$ .

# Idempotent completeness, separable monads, splittings

## Definition

Let  $r \vdash l : X \rightarrow Y$  be an adjunction with unit  $\eta : \text{id}_X \Rightarrow rl$  and counit  $\varepsilon : lr \Rightarrow \text{id}_Y$ . We say the adjunction  $l \dashv r$  is *separable* if  $\varepsilon$  admits a section.

# Idempotent completeness, separable monads, splittings

## Definition

Let  $r \vdash l : X \rightarrow Y$  be an adjunction with unit  $\eta : \text{id}_X \Rightarrow rl$  and counit  $\varepsilon : lr \Rightarrow \text{id}_Y$ . We say the adjunction  $l \dashv r$  is *separable* if  $\varepsilon$  admits a section.

Clearly, if an adjunction  $l \dashv r$  is separable, then the monad  $rl$  is separable - the splitting gives  $rl \simeq r(\text{id}_Y)l \Rightarrow r(lr)l \simeq (rl)(rl)$

# Idempotent completeness, separable monads, splittings

## Definition

Let  $r \vdash l : X \rightarrow Y$  be an adjunction with unit  $\eta : \text{id}_X \Rightarrow rl$  and counit  $\varepsilon : lr \Rightarrow \text{id}_Y$ . We say the adjunction  $l \dashv r$  is *separable* if  $\varepsilon$  admits a section.

Clearly, if an adjunction  $l \dashv r$  is separable, then the monad  $rl$  is separable - the splitting gives  $rl \simeq r(\text{id}_Y)l \Rightarrow r(lr)l \simeq (rl)(rl)$

## Definition

Let  $(t, \mu, \eta)$  be a separable monad on an object  $X \in \mathcal{C}$ . A *(separable) splitting* of  $t$  is a (separable) adjunction  $r \vdash l : X \rightarrow Y$  together with an isomorphism  $\psi : rl \simeq t$  as monads on  $X$ .

# Idempotent completeness, separable monads, splittings

## Definition

Let  $r \vdash l : X \rightarrow Y$  be an adjunction with unit  $\eta : \text{id}_X \Rightarrow rl$  and counit  $\varepsilon : lr \Rightarrow \text{id}_Y$ . We say the adjunction  $l \dashv r$  is *separable* if  $\varepsilon$  admits a section.

Clearly, if an adjunction  $l \dashv r$  is separable, then the monad  $rl$  is separable - the splitting gives  $rl \simeq r(\text{id}_Y)l \Rightarrow r(lr)l \simeq (rl)(rl)$

## Definition

Let  $(t, \mu, \eta)$  be a separable monad on an object  $X \in \mathcal{C}$ . A *(separable) splitting* of  $t$  is a (separable) adjunction  $r \vdash l : X \rightarrow Y$  together with an isomorphism  $\psi : rl \simeq t$  as monads on  $X$ .

## Proposition (Uniqueness of splitting)

*[[1], Theorem A.3.1] In locally idempotent complete 2-category  $\mathcal{C}$ , separable splitting of monad, if exists, is unique up to equivalence.*

# Idempotent completeness, 2-category

## Definition

A 2-category  $\mathcal{C}$  is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

# Idempotent completeness, 2-category

## Definition

A 2-category  $\mathcal{C}$  is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

## Definition (Idempotent completion)

Let  $\mathcal{C}$  be a locally idempotent complete 2-category. The *idempotent completion of  $\mathcal{C}$* , denoted  $\mathcal{C}^\nabla$ , with:

- ▶ Objects:  $(X, p)$  separable monad in  $\mathcal{C}$ ,
- ▶  $\mathcal{C}^\nabla((X, p), (Y, q)) = q\text{-bimod-}p(\mathcal{C}(X, Y))$



# Idempotent completeness, 2-category

## Definition

A 2-category  $\mathcal{C}$  is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

## Definition (Idempotent completion)

Let  $\mathcal{C}$  be a locally idempotent complete 2-category. The *idempotent completion of  $\mathcal{C}$* , denoted  $\mathcal{C}^\nabla$ , with:

- ▶ Objects:  $(X, p)$  separable monad in  $\mathcal{C}$ ,
- ▶  $\mathcal{C}^\nabla((X, p), (Y, q)) = q\text{-bimod-}p(\mathcal{C}(X, Y))$

There is a natural 2-functor  $I : \mathcal{C} \rightarrow \mathcal{C}^\nabla$  that is fully faithful.

# Idempotent completeness, 2-category

## Definition

A 2-category  $\mathcal{C}$  is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

## Definition (Idempotent completion)

Let  $\mathcal{C}$  be a locally idempotent complete 2-category. The *idempotent completion* of  $\mathcal{C}$ , denoted  $\mathcal{C}^\nabla$ , with:

- ▶ Objects:  $(X, p)$  separable monad in  $\mathcal{C}$ ,
- ▶  $\mathcal{C}^\nabla((X, p), (Y, q)) = q\text{-bimod-}p(\mathcal{C}(X, Y))$

There is a natural 2-functor  $I : \mathcal{C} \rightarrow \mathcal{C}^\nabla$  that is fully faithful.

A 2-functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  extends to a 2-functor  $F^\nabla : \mathcal{C}^\nabla \rightarrow \mathcal{D}^\nabla$  that commutes with  $I$ 's.

# Idempotent completeness, 2-category

Key example:

$\mathcal{BC}$ : one object  $*$  with endomorphism category  $\mathcal{BC}(*, *) = C$   
(with  $x \otimes y = x \circ y$ )

$$(\mathcal{BC})^\nabla = \begin{cases} \text{Obj} : \text{separable algebras in } C \\ \text{Mor} : (\mathcal{BC})^\nabla(a, b) = b\text{-bimod-}a(C) \end{cases}$$

# Idempotent completeness, 2-category

## Proposition

$\mathcal{C}^\nabla$  is idempotent complete. Moreover, if  $\mathcal{C}$  is already idempotent complete, then  $I : \mathcal{C} \simeq \mathcal{C}^\nabla$  is an equivalence.

# Idempotent completeness, 2-category

## Proposition

$\mathcal{C}^\nabla$  is idempotent complete. Moreover, if  $\mathcal{C}$  is already idempotent complete, then  $I : \mathcal{C} \simeq \mathcal{C}^\nabla$  is an equivalence.

As a consequence, if  $\mathcal{D}$  is idempotent complete, then we have equivalences of 2-categories

$$\mathcal{F}un(\mathcal{C}, \mathcal{D}) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D}^\nabla) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D})$$

# Idempotent completeness, 2-category

## Proposition

$\mathcal{C}^\nabla$  is idempotent complete. Moreover, if  $\mathcal{C}$  is already idempotent complete, then  $I : \mathcal{C} \simeq \mathcal{C}^\nabla$  is an equivalence.

As a consequence, if  $\mathcal{D}$  is idempotent complete, then we have equivalences of 2-categories

$$\mathcal{F}un(\mathcal{C}, \mathcal{D}) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D}^\nabla) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D})$$

## Proposition

If  $F : \mathcal{C} \rightarrow \mathcal{D}$  is fully faithful, then  $F^\nabla : \mathcal{C}^\nabla \rightarrow \mathcal{D}^\nabla$  is also fully faithful.

# Idempotent completeness, 2-category

## Proposition ([1]Prop 1.3.13)

*For a multifusion category  $C$ , the following 2-functor is an equivalence:*

$$(-)\text{-mod}(C) : (\mathcal{BC})^{\nabla} \rightarrow \mathcal{Mod}(C)$$

$$a \mapsto a\text{-mod}(C)$$

$${}_b m_a \mapsto m \otimes_a -$$

$$\varphi \mapsto \varphi \otimes_a -$$

# Idempotent completeness, 2-category

Proof.

Essential surjectivity follows from:

-[2]Theorem 1: for a finite semisimple right module category over multifusion  $C$ , there exists a semisimple algebra  $a$  in  $C$  such that  $M \simeq a\text{-mod}(C)$  as right  $C$ -module categories; and



# Idempotent completeness, 2-category

## Proof.

Essential surjectivity follows from:

- [2]Theorem 1: for a finite semisimple right module category over multifusion  $C$ , there exists a semisimple algebra  $a$  in  $C$  such that  $M \simeq a\text{-mod}(C)$  as right  $C$ -module categories; and
- [4]Corollary 2.6.9: When  $C$  is multifusion over a field of characteristic 0, a right  $C$ -module category  $M$  is separable ( $\simeq a\text{-mod}(C)$  for a separable algebra  $a$ ) if and only if it is semisimple.

# Idempotent completeness, 2-category

## Proof.

Essential surjectivity follows from:

- [2]Theorem 1: for a finite semisimple right module category over multifusion  $C$ , there exists a semisimple algebra  $a$  in  $C$  such that  $M \simeq a\text{-mod}(C)$  as right  $C$ -module categories; and
- [4]Corollary 2.6.9: When  $C$  is multifusion over a field of characteristic 0, a right  $C$ -module category  $M$  is separable ( $\simeq a\text{-mod}(C)$  for a separable algebra  $a$ ) if and only if it is semisimple.

Fully faithfulness:

# Idempotent completeness, 2-category

## Proof.

Essential surjectivity follows from:

- [2]Theorem 1: for a finite semisimple right module category over multifusion  $C$ , there exists a semisimple algebra  $a$  in  $C$  such that  $M \simeq a\text{-mod}(C)$  as right  $C$ -module categories; and
- [4]Corollary 2.6.9: When  $C$  is multifusion over a field of characteristic 0, a right  $C$ -module category  $M$  is separable ( $\simeq a\text{-mod}(C)$  for a separable algebra  $a$ ) if and only if it is semisimple.

Fully faithfulness: on free modules:  $F(a \otimes c) \simeq F(a) \otimes c$ ;

# Idempotent completeness, 2-category

## Proof.

Essential surjectivity follows from:

-[2]Theorem 1: for a finite semisimple right module category over multifusion  $C$ , there exists a semisimple algebra  $a$  in  $C$  such that  $M \simeq a\text{-mod}(C)$  as right  $C$ -module categories; and

-[4]Corollary 2.6.9: When  $C$  is multifusion over a field of characteristic 0, a right  $C$ -module category  $M$  is separable ( $\simeq a\text{-mod}(C)$  for a separable algebra  $a$ ) if and only if it is semisimple.

Fully faithfulness: on free modules:  $F(a \otimes c) \simeq F(a) \otimes c$ ; (right)

exactness:  $F(m) = F(a \otimes_a m) \simeq F(a) \otimes_a m$

# Idempotent completeness, 2-category

## Proof.

Essential surjectivity follows from:

-[2]Theorem 1: for a finite semisimple right module category over multifusion  $C$ , there exists a semisimple algebra  $a$  in  $C$  such that  $M \simeq a\text{-mod}(C)$  as right  $C$ -module categories; and

-[4]Corollary 2.6.9: When  $C$  is multifusion over a field of characteristic 0, a right  $C$ -module category  $M$  is separable ( $\simeq a\text{-mod}(C)$  for a separable algebra  $a$ ) if and only if it is semisimple.

Fully faithfulness: on free modules:  $F(a \otimes c) \simeq F(a) \otimes c$ ; (right)

exactness:  $F(m) = F(a \otimes_a m) \simeq F(a) \otimes_a m$  ([5]Prop 7.11.1) □

# Idempotent completeness, 2-category

Slightly more detailed proof.

$$M_C \simeq \mathbf{a}\text{-mod}(C):$$

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$   
to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$



# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ;

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

$\mathbf{1} \in C$  is projective,

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

$1 \in C$  is projective,  $a \in a\text{-mod}(C)$  is projective,

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

$1 \in C$  is projective,  $a \in a\text{-mod}(C)$  is projective, free modules  $a \otimes c$  are projective



# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

$1 \in C$  is projective,  $a \in a\text{-mod}(C)$  is projective, free modules  $a \otimes c$  are projective

for left  $a$ -module  $m$ ,

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

$1 \in C$  is projective,  $a \in a\text{-mod}(C)$  is projective, free modules  $a \otimes c$  are projective

for left  $a$ -module  $m$ , separability of  $a$  gives section to quotient  $a \otimes m \rightarrow a \otimes_a m \simeq m$ , so  $m$  is projective.

# Idempotent completeness, 2-category

Slightly more detailed proof.

$M_C \simeq a\text{-mod}(C)$ : use  $C$ -action on  $M$  to enrich  $M(m_1, m_2) \in \text{Vec}$  to object  $\underline{\text{Hom}}(m_1, m_2) \in C$ , “internal hom”

$$C(c, \underline{\text{Hom}}(m_1, m_2)) \simeq M(m_1 \triangleleft c, m_2)$$

fix some  $x \in M$ ;  $a := \underline{\text{Hom}}(x, x)$ ; composition gives  $a$ -module structure on  $\underline{\text{Hom}}(x, m)$

$$M \ni m \mapsto \underline{\text{Hom}}(x, m) \in a\text{-mod}(C)$$

separable  $\Rightarrow$  semisimple:

$1 \in C$  is projective,  $a \in a\text{-mod}(C)$  is projective, free modules  $a \otimes c$  are projective

for left  $a$ -module  $m$ , separability of  $a$  gives section to quotient  $a \otimes m \rightarrow a \otimes_a m \simeq m$ , so  $m$  is projective.

semisimple  $\Rightarrow$  separable is more challenging



# Simple objects

# Simple objects

## Proposition (equivalent notions of simple-ness)

*Let  $\mathcal{C}$  be an idempotent complete and locally finite semisimple 2-category, and let  $X \in \mathcal{C}$  be a nonzero object. Then the following notions of  $X$  being simple are equivalent:*

# Simple objects

## Proposition (equivalent notions of simple-ness)

*Let  $\mathcal{C}$  be an idempotent complete and locally finite semisimple 2-category, and let  $X \in \mathcal{C}$  be a nonzero object. Then the following notions of  $X$  being simple are equivalent:*

*(1) any subobject  $i : Y \rightarrow X$  of  $X$  is either 0 or an equivalence;*

# Simple objects

## Proposition (equivalent notions of simple-ness)

*Let  $\mathcal{C}$  be an idempotent complete and locally finite semisimple 2-category, and let  $X \in \mathcal{C}$  be a nonzero object. Then the following notions of  $X$  being simple are equivalent:*

- (1) any subobject  $i : Y \rightarrow X$  of  $X$  is either 0 or an equivalence;*
- (2)  $X$  cannot be written as a non-trivial direct sum, i.e. if  $X = \boxplus X_i$ , then  $X_i \simeq 0$  for all but one  $i$ ;*

# Simple objects

## Proposition (equivalent notions of simple-ness)

*Let  $\mathcal{C}$  be an idempotent complete and locally finite semisimple 2-category, and let  $X \in \mathcal{C}$  be a nonzero object. Then the following notions of  $X$  being simple are equivalent:*

- (1) any subobject  $i : Y \rightarrow X$  of  $X$  is either 0 or an equivalence;*
- (2)  $X$  cannot be written as a non-trivial direct sum, i.e. if  $X = \boxplus X_i$ , then  $X_i \simeq 0$  for all but one  $i$ ;*
- (3)  $\text{id}_X$  is a simple object in  $\mathcal{C}(X, X)$ .*



# Simple objects

Proof sketch.

(1)  $\Rightarrow$  (2): Contravariant statement follows from fully faithfulness of  $i_k : A_k \rightarrow A_1 \boxplus A_2$ .

# Simple objects

## Proof sketch.

(1)  $\Rightarrow$  (2): Contravariant statement follows from fully faithfulness of  $i_k : A_k \rightarrow A_1 \boxplus A_2$ .

(2)  $\Rightarrow$  (3): Contravariant statement is “identity splitting implies object splitting”, uses idempotent completeness of  $\mathcal{C}$  to split out objects corresponding to summands of  $\text{id}_X$  (which are separable monads) (see [1]Prop 1.3.16).

# Simple objects

## Proof sketch.

(1)  $\Rightarrow$  (2): Contravariant statement follows from fully faithfulness of  $i_k : A_k \rightarrow A_1 \boxplus A_2$ .

(2)  $\Rightarrow$  (3): Contravariant statement is “identity splitting implies object splitting”, uses idempotent completeness of  $\mathcal{C}$  to split out objects corresponding to summands of  $\text{id}_X$  (which are separable monads) (see [1]Prop 1.3.16).

(3)  $\Rightarrow$  (1): for non-zero fully faithful  $r : Y \rightarrow X$ , with  $\text{id}_X$  simple, consider the left adjoint  $l : X \rightarrow Y$ , use fully faithfulness to get a preimage  $\delta : \text{id}_Y \Rightarrow lr$  of  $\eta \circ r : r \Rightarrow rlr$ . Show  $\delta$  is a section of the counit. Etc. (See [1]Prop 1.2.14) □

# Semisimple 2-category

Definition ((finite) semisimple 2-category)

# Semisimple 2-category

## Definition ((finite) semisimple 2-category)

A 2-category  $\mathcal{C}$  is *semisimple* if it is:

- ▶ locally semisimple,
- ▶ admits left and right adjoints for every 1-morphism,
- ▶ additive,
- ▶ idempotent complete.

It is furthermore *finite semisimple* if it is also locally finite and has finitely many equivalence classes of simple objects.


# New Stuff

# Schur's lemma

The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case.

# Schur's lemma


The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case. However, recall  $\mathcal{M}\text{od}(\text{Vec}_{\mathbb{Z}/2})$ :





# Schur's lemma


The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case. However, recall  $\mathcal{M}\text{od}(\text{Vec}_{\mathbb{Z}/2})$ :



There can be nonzero morphisms between non-equivalent simple objects - weird!

## Schur's lemma

The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case. However, recall  $\mathcal{M}\text{od}(\text{Vec}_{\mathbb{Z}/2})$ :



There can be nonzero morphisms between non-equivalent simple objects - weird!

“2-Morita equivalences” between fusion categories

# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

Proof.

$$A \xrightarrow{\quad f \quad} B \xrightarrow{\quad g \quad} C$$




# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

Proof.




# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

Proof.




# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

Proof.



$$\text{id}_{\text{id}_B} = \varepsilon \cdot \delta$$




# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

Proof.



$$\text{id}_{\text{id}_B} = \epsilon \cdot \delta$$

$$\text{id}_g = (\text{id}_g \circ \epsilon) \cdot (\text{id}_g \circ \delta) : g \Rightarrow gff^* \Rightarrow g$$






# Schur's lemma

## Proposition (Schur's Lemma, [1]Prop 1.2.19)

*In a semisimple 2-category  $\mathcal{C}$ , if  $f : A \rightarrow B, g : B \rightarrow C$  are nonzero 1-morphisms between simple objects  $A, B, C$ , then  $g \circ f$  is also nonzero.*

Proof.



$$\mathrm{id}_{\mathrm{id}_B} = \varepsilon \cdot \delta$$

$$\mathrm{id}_g = (\mathrm{id}_g \circ \varepsilon) \cdot (\mathrm{id}_g \circ \delta) : g \Rightarrow g f f^* \Rightarrow g$$

$$\text{if } g f \cong 0, \text{ then } \mathrm{id}_g = 0, \text{ and } g \cong 0$$



# Schur's lemma - components

## Definition (component of semisimple 2-category)

In a semisimple 2-category  $\mathcal{C}$ , we say two simple objects  $A, B$  belong to the same *component* if there exists a nonzero 1-morphism  $f : A \rightarrow B$ .

# Schur's lemma - components

## Definition (component of semisimple 2-category)

In a semisimple 2-category  $\mathcal{C}$ , we say two simple objects  $A, B$  belong to the same *component* if there exists a nonzero 1-morphism  $f : A \rightarrow B$ .

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

# Schur's lemma - components

## Definition (component of semisimple 2-category)

In a semisimple 2-category  $\mathcal{C}$ , we say two simple objects  $A, B$  belong to the same *component* if there exists a nonzero 1-morphism  $f : A \rightarrow B$ .

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

In a finite semisimple 2-category  $\mathcal{C}$ , there will be finite set  $J$  of components;

# Schur's lemma - components

## Definition (component of semisimple 2-category)

In a semisimple 2-category  $\mathcal{C}$ , we say two simple objects  $A, B$  belong to the same *component* if there exists a nonzero 1-morphism  $f : A \rightarrow B$ .

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

In a finite semisimple 2-category  $\mathcal{C}$ , there will be finite set  $J$  of components;

full subcategory  $\mathcal{C}_j$ ,  $j \in J$ : direct sum of simple objects in  $j$

# Schur's lemma - components

## Definition (component of semisimple 2-category)

In a semisimple 2-category  $\mathcal{C}$ , we say two simple objects  $A, B$  belong to the same *component* if there exists a nonzero 1-morphism  $f : A \rightarrow B$ .

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

In a finite semisimple 2-category  $\mathcal{C}$ , there will be finite set  $J$  of components;

full subcategory  $\mathcal{C}_j$ ,  $j \in J$ : direct sum of simple objects in  $j$   
gives direct sum decomposition  $\mathcal{C} \simeq \bigoplus_{j \in J} \mathcal{C}_j$ .

# Main results

## Theorem ([1]Theorem 1.4.8)

*The 2-category of finite semisimple module categories of a multifusion category  $\mathcal{C}$  is a finite semisimple 2-category.*

# Main results

## Theorem ([1]Theorem 1.4.8)

*The 2-category of finite semisimple module categories of a multifusion category  $C$  is a finite semisimple 2-category.*

## Proof.

$\mathcal{M}\mathrm{od}(C) \simeq (\mathcal{B}C)^\nabla$  is idempotent complete.  $\mathcal{M}\mathrm{od}(C)$  is clearly already additive.

Locally semisimple-ness follows directly from [4]Corollary 2.5.6;



# Main results

## Theorem ([1]Theorem 1.4.8)

*The 2-category of finite semisimple module categories of a multifusion category  $\mathcal{C}$  is a finite semisimple 2-category.*

## Proof.

$\mathcal{M}\mathrm{od}(\mathcal{C}) \simeq (\mathcal{B}\mathcal{C})^\nabla$  is idempotent complete.  $\mathcal{M}\mathrm{od}(\mathcal{C})$  is clearly already additive.

Locally semisimple-ness follows directly from [4]Corollary 2.5.6; relative Deligne tensor product of separable module categories is also separable

# Main results

## Theorem ([1]Theorem 1.4.8)

*The 2-category of finite semisimple module categories of a multifusion category  $C$  is a finite semisimple 2-category.*

### Proof.

$\mathcal{M}\mathrm{od}(C) \simeq (\mathcal{B}C)^\nabla$  is idempotent complete.  $\mathcal{M}\mathrm{od}(C)$  is clearly already additive.

Locally semisimple-ness follows directly from [4]Corollary 2.5.6; relative Deligne tensor product of separable module categories is also separable

Existence of adjoints for 1-morphisms follows from [3]Corollary 2.13: left/right adjoint to underlying functor of a  $C$ -module functor also has  $C$ -module structure.

# Main results

## Theorem ([1]Theorem 1.4.8)

*The 2-category of finite semisimple module categories of a multifusion category  $C$  is a finite semisimple 2-category.*

## Proof.

$\text{Mod}(C) \simeq (\mathcal{BC})^\nabla$  is idempotent complete.  $\text{Mod}(C)$  is clearly already additive.

Locally semisimple-ness follows directly from [4]Corollary 2.5.6; relative Deligne tensor product of separable module categories is also separable

Existence of adjoints for 1-morphisms follows from [3]Corollary 2.13: left/right adjoint to underlying functor of a  $C$ -module functor also has  $C$ -module structure.

(explicit right adjoint to  ${}_b m_a : a\text{-mod}(C) \rightarrow b\text{-mod}(C)$  is given by  $\underline{\text{Hom}}_B(m, -)$ )



# Main results

## Theorem ([1]Theorem 1.4.9)

*Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.*

## Proof.

Assume that  $\mathcal{C}$  has only one component.

# Main results

## Theorem ([1]Theorem 1.4.9)

*Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.*

## Proof.

Assume that  $\mathcal{C}$  has only one component.

( $\mathcal{C} \simeq \boxplus_{j \in J} \mathcal{C}_j$  and  $\mathcal{C}_j \simeq \text{Mod}(C_j)$  for multifusion  $C_j$ , then  
 $\mathcal{C} \simeq \text{Mod}(\oplus_{j \in J} C_j)$ .)

# Main results

## Theorem ([1]Theorem 1.4.9)

*Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.*

### Proof.

Assume that  $\mathcal{C}$  has only one component.

( $\mathcal{C} \simeq \boxplus_{j \in J} \mathcal{C}_j$  and  $\mathcal{C}_j \simeq \text{Mod}(C_j)$  for multifusion  $C_j$ , then  
 $\mathcal{C} \simeq \text{Mod}(\oplus_{j \in J} C_j)$ .)

Fix simple object  $X$ , let  $C = \mathcal{C}(X, X)$ .

# Main results

## Theorem ([1]Theorem 1.4.9)

*Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.*

### Proof.

Assume that  $\mathcal{C}$  has only one component.

( $\mathcal{C} \simeq \boxplus_{j \in J} \mathcal{C}_j$  and  $\mathcal{C}_j \simeq \text{Mod}(C_j)$  for multifusion  $C_j$ , then  
 $\mathcal{C} \simeq \text{Mod}(\oplus_{j \in J} C_j)$ .)

Fix simple object  $X$ , let  $C = \mathcal{C}(X, X)$ .

We show  $\mathcal{C} \simeq (\mathcal{B}C)^\nabla$ .

# Main results

## Proof (Cont.)

Consider the inclusion 2-functor

$$\begin{aligned} F : \mathcal{BC} &\rightarrow \mathcal{C} \\ * &\mapsto X \end{aligned}$$

which is fully faithful by construction.



# Main results

## Proof (Cont.)

Consider the inclusion 2-functor

$$\begin{aligned} F : \mathcal{BC} &\rightarrow \mathcal{C} \\ * &\mapsto X \end{aligned}$$

which is fully faithful by construction. Then

$$F^{\nabla} : (\mathcal{BC})^{\nabla} \rightarrow \mathcal{C}$$

is fully faithful.

# Main results

## Proof (Cont.)

Consider the inclusion 2-functor

$$\begin{aligned} F : \mathcal{BC} &\rightarrow \mathcal{C} \\ * &\mapsto X \end{aligned}$$

which is fully faithful by construction. Then

$$F^\nabla : (\mathcal{BC})^\nabla \rightarrow \mathcal{C}$$

is fully faithful. Remains to show other simples are in essential image.

# Main results

Proof (Cont.)


$Y$

$$* \xrightarrow{F^\nabla} X$$




# Main results

Proof (Cont.)




# Main results

## Proof (Cont.)



# Main results

## Proof (Cont.)




$\text{id}_Y$  simple  
 $\varepsilon$  has section  
 $\Rightarrow$  sep. adj.



# Main results

## Proof (Cont.)




$\text{id}_Y$  simple  
 $\varepsilon$  has section  
 $\Rightarrow$  sep. adj.



# Main results

## Proof (Cont.)




$\text{id}_Y$  simple  
 $\varepsilon$  has section  
 $\Rightarrow$  sep. adj.





# Main results

## Proof (Cont.)



$\text{id}_Y$  simple  
 $\varepsilon$  has section  
 $\Rightarrow$  sep. adj.



# Main results

## Proof (Cont.)

$$\begin{array}{ccccc} (*, gf) & \xrightarrow{F^\nabla} & F^\nabla(*, gf) & \xrightarrow{\cong} & Y \\ & \nwarrow & \nwarrow & \nearrow f & \nearrow g \\ & & * & \xrightarrow{F^\nabla} & X \end{array}$$

Diagram illustrating the proof structure. The top row shows a sequence of maps:  $(*, gf) \xrightarrow{F^\nabla} F^\nabla(*, gf) \xrightarrow{\cong} Y$ . The bottom row shows  $* \xrightarrow{F^\nabla} X$ . A curved arrow labeled  $gf$  points from  $*$  to  $(*, gf)$ . A curved arrow labeled  $f$  points from  $X$  to  $F^\nabla(*, gf)$ . A curved arrow labeled  $g$  points from  $X$  to  $Y$ . There is also a curved arrow labeled  $\dashv$  from  $Y$  to  $X$ .

$\text{id}_Y$  simple  
 $\varepsilon$  has section  
 $\Rightarrow$  sep. adj.



# Main results

The only property of  $X$  that we used is the fact that there exists a nonzero 1-morphism from  $X$  to every simple in  $\mathcal{C}$ , and thus any nonzero  $X$  will do.

# Main results

The only property of  $X$  that we used is the fact that there exists a nonzero 1-morphism from  $X$  to every simple in  $\mathcal{C}$ , and thus any nonzero  $X$  will do. Taking, say,  $X = \boxplus X_i$ , where the sum is over equivalence classes of simples, would result in a multifusion  $\mathcal{C} = \mathcal{C}(X, X)$ .



# Main results

The only property of  $X$  that we used is the fact that there exists a nonzero 1-morphism from  $X$  to every simple in  $\mathcal{C}$ , and thus any nonzero  $X$  will do. Taking, say,  $X = \boxplus X_i$ , where the sum is over equivalence classes of simples, would result in a multifusion  $\mathcal{C} = \mathcal{C}(X, X)$ .


We can also avoid the first step of taking only one component of  $\mathcal{C}$ , take  $X$  with at least one object from each  $\mathcal{C}_j$  in its direct sum decomposition.

# Main results

Every object contains within them the data to reconstruct the whole 2-category.



# Indra's Net of Pearls (or $T^3 \# T^3$ )



## Example

$C = \text{Vec}_G$  for a finite group  $G$ .



## Example

$C = \text{Vec}_G$  for a finite group  $G$ .

$e$ : trivial algebra  $\mathbf{k}_e$      $e\text{-mod}(C) \simeq C_C$

# Example

$C = \text{Vec}_G$  for a finite group  $G$ .

$e$ : trivial algebra  $\mathbf{k}_e$      $e\text{-mod}(C) \simeq C_C$

$f$ : the group algebra  $\mathbf{k}[G] = \bigoplus \mathbf{k}_g$      $f\text{-mod}(C) \simeq \text{Vec}$   
(right action on  $\text{Vec}$  = forget  $G$ -grading)

## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

For  $f\text{-mod}(C)$ , an endofunctor is given by some  $m \in f\text{-bimod-}f(C)$ ; write  $m = \bigoplus m_g$ .

## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

For  $f\text{-mod}(C)$ , an endofunctor is given by some  $m \in f\text{-bimod-}f(C)$ ; write  $m = \bigoplus m_g$ . The right  $f$ -action on  $m$  makes all the  $m_g$  isomorphic in a coherent manner.

## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

For  $f\text{-mod}(C)$ , an endofunctor is given by some  $m \in f\text{-bimod-}f(C)$ ; write  $m = \bigoplus m_g$ . The right  $f$ -action on  $m$  makes all the  $m_g$  isomorphic in a coherent manner. For  $h \in G$ , conjugation (left action by  $h$  and right action by  $h^{-1}$ ) gives an action of  $G$  on  $m_e$ ; determines  $m$  completely.

## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

For  $f\text{-mod}(C)$ , an endofunctor is given by some  $m \in f\text{-bimod-}f(C)$ ; write  $m = \bigoplus m_g$ . The right  $f$ -action on  $m$  makes all the  $m_g$  isomorphic in a coherent manner. For  $h \in G$ , conjugation (left action by  $h$  and right action by  $h^{-1}$ ) gives an action of  $G$  on  $m_e$ ; determines  $m$  completely.

Thus,  $\mathcal{M}\text{od}(C)(f, f) \simeq f\text{-bimod-}f(C) \simeq \text{Rep}(G)$ .

## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

For  $f\text{-mod}(C)$ , an endofunctor is given by some  $m \in f\text{-bimod-}f(C)$ ; write  $m = \bigoplus m_g$ . The right  $f$ -action on  $m$  makes all the  $m_g$  isomorphic in a coherent manner. For  $h \in G$ , conjugation (left action by  $h$  and right action by  $h^{-1}$ ) gives an action of  $G$  on  $m_e$ ; determines  $m$  completely.

Thus,  $\text{Mod}(C)(f, f) \simeq f\text{-bimod-}f(C) \simeq \text{Rep}(G)$ .

(Alternatively, one can also check that  $C$ -module structure on  $m : \text{Vec} \rightarrow \text{Vec}$  amounts to  $G$ -action on  $m(\mathbf{k}).$ )



## Example

Next we study the functor categories. Clearly the  $C$ -endofunctors of  $C_C$  is  $C$  itself.

For  $f\text{-mod}(C)$ , an endofunctor is given by some  $m \in f\text{-bimod-}f(C)$ ; write  $m = \bigoplus m_g$ . The right  $f$ -action on  $m$  makes all the  $m_g$  isomorphic in a coherent manner. For  $h \in G$ , conjugation (left action by  $h$  and right action by  $h^{-1}$ ) gives an action of  $G$  on  $m_e$ ; determines  $m$  completely.

Thus,  $\text{Mod}(C)(f, f) \simeq f\text{-bimod-}f(C) \simeq \text{Rep}(G)$ .

(Alternatively, one can also check that  $C$ -module structure on  $m: \text{Vec} \rightarrow \text{Vec}$  amounts to  $G$ -action on  $m(\mathbf{k})$ .)

Thus we have:

$$\text{Mod}(\text{Vec}_G) \simeq \text{Mod}(\text{Rep}(G))$$

## Example

For functor categories between them, we see that

$$\mathcal{M}\mathrm{od}(C)(e, f) \simeq f\text{-bimod-}e(C) \simeq f\text{-mod}(C) \simeq \mathbf{Vec}$$

$$\mathcal{M}\mathrm{od}(C)(f, e) \simeq e\text{-bimod-}f(C) \simeq \mathrm{mod-}f(C) \simeq \mathbf{Vec}$$

## Example

For functor categories between them, we see that

$$\mathcal{M}\mathrm{od}(C)(e, f) \simeq f\text{-bimod-}e(C) \simeq f\text{-mod}(C) \simeq \mathrm{Vec}$$

$$\mathcal{M}\mathrm{od}(C)(f, e) \simeq e\text{-bimod-}f(C) \simeq \mathrm{mod-}f(C) \simeq \mathrm{Vec}$$

Note for  $G = \mathbb{Z}/2$ ,  $\mathrm{Vec}_{\mathbb{Z}/2} \simeq \mathrm{Rep}(\mathbb{Z}/2)$ , don't see difference in endomorphism categories.

## Example

For functor categories between them, we see that






$$\mathcal{M}\mathrm{od}(C)(e, f) \simeq f\text{-bimod-}e(C) \simeq f\text{-mod}(C) \simeq \mathrm{Vec}$$

$$\mathcal{M}\mathrm{od}(C)(f, e) \simeq e\text{-bimod-}f(C) \simeq \mathrm{mod-}f(C) \simeq \mathrm{Vec}$$

Note for  $G = \mathbb{Z}/2$ ,  $\mathrm{Vec}_{\mathbb{Z}/2} \simeq \mathrm{Rep}(\mathbb{Z}/2)$ , don't see difference in endomorphism categories.

Of course, there are many other objects in  $\mathcal{M}\mathrm{od}(C)$ , e.g. for a subgroup  $H \subseteq G$ , have group algebra  $\mathbf{k}[H]$ .

End!

-  Douglas, Christopher L., and David J. Reutter. “Fusion 2-categories and a state-sum invariant for 4-manifolds.” arXiv preprint arXiv:1812.11933 (2018).
-  Ostrik, Victor. “Module categories, weak Hopf algebras and modular invariants.” Transformation groups 8, no. 2 (2003): 177-206.
-  C. L. Douglas, C. Schommer-Pries, and N. Snyder. The balanced tensor product of module categories. Kyoto J. Math., 2017. arXiv:1406.4204.
-  C. L. Douglas, C. Schommer-Pries, and N. Snyder. Dualizable tensor categories. Mem. Amer. Math. Soc., 2017. arXiv:1312.7188.
-  P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. Tensor Categories. American Mathematical Society, 2015. Available online at <http://www.math.mit.edu/~etingof/egnobookfinal.pdf>.