

Semisimple 2-categories

Ying Hong Tham

UHH

2022

Goal

$$\{\text{multifusion category}\} \xrightarrow[\simeq]{\mathcal{M}\text{od}_{s.s.}^{fin}(-)} \{\text{finite semisimple 2-cat}\}$$

- ▶ The 2-category $\mathcal{M}\text{od}_{s.s.}^{fin}(C)$ of finite semisimple module categories over a multifusion category C is semisimple.
- ▶ For any finite semisimple 2-category \mathcal{C} , there exists a multifusion category C such that $\mathcal{C} \simeq \mathcal{M}\text{od}_{s.s.}^{fin}(C)$

Overview

- ▶ direct sum

Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff

Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff
- ▶ simple objects

Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff
- ▶ simple objects
- ▶ 'Schur lemma'

Overview

- ▶ direct sum
- ▶ idempotent completion, separable stuff
- ▶ simple objects
- ▶ 'Schur lemma'
- ▶ main results

Conventions

In relation to a 2-category:

- ▶ \mathcal{C} (caligraphic font): 2-category;
- ▶ X, Y, F (upper case latin): object of 2-category, functor between 2-categories;
- ▶ f, g (lower case latin): 1-morphism; we write $\mathcal{C}(X, Y)$ for the category of morphisms from X to Y ;
- ▶ $\eta, \varepsilon, \delta$ (lower case greek): 2-morphism; for a 2-morphism $\alpha : f \Rightarrow g : X \rightarrow Y$, we may write $\alpha \in \mathcal{C}(X, Y)(f, g)$ to indicate its sources and targets, or simply $\alpha \in \text{Hom}(f, g)$ if the objects are clear

Conventions

In relation to a 1-category:

- ▶ C, A (upper case latin): category;
- ▶ a, b, f, g (lower case latin): objects in category, functor between categories;
- ▶ α, β (lower case greek): morphism in category

Conventions

We also compose morphisms from right to left: in a 2-category \mathcal{C} , for $\alpha \in \mathcal{C}(X, Y)(f, f')$, $\beta \in \mathcal{C}(Y, Z)(g, g')$, $\gamma \in \mathcal{C}(X, Y)(f', f'')$, we write:

for composition of 1-morphisms,

$$g \circ f, g \circ f', \dots : X \rightarrow Z$$

for horizontal composition of 2-morphisms,

$$\beta \circ \alpha : (g \circ f) \Rightarrow (g' \circ f') : X \rightarrow Z$$

for vertical composition of 2-morphisms,

$$\gamma \cdot \alpha : f \Rightarrow f'' : X \rightarrow Y$$

Conventions

In general, if P is a property of a 1-category, we say that a 2-category \mathcal{C} is *locally P* if every hom-category $\mathcal{C}(X, Y)$ satisfies P .

Conventions

In general, if P is a property of a 1-category, we say that a 2-category \mathcal{C} is *locally P* if every hom-category $\mathcal{C}(X, Y)$ satisfies P . By 2-category we always mean a weak 2-category that is furthermore locally additive over \mathbf{k} . By 2-functor (sometimes just functor for simplicity) between 2-categories will always be locally \mathbf{k} -linear.

Review

Additive 2-category, direct sum of objects

Additive 2-category, direct sum of objects

Definition (direct sum of objects in 2-category)

A *direct sum* of two objects A_1, A_2 in \mathcal{C} is an object $A_1 \boxplus A_2$ together with inclusion and projection 1-morphisms

$i_k : A_k \Rightarrow A_1 \boxplus A_2 : p_k$, such that

- ▶ $p_k \circ i_k \simeq \text{id}_{A_k}$,
- ▶ $p_2 \circ i_1, p_1 \circ i_2$ are zero 1-morphisms,
- ▶ $\text{id}_{A_1 \boxplus A_2} \simeq i_1 \circ p_1 \oplus i_2 \circ p_2$

Additive 2-category, direct sum of objects

Definition (direct sum of objects in 2-category)

A *direct sum* of two objects A_1, A_2 in \mathcal{C} is an object $A_1 \boxplus A_2$ together with inclusion and projection 1-morphisms

$i_k : A_k \Rightarrow A_1 \boxplus A_2 : p_k$, such that

- ▶ $p_k \circ i_k \simeq \text{id}_{A_k}$,
- ▶ $p_2 \circ i_1, p_1 \circ i_2$ are zero 1-morphisms,
- ▶ $\text{id}_{A_1 \boxplus A_2} \simeq i_1 \circ p_1 \oplus i_2 \circ p_2$

Definition (zero object)

A *zero object* in \mathcal{C} is an object 0 with trivial endomorphism category $\mathcal{C}(0, 0)$ (has one object id_0 with only identity morphism id_{id_0}).

Additive 2-category, direct sum of objects

Definition (direct sum of objects in 2-category)

A *direct sum* of two objects A_1, A_2 in \mathcal{C} is an object $A_1 \boxplus A_2$ together with inclusion and projection 1-morphisms

$i_k : A_k \Rightarrow A_1 \boxplus A_2 : p_k$, such that

- ▶ $p_k \circ i_k \simeq \text{id}_{A_k}$,
- ▶ $p_2 \circ i_1, p_1 \circ i_2$ are zero 1-morphisms,
- ▶ $\text{id}_{A_1 \boxplus A_2} \simeq i_1 \circ p_1 \oplus i_2 \circ p_2$

Definition (zero object)

A *zero object* in \mathcal{C} is an object 0 with trivial endomorphism category $\mathcal{C}(0, 0)$ (has one object id_0 with only identity morphism id_{id_0}).

Definition

2-category \mathcal{C} is *additive* if finite direct sums of objects exist, has a zero object, (and is locally additive).

Additive 2-category, direct sum of objects

Definition (subobject)

A 1-morphism $i : X \rightarrow Y$ is *fully faithful* (or (X, i) is a *subobject* of Y) if it induces fully faithful functors between hom-categories by post-composition, i.e. for all objects A , $i \circ - : \mathcal{C}(A, X) \rightarrow \mathcal{C}(A, Y)$ is fully faithful.

Additive 2-category, direct sum of objects

Definition (subobject)

A 1-morphism $i : X \rightarrow Y$ is *fully faithful* (or (X, i) is a *subobject* of Y) if it induces fully faithful functors between hom-categories by post-composition, i.e. for all objects A , $i \circ - : \mathcal{C}(A, X) \rightarrow \mathcal{C}(A, Y)$ is fully faithful.

Proposition

$i_k : A_k \rightarrow A_1 \boxplus A_2$ is fully faithful.

Additive 2-category, direct sum of objects

Definition (Direct sum of 2-categories)

Given additive 2-categories \mathcal{C}_j , $j \in J$, we may consider the direct sum 2-category $\mathcal{C} := \boxplus_{j \in J} \mathcal{C}_j$:

- ▶ $\text{Obj } \mathcal{C} = \prod_{j \in J} \text{Obj } \mathcal{C}_j$

Additive 2-category, direct sum of objects

Definition (Direct sum of 2-categories)

Given additive 2-categories \mathcal{C}_j , $j \in J$, we may consider the direct sum 2-category $\mathcal{C} := \boxplus_{j \in J} \mathcal{C}_j$:

- ▶ $\text{Obj } \mathcal{C} = \prod_{j \in J} \text{Obj } \mathcal{C}_j$
- ▶ $X_i \in \mathcal{C}_i$ becomes $(\dots, 0, X_i, 0, \dots) \in \mathcal{C}$.

Additive 2-category, direct sum of objects

Definition (Direct sum of 2-categories)

Given additive 2-categories \mathcal{C}_j , $j \in J$, we may consider the direct sum 2-category $\mathcal{C} := \boxplus_{j \in J} \mathcal{C}_j$:

- ▶ $\text{Obj } \mathcal{C} = \prod_{j \in J} \text{Obj } \mathcal{C}_j$
- ▶ $X_i \in \mathcal{C}_i$ becomes $(\dots, 0, X_i, 0, \dots) \in \mathcal{C}$.
- ▶ $\mathcal{C}((X_i)_{i \in J}, (Y_j)_{j \in J}) = \bigoplus \mathcal{C}_i(X_i, Y_i)$

Idempotent completeness, separable monads, splittings

Idempotent completeness, separable monads, splittings

Definition

A *separable algebra* (a, μ, η) in a tensor category \mathcal{C} is an algebra that admits an a - a -bimodule section ${}_a a_a \rightarrow {}_a a \otimes a_a$ to μ .

Idempotent completeness, separable monads, splittings

Definition

A *separable algebra* (a, μ, η) in a tensor category \mathcal{C} is an algebra that admits an a - a -bimodule section ${}_a a_a \rightarrow {}_a a \otimes a_a$ to μ .

Definition

Let (t, μ, η) be a monad on an object X in a 2-category \mathcal{C} . We say t is *separable* if there is a t - t -bimodule section $t \Rightarrow t \circ t$ to μ .

Idempotent completeness, separable monads, splittings

Definition

A *separable algebra* (a, μ, η) in a tensor category \mathcal{C} is an algebra that admits an a - a -bimodule section ${}_a a_a \rightarrow {}_a a \otimes a_a$ to μ .

Definition

Let (t, μ, η) be a monad on an object X in a 2-category \mathcal{C} . We say t is *separable* if there is a t - t -bimodule section $t \Rightarrow t \circ t$ to μ .

In other words, a separable monad over X is a separable algebra in $\mathcal{C}(X, X)$.

Idempotent completeness, separable monads, splittings

Definition

Let $r \vdash l : X \rightarrow Y$ be an adjunction with unit $\eta : \text{id}_X \Rightarrow rl$ and counit $\varepsilon : lr \Rightarrow \text{id}_Y$. We say the adjunction $l \dashv r$ is *separable* if ε admits a section.

Idempotent completeness, separable monads, splittings

Definition

Let $r \vdash l : X \rightarrow Y$ be an adjunction with unit $\eta : \text{id}_X \Rightarrow rl$ and counit $\varepsilon : lr \Rightarrow \text{id}_Y$. We say the adjunction $l \dashv r$ is *separable* if ε admits a section.

Clearly, if an adjunction $l \dashv r$ is separable, then the monad rl is separable - the splitting gives $rl \simeq r(\text{id}_Y)l \Rightarrow r(lr)l \simeq (rl)(rl)$

Idempotent completeness, separable monads, splittings

Definition

Let $r \vdash l : X \rightarrow Y$ be an adjunction with unit $\eta : \text{id}_X \Rightarrow rl$ and counit $\varepsilon : lr \Rightarrow \text{id}_Y$. We say the adjunction $l \dashv r$ is *separable* if ε admits a section.

Clearly, if an adjunction $l \dashv r$ is separable, then the monad rl is separable - the splitting gives $rl \simeq r(\text{id}_Y)l \Rightarrow r(lr)l \simeq (rl)(rl)$

Definition

Let (t, μ, η) be a separable monad on an object $X \in \mathcal{C}$. A *(separable) splitting* of t is a (separable) adjunction $r \vdash l : X \rightarrow Y$ together with an isomorphism $\psi : rl \simeq t$ as monads on X .

Idempotent completeness, separable monads, splittings

Definition

Let $r \vdash l : X \rightarrow Y$ be an adjunction with unit $\eta : \text{id}_X \Rightarrow rl$ and counit $\varepsilon : lr \Rightarrow \text{id}_Y$. We say the adjunction $l \dashv r$ is *separable* if ε admits a section.

Clearly, if an adjunction $l \dashv r$ is separable, then the monad rl is separable - the splitting gives $rl \simeq r(\text{id}_Y)l \Rightarrow r(lr)l \simeq (rl)(rl)$

Definition

Let (t, μ, η) be a separable monad on an object $X \in \mathcal{C}$. A *(separable) splitting* of t is a (separable) adjunction $r \vdash l : X \rightarrow Y$ together with an isomorphism $\psi : rl \simeq t$ as monads on X .

Proposition (Uniqueness of splitting)

[[1], Theorem A.3.1] In locally idempotent complete 2-category \mathcal{C} , separable splitting of monad, if exists, is unique up to equivalence.

Idempotent completeness, 2-category

Definition

A 2-category \mathcal{C} is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

Idempotent completeness, 2-category

Definition

A 2-category \mathcal{C} is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

Definition (Idempotent completion)

Let \mathcal{C} be a locally idempotent complete 2-category. The *idempotent completion of \mathcal{C}* , denoted \mathcal{C}^∇ , with:

- ▶ Objects: (X, p) separable monad in \mathcal{C} ,
- ▶ $\mathcal{C}^\nabla((X, p), (Y, q)) = q\text{-bimod-}p(\mathcal{C}(X, Y))$

Idempotent completeness, 2-category

Definition

A 2-category \mathcal{C} is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

Definition (Idempotent completion)

Let \mathcal{C} be a locally idempotent complete 2-category. The *idempotent completion* of \mathcal{C} , denoted \mathcal{C}^∇ , with:

- ▶ Objects: (X, p) separable monad in \mathcal{C} ,
- ▶ $\mathcal{C}^\nabla((X, p), (Y, q)) = q\text{-bimod-}p(\mathcal{C}(X, Y))$

There is a natural 2-functor $I : \mathcal{C} \rightarrow \mathcal{C}^\nabla$ that is fully faithful.

Idempotent completeness, 2-category

Definition

A 2-category \mathcal{C} is *idempotent complete* if every separable monad admits a splitting and is locally idempotent complete.

Definition (Idempotent completion)

Let \mathcal{C} be a locally idempotent complete 2-category. The *idempotent completion* of \mathcal{C} , denoted \mathcal{C}^∇ , with:

- ▶ Objects: (X, p) separable monad in \mathcal{C} ,
- ▶ $\mathcal{C}^\nabla((X, p), (Y, q)) = q\text{-bimod-}p(\mathcal{C}(X, Y))$

There is a natural 2-functor $I : \mathcal{C} \rightarrow \mathcal{C}^\nabla$ that is fully faithful.

A 2-functor $F : \mathcal{C} \rightarrow \mathcal{D}$ extends to a 2-functor $F^\nabla : \mathcal{C}^\nabla \rightarrow \mathcal{D}^\nabla$ that commutes with I 's.

Idempotent completeness, 2-category

Key example:

\mathcal{BC} : one object $*$ with endomorphism category $\mathcal{BC}(*, *) = C$
(with $x \otimes y = x \circ y$)

$$(\mathcal{BC})^\nabla = \begin{cases} \text{Obj} : \text{separable algebras in } C \\ \text{Mor} : (\mathcal{BC})^\nabla(a, b) = b\text{-bimod-}a(C) \end{cases}$$

Idempotent completeness, 2-category

Proposition

\mathcal{C}^∇ is idempotent complete. Moreover, if \mathcal{C} is already idempotent complete, then $I : \mathcal{C} \simeq \mathcal{C}^\nabla$ is an equivalence.

Idempotent completeness, 2-category

Proposition

\mathcal{C}^∇ is idempotent complete. Moreover, if \mathcal{C} is already idempotent complete, then $I : \mathcal{C} \simeq \mathcal{C}^\nabla$ is an equivalence.

As a consequence, if \mathcal{D} is idempotent complete, then we have equivalences of 2-categories

$$\mathcal{F}un(\mathcal{C}, \mathcal{D}) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D}^\nabla) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D})$$

Idempotent completeness, 2-category

Proposition

\mathcal{C}^∇ is idempotent complete. Moreover, if \mathcal{C} is already idempotent complete, then $I : \mathcal{C} \simeq \mathcal{C}^\nabla$ is an equivalence.

As a consequence, if \mathcal{D} is idempotent complete, then we have equivalences of 2-categories

$$\mathcal{F}un(\mathcal{C}, \mathcal{D}) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D}^\nabla) \simeq \mathcal{F}un(\mathcal{C}^\nabla, \mathcal{D})$$

Proposition

If $F : \mathcal{C} \rightarrow \mathcal{D}$ is fully faithful, then $F^\nabla : \mathcal{C}^\nabla \rightarrow \mathcal{D}^\nabla$ is also fully faithful.

Idempotent completeness, 2-category

Proposition ([1]Prop 1.3.13)

For a multifusion category C , the following 2-functor is an equivalence:

$$(-)\text{-mod}(C) : (\mathcal{BC})^{\nabla} \rightarrow \mathcal{Mod}(C)$$

$$a \mapsto a\text{-mod}(C)$$

$${}_b m_a \mapsto m \otimes_a -$$

$$\varphi \mapsto \varphi \otimes_a -$$

Idempotent completeness, 2-category

Proof.

Essential surjectivity follows from:

-[2] Theorem 1: for a finite semisimple right module category over multifusion C , there exists a semisimple algebra a in C such that $M \simeq a\text{-mod}(C)$ as right C -module categories; and

Idempotent completeness, 2-category

Proof.

Essential surjectivity follows from:

- [2] Theorem 1: for a finite semisimple right module category over multifusion C , there exists a semisimple algebra a in C such that $M \simeq a\text{-mod}(C)$ as right C -module categories; and
- [4] Corollary 2.6.9: When C is multifusion over a field of characteristic 0, a right C -module category M is separable ($\simeq a\text{-mod}(C)$ for a separable algebra a) if and only if it is semisimple.

Idempotent completeness, 2-category

Proof.

Essential surjectivity follows from:

- [2]Theorem 1: for a finite semisimple right module category over multifusion C , there exists a semisimple algebra a in C such that $M \simeq a\text{-mod}(C)$ as right C -module categories; and
- [4]Corollary 2.6.9: When C is multifusion over a field of characteristic 0, a right C -module category M is separable ($\simeq a\text{-mod}(C)$ for a separable algebra a) if and only if it is semisimple.

Fully faithfulness follows from [5]Prop 7.11.1.



Idempotent completeness, 2-category

Proof.

Essential surjectivity follows from:

-[2]Theorem 1: for a finite semisimple right module category over multifusion C , there exists a semisimple algebra a in C such that $M \simeq a\text{-mod}(C)$ as right C -module categories; and

-[4]Corollary 2.6.9: When C is multifusion over a field of characteristic 0, a right C -module category M is separable ($\simeq a\text{-mod}(C)$ for a separable algebra a) if and only if it is semisimple.

Fully faithfulness follows from [5]Prop 7.11.1. □

This is almost one half of the main result; one still needs to prove local semisimplicity and existence of adjoints. This will follow from more results from [3],[4], which we show later.

Simple objects

Simple objects

Proposition (equivalent notions of simple-ness)

Let \mathcal{C} be an idempotent complete and locally finite semisimple 2-category, and let $X \in \mathcal{C}$ be a nonzero object. Then the following notions of X being simple are equivalent:

Simple objects

Proposition (equivalent notions of simple-ness)

Let \mathcal{C} be an idempotent complete and locally finite semisimple 2-category, and let $X \in \mathcal{C}$ be a nonzero object. Then the following notions of X being simple are equivalent:

(1) any subobject $i : Y \rightarrow X$ of X is either 0 or an equivalence;

Simple objects

Proposition (equivalent notions of simple-ness)

Let \mathcal{C} be an idempotent complete and locally finite semisimple 2-category, and let $X \in \mathcal{C}$ be a nonzero object. Then the following notions of X being simple are equivalent:

- (1) any subobject $i : Y \rightarrow X$ of X is either 0 or an equivalence;*
- (2) X cannot be written as a non-trivial direct sum, i.e. if $X = \boxplus X_i$, then $X_i \simeq 0$ for all but one i ;*

Simple objects

Proposition (equivalent notions of simple-ness)

Let \mathcal{C} be an idempotent complete and locally finite semisimple 2-category, and let $X \in \mathcal{C}$ be a nonzero object. Then the following notions of X being simple are equivalent:

- (1) any subobject $i : Y \rightarrow X$ of X is either 0 or an equivalence;*
- (2) X cannot be written as a non-trivial direct sum, i.e. if $X = \boxplus X_i$, then $X_i \simeq 0$ for all but one i ;*
- (3) id_X is a simple object in $\mathcal{C}(X, X)$.*

Simple objects

Proof sketch.

(1) \Rightarrow (2): Contravariant statement follows from fully faithfulness of $i_k : A_k \rightarrow A_1 \boxplus A_2$.

Simple objects

Proof sketch.

(1) \Rightarrow (2): Contravariant statement follows from fully faithfulness of $i_k : A_k \rightarrow A_1 \boxplus A_2$.

(2) \Rightarrow (3): Contravariant statement is “identity splitting implies object splitting”, uses idempotent completeness of \mathcal{C} to split out objects corresponding to summands of id_X (which are separable monads) (see [1]Prop 1.3.16).

Simple objects

Proof sketch.

(1) \Rightarrow (2): Contravariant statement follows from fully faithfulness of $i_k : A_k \rightarrow A_1 \boxplus A_2$.

(2) \Rightarrow (3): Contravariant statement is “identity splitting implies object splitting”, uses idempotent completeness of \mathcal{C} to split out objects corresponding to summands of id_X (which are separable monads) (see [1]Prop 1.3.16).

(3) \Rightarrow (1): for non-zero fully faithful $r : Y \rightarrow X$, with id_X simple, consider the left adjoint $l : X \rightarrow Y$, use fully faithfulness to get a preimage $\delta : \text{id}_Y \Rightarrow lr$ of $\eta \circ r : r \Rightarrow rlr$. Show δ is a section of the counit. Etc. (See [1]Prop 1.2.14) □

Semisimple 2-category

Definition ((finite) semisimple 2-category)

Semisimple 2-category

Definition ((finite) semisimple 2-category)

A 2-category \mathcal{C} is *semisimple* if it is:

- ▶ locally semisimple,
- ▶ admits left and right adjoints for every 1-morphism,
- ▶ additive,
- ▶ idempotent complete.

It is furthermore *finite semisimple* if it is also locally finite and has finitely many equivalence classes of simple objects.


New Stuff

Schur's lemma

The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case.


Schur's lemma

The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case. However, recall $\mathcal{M}\text{od}(\text{Vec}_{\mathbb{Z}/2})$:



Schur's lemma


The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case. However, recall $\mathcal{M}\text{od}(\text{Vec}_{\mathbb{Z}/2})$:



There can be nonzero morphisms between non-equivalent simple objects - weird!

Schur's lemma

The equivalence between notions of a simple object in a semisimple 2-category, is similar to the semisimple 1-category case. However, recall $\mathcal{M}\text{od}(\text{Vec}_{\mathbb{Z}/2})$:



There can be nonzero morphisms between non-equivalent simple objects - weird!

“2-Morita equivalences” between fusion categories

Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Proof.

$$A \xrightarrow{\quad f \quad} B \xrightarrow{\quad g \quad} C$$




Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Proof.




Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Proof.




Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Proof.



$$\text{id}_{\text{id}_B} = \varepsilon \cdot \delta$$




Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Proof.



$$\text{id}_{\text{id}_B} = \varepsilon \cdot \delta$$

$$\text{id}_g = (\text{id}_g \circ \varepsilon) \cdot (\text{id}_g \circ \delta) : g \Rightarrow gff^* \Rightarrow g$$




Schur's lemma

Proposition (Schur's Lemma, [1]Prop 1.2.19)

In a semisimple 2-category \mathcal{C} , if $f : A \rightarrow B, g : B \rightarrow C$ are nonzero 1-morphisms between simple objects A, B, C , then $g \circ f$ is also nonzero.

Proof.



$$\text{id}_{\text{id}_B} = \epsilon \cdot \delta$$

$$\text{id}_g = (\text{id}_g \circ \epsilon) \cdot (\text{id}_g \circ \delta) : g \Rightarrow gff^* \Rightarrow g$$

$$\text{if } gf \cong 0, \text{ then } \text{id}_g = 0, \text{ and } g \cong 0$$



Schur's lemma - components

Definition (component of semisimple 2-category)

In a semisimple 2-category \mathcal{C} , we say two simple objects A, B belong to the same *component* if there exists a nonzero 1-morphism $f : A \rightarrow B$.

Schur's lemma - components

Definition (component of semisimple 2-category)

In a semisimple 2-category \mathcal{C} , we say two simple objects A, B belong to the same *component* if there exists a nonzero 1-morphism $f : A \rightarrow B$.

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

Schur's lemma - components

Definition (component of semisimple 2-category)

In a semisimple 2-category \mathcal{C} , we say two simple objects A, B belong to the same *component* if there exists a nonzero 1-morphism $f : A \rightarrow B$.

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

In a finite semisimple 2-category \mathcal{C} , there will be finite set J of components;

Schur's lemma - components

Definition (component of semisimple 2-category)

In a semisimple 2-category \mathcal{C} , we say two simple objects A, B belong to the same *component* if there exists a nonzero 1-morphism $f : A \rightarrow B$.

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

In a finite semisimple 2-category \mathcal{C} , there will be finite set J of components;

full subcategory \mathcal{C}_j , $j \in J$: direct sum of simple objects in j

Schur's lemma - components

Definition (component of semisimple 2-category)

In a semisimple 2-category \mathcal{C} , we say two simple objects A, B belong to the same *component* if there exists a nonzero 1-morphism $f : A \rightarrow B$.

-transitivity from Schur's lemma, symmetry from adjoints, reflexivity from identity

In a finite semisimple 2-category \mathcal{C} , there will be finite set J of components;

full subcategory \mathcal{C}_j , $j \in J$: direct sum of simple objects in j gives direct sum decomposition $\mathcal{C} \simeq \bigoplus_{j \in J} \mathcal{C}_j$.

Main results

Theorem ([1]Theorem 1.4.8)

The 2-category of finite semisimple module categories of a multifusion category \mathcal{C} is a finite semisimple 2-category.

Main results

Theorem ([1]Theorem 1.4.8)

The 2-category of finite semisimple module categories of a multifusion category \mathcal{C} is a finite semisimple 2-category.

Proof.

$\text{Mod}(\mathcal{C}) \simeq (\mathcal{B}\mathcal{C})^\nabla$ is idempotent complete. $\text{Mod}(\mathcal{C})$ is clearly already additive.

Locally semisimple-ness follows directly from [4]Corollary 2.5.6, and existence of adjoints for 1-morphisms follows from [3]Corollary 2.13.



Main results

Theorem ([1]Theorem 1.4.9)

Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.

Proof.

Assume that \mathcal{C} has only one component.

Main results

Theorem ([1]Theorem 1.4.9)

Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.

Proof.

Assume that \mathcal{C} has only one component.

($\mathcal{C} \simeq \boxplus_{j \in J} \mathcal{C}_j$ and $\mathcal{C}_j \simeq \text{Mod}(C_j)$ for multifusion C_j , then
 $\mathcal{C} \simeq \text{Mod}(\oplus_{j \in J} C_j)$.)

Main results

Theorem ([1]Theorem 1.4.9)

Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.

Proof.

Assume that \mathcal{C} has only one component.

($\mathcal{C} \simeq \boxplus_{j \in J} \mathcal{C}_j$ and $\mathcal{C}_j \simeq \text{Mod}(C_j)$ for multifusion C_j , then
 $\mathcal{C} \simeq \text{Mod}(\oplus_{j \in J} C_j)$.)

Fix simple object X , let $C = \mathcal{C}(X, X)$.

Main results

Theorem ([1]Theorem 1.4.9)

Every finite semisimple 2-category is equivalent to the 2-category of finite semisimple module categories of a multifusion category.

Proof.

Assume that \mathcal{C} has only one component.

($\mathcal{C} \simeq \boxplus_{j \in J} \mathcal{C}_j$ and $\mathcal{C}_j \simeq \text{Mod}(C_j)$ for multifusion C_j , then
 $\mathcal{C} \simeq \text{Mod}(\oplus_{j \in J} C_j)$.)

Fix simple object X , let $C = \mathcal{C}(X, X)$.

We show $\mathcal{C} \simeq (\mathcal{B}C)^\nabla$.

Main results

Proof (Cont.)

Consider the inclusion 2-functor

$$F : \mathcal{BC} \rightarrow \mathcal{C}$$

$$* \mapsto X$$

which is fully faithful by construction.

Main results

Proof (Cont.)

Consider the inclusion 2-functor

$$F : \mathcal{BC} \rightarrow \mathcal{C}$$

$$* \mapsto X$$

which is fully faithful by construction. Then

$$F^\nabla : (\mathcal{BC})^\nabla \rightarrow \mathcal{C}$$

is fully faithful.

Main results

Proof (Cont.)

Consider the inclusion 2-functor

$$\begin{aligned} F : \mathcal{BC} &\rightarrow \mathcal{C} \\ * &\mapsto X \end{aligned}$$

which is fully faithful by construction. Then

$$F^\nabla : (\mathcal{BC})^\nabla \rightarrow \mathcal{C}$$

is fully faithful. Remains to show other simples are in essential image.

Main results

Proof (Cont.)


Y

$$* \xrightarrow{F^\nabla} X$$




Main results

Proof (Cont.)




Main results

Proof (Cont.)



Main results

Proof (Cont.)




id_Y simple
 ε has section
 \Rightarrow sep. adj.



Main results

Proof (Cont.)




id_Y simple
 ε has section
 \Rightarrow sep. adj.



Main results

Proof (Cont.)




id_Y simple
 ε has section
 \Rightarrow sep. adj.



Main results

Proof (Cont.)



id_Y simple
 ε has section
 \Rightarrow sep. adj.



Main results

Proof (Cont.)

$$\begin{array}{ccccc} (*, gf) & \xrightarrow{F^\nabla} & F^\nabla(*, gf) & \xrightarrow{\cong} & Y \\ & \swarrow & \swarrow & \nearrow f & \nearrow g \\ & & * & \xrightarrow{F^\nabla} & X \end{array}$$

Diagram illustrating the proof structure. The top row shows a sequence of maps: $(*, gf) \xrightarrow{F^\nabla} F^\nabla(*, gf) \xrightarrow{\cong} Y$. The bottom row shows $* \xrightarrow{F^\nabla} X$. A curved arrow labeled gf points from $*$ to $(*, gf)$. A curved arrow labeled f points from X to $F^\nabla(*, gf)$. A curved arrow labeled g points from X to Y . A curved arrow labeled \dashv points from Y to X .

id_Y simple
 ε has section
 \Rightarrow sep. adj.



Main results

The only property of X that we used is the fact that there exists a nonzero 1-morphism from X to every simple in \mathcal{C} , and thus any nonzero X will do.

Main results

The only property of X that we used is the fact that there exists a nonzero 1-morphism from X to every simple in \mathcal{C} , and thus any nonzero X will do. Taking, say, $X = \boxplus X_i$, where the sum is over equivalence classes of simples, would result in a multifusion $\mathcal{C} = \mathcal{C}(X, X)$.



Main results

The only property of X that we used is the fact that there exists a nonzero 1-morphism from X to every simple in \mathcal{C} , and thus any nonzero X will do. Taking, say, $X = \boxplus X_i$, where the sum is over equivalence classes of simples, would result in a multifusion $\mathcal{C} = \mathcal{C}(X, X)$.


We can also avoid the first step of taking only one component of \mathcal{C} , take X with at least one object from each \mathcal{C}_j in its direct sum decomposition.

Main results

Every object contains within them the data to reconstruct the whole 2-category.



Indra's Net of Pearls (or $T^3 \# T^3$)



Example

$C = \text{Vec}_G$ for a finite group G .

Example

$C = \text{Vec}_G$ for a finite group G .

e : trivial algebra \mathbf{k}_e $e\text{-mod}(C) \simeq C_C$

Example

$C = \text{Vec}_G$ for a finite group G .

e : trivial algebra \mathbf{k}_e $e\text{-mod}(C) \simeq C_C$

f : the group algebra $\mathbf{k}[G] = \bigoplus \mathbf{k}_g$ $f\text{-mod}(C) \simeq \text{Vec}$
(right action on Vec = forget G -grading)

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

For $f\text{-mod}(C)$, an endofunctor is given by some $m \in f\text{-bimod-}f(C)$; write $m = \bigoplus m_g$.

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

For $f\text{-mod}(C)$, an endofunctor is given by some $m \in f\text{-bimod-}f(C)$; write $m = \bigoplus m_g$. The right f -action on m makes all the m_g isomorphic in a coherent manner.

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

For $f\text{-mod}(C)$, an endofunctor is given by some $m \in f\text{-bimod-}f(C)$; write $m = \bigoplus m_g$. The right f -action on m makes all the m_g isomorphic in a coherent manner. For $h \in G$, conjugation (left action by h and right action by h^{-1}) gives an action of G on m_e ; determines m completely.

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

For $f\text{-mod}(C)$, an endofunctor is given by some $m \in f\text{-bimod-}f(C)$; write $m = \bigoplus m_g$. The right f -action on m makes all the m_g isomorphic in a coherent manner. For $h \in G$, conjugation (left action by h and right action by h^{-1}) gives an action of G on m_e ; determines m completely.

Thus, $\mathcal{M}\text{od}(C)(f, f) \simeq f\text{-bimod-}f(C) \simeq \text{Rep}(G)$.

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

For $f\text{-mod}(C)$, an endofunctor is given by some $m \in f\text{-bimod-}f(C)$; write $m = \bigoplus m_g$. The right f -action on m makes all the m_g isomorphic in a coherent manner. For $h \in G$, conjugation (left action by h and right action by h^{-1}) gives an action of G on m_e ; determines m completely.

Thus, $\text{Mod}(C)(f, f) \simeq f\text{-bimod-}f(C) \simeq \text{Rep}(G)$.

(Alternatively, one can also check that C -module structure on $m : \text{Vec} \rightarrow \text{Vec}$ amounts to G -action on $m(\mathbf{k}).$)

Example

Next we study the functor categories. Clearly the C -endofunctors of C_C is C itself.

For $f\text{-mod}(C)$, an endofunctor is given by some $m \in f\text{-bimod-}f(C)$; write $m = \bigoplus m_g$. The right f -action on m makes all the m_g isomorphic in a coherent manner. For $h \in G$, conjugation (left action by h and right action by h^{-1}) gives an action of G on m_e ; determines m completely.

Thus, $\text{Mod}(C)(f, f) \simeq f\text{-bimod-}f(C) \simeq \text{Rep}(G)$.

(Alternatively, one can also check that C -module structure on $m: \text{Vec} \rightarrow \text{Vec}$ amounts to G -action on $m(\mathbf{k})$.)

Thus we have:

$$\text{Mod}(\text{Vec}_G) \simeq \text{Mod}(\text{Rep}(G))$$

Example

For functor categories between them, we see that

$$\mathcal{M}\mathrm{od}(C)(e, f) \simeq f\text{-bimod-}e(C) \simeq f\text{-mod}(C) \simeq \mathbf{Vec}$$

$$\mathcal{M}\mathrm{od}(C)(f, e) \simeq e\text{-bimod-}f(C) \simeq \mathrm{mod-}f(C) \simeq \mathbf{Vec}$$

Example

For functor categories between them, we see that

$$\mathcal{M}\mathrm{od}(C)(e, f) \simeq f\text{-bimod-}e(C) \simeq f\text{-mod}(C) \simeq \mathrm{Vec}$$

$$\mathcal{M}\mathrm{od}(C)(f, e) \simeq e\text{-bimod-}f(C) \simeq \mathrm{mod-}f(C) \simeq \mathrm{Vec}$$

Note for $G = \mathbb{Z}/2$, $\mathrm{Vec}_{\mathbb{Z}/2} \simeq \mathrm{Rep}(\mathbb{Z}/2)$, don't see difference in endomorphism categories.

Example

For functor categories between them, we see that






$$\mathcal{M}\mathrm{od}(C)(e, f) \simeq f\text{-bimod-}e(C) \simeq f\text{-mod}(C) \simeq \mathrm{Vec}$$

$$\mathcal{M}\mathrm{od}(C)(f, e) \simeq e\text{-bimod-}f(C) \simeq \mathrm{mod-}f(C) \simeq \mathrm{Vec}$$

Note for $G = \mathbb{Z}/2$, $\mathrm{Vec}_{\mathbb{Z}/2} \simeq \mathrm{Rep}(\mathbb{Z}/2)$, don't see difference in endomorphism categories.

Of course, there are many other objects in $\mathcal{M}\mathrm{od}(C)$, e.g. for a subgroup $H \subseteq G$, have group algebra $\mathbf{k}[H]$.

End!

-  Douglas, Christopher L., and David J. Reutter. “Fusion 2-categories and a state-sum invariant for 4-manifolds.” arXiv preprint arXiv:1812.11933 (2018).
-  Ostrik, Victor. “Module categories, weak Hopf algebras and modular invariants.” Transformation groups 8, no. 2 (2003): 177-206.
-  C. L. Douglas, C. Schommer-Pries, and N. Snyder. The balanced tensor product of module categories. Kyoto J. Math., 2017. arXiv:1406.4204.
-  C. L. Douglas, C. Schommer-Pries, and N. Snyder. Dualizable tensor categories. Mem. Amer. Math. Soc., 2017. arXiv:1312.7188.
-  P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. Tensor Categories. American Mathematical Society, 2015. Available online at <http://www.math.mit.edu/~etingof/egnobookfinal.pdf>.