Simple Auctions and Approximations

Yingkai Li

EC4501/EC4501HM

Mechanism Design

A mechanism design instance is denoted as $\Gamma_M = \left(N, \Omega, (v_i)_{i \in N}, (\Theta_i)_{i \in N}, F\right)$ where

- ullet N is the set of agents;
- Ω is the set of outcomes;
- Θ_i is the set of agent i's "types" where $\theta_i \in \Theta_i$ is private information of i;
- $v_i: \Omega \times \Theta_i \to \mathbb{R}$ is agent *i*'s value function;
- $F = F_1 \times \cdots \times F_n$ prior distribution over types.

Mechanism Design

A mechanism design instance is denoted as $\Gamma_{M}=\left(N,\Omega,\left(v_{i}\right)_{i\in N},\left(\Theta_{i}\right)_{i\in N},F\right)$ where

- ullet N is the set of agents;
- Ω is the set of outcomes;
- Θ_i is the set of agent i's "types" where $\theta_i \in \Theta_i$ is private information of i;
- $v_i: \Omega \times \Theta_i \to \mathbb{R}$ is agent *i*'s value function;
- $F = F_1 \times \cdots \times F_n$ prior distribution over types.

Let B_i be the report space of agent i.

A mechanism M = (x, p):

- $x: B \to \Delta(\Omega)$;
- $p_i: B \to \mathbb{R}, \forall i$.

Single-item Auctions

Auctions: a single item, n agents.

- each agent i has a private value $v_i \sim F_i \in \Delta(\mathbb{R}_+)$;
- each agent i has linear utility $u_i = v_i x_i p_i$ where $x_i \in [0, 1], p_i \in \mathbb{R}$;
- feasibility: $\sum_i x_i \leq 1$.

Single-item Auctions

Auctions: a single item, n agents.

- each agent i has a private value $v_i \sim F_i \in \Delta(\mathbb{R}_+)$;
- each agent i has linear utility $u_i = v_i x_i p_i$ where $x_i \in [0,1], p_i \in \mathbb{R}$;
- feasibility: $\sum_i x_i \leq 1$.

Welfare: $\mathbf{E}[\sum_i v_i x_i]$.

Revenue: $E[\sum_i p_i]$.

Welfare Approximations

Inefficiency of Standard Auctions

First-price auctions / all-pay auctions are not efficient in general.

ullet e.g., two agents, $v_1 \sim U[0,1], v_2 \sim U[0,2].$

Inefficiency of Standard Auctions

First-price auctions / all-pay auctions are not efficient in general.

ullet e.g., two agents, $v_1 \sim U[0,1], v_2 \sim U[0,2].$

Even though those auctions are inefficient, they are approximately efficient.

Inefficiency of Standard Auctions

First-price auctions / all-pay auctions are not efficient in general.

ullet e.g., two agents, $v_1 \sim U[0,1], v_2 \sim U[0,2].$

Even though those auctions are inefficient, they are approximately efficient.

Price of Anarchy (PoA): given any auction M, letting M(F) be the equilibrium welfare for auction M,

$$\operatorname{PoA}(M) = \max_{F} \frac{\operatorname{Wel}(F)}{M(F)}.$$

Smooth Auctions

Let $u_i(\mathbf{b}; \theta_i) = v_i(x(\mathbf{b}), \theta_i) - p_i(\mathbf{b})$ be the utility of agent i given bid profile b. Let $\mathcal{R}(\mathbf{b}) = \sum_i p_i(\mathbf{b})$.

Smooth Auctions

Let $u_i(\mathbf{b}; \theta_i) = v_i(x(\mathbf{b}), \theta_i) - p_i(\mathbf{b})$ be the utility of agent i given bid profile b. Let $\mathcal{R}(\mathbf{b}) = \sum_i p_i(\mathbf{b})$.

Definition (Smooth Auctions)

For parameters $\lambda \geq 0$ and $\mu \geq 1$, an auction is (λ, μ) -smooth if for every valuation profile $\mathbf{v} \in \mathcal{V}$ there exist bidding distributions $D_1^*(\mathbf{v}), \dots, D_n^*(\mathbf{v})$ such that, for every bid profile \mathbf{b} ,

$$\sum_{i} \mathbb{E}_{b_{i}^{*} \sim D_{i}^{*}(\mathbf{v})}[u_{i}(b_{i}^{*}, \mathbf{b}_{-i}; \mathbf{v}_{i})] \geq \lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}).$$

Smooth Auctions

Let $u_i(\mathbf{b}; \theta_i) = v_i(x(\mathbf{b}), \theta_i) - p_i(\mathbf{b})$ be the utility of agent i given bid profile b. Let $\mathcal{R}(\mathbf{b}) = \sum_i p_i(\mathbf{b})$.

Definition (Smooth Auctions)

For parameters $\lambda \geq 0$ and $\mu \geq 1$, an auction is (λ, μ) -smooth if for every valuation profile $\mathbf{v} \in \mathcal{V}$ there exist bidding distributions $D_1^*(\mathbf{v}), \ldots, D_n^*(\mathbf{v})$ such that, for every bid profile \mathbf{b} ,

$$\sum_{i} \mathbb{E}_{b_{i}^{*} \sim D_{i}^{*}(\mathbf{v})}[u_{i}(b_{i}^{*}, \mathbf{b}_{-i}; \mathbf{v}_{i})] \geq \lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}).$$

First-price auction is $(\frac{1}{2}, 1)$ -smooth.

ullet by bidding $\frac{v_i}{2}$, either wins and the utility is high, or loses and the total payment is high.

Smoothness of First-Price Auction

For each agent i, one possible strategy is to bid $b_i^* = \frac{v_i}{2}$ regardless of the opponents' strategy.

$$u_i(b_i^*, \mathbf{b}_{-i}; v_i) \ge \frac{1}{2}v_i - \mathcal{R}(\mathbf{b}).$$

since the bidder either wins and obtains utility $v_i - b_i^* = v_i - \frac{1}{2}v_i = \frac{1}{2}v_i \geq \frac{1}{2}v_i - \mathcal{R}(\mathbf{b})$, or loses and obtains utility $0 \geq \frac{1}{2}v_i - \mathcal{R}(\mathbf{b})$.

Smoothness of First-Price Auction

For each agent i, one possible strategy is to bid $b_i^* = \frac{v_i}{2}$ regardless of the opponents' strategy.

$$u_i(b_i^*, \mathbf{b}_{-i}; v_i) \ge \frac{1}{2}v_i - \mathcal{R}(\mathbf{b}).$$

since the bidder either wins and obtains utility $v_i - b_i^* = v_i - \frac{1}{2}v_i = \frac{1}{2}v_i \geq \frac{1}{2}v_i - \mathcal{R}(\mathbf{b})$, or loses and obtains utility $0 \geq \frac{1}{2}v_i - \mathcal{R}(\mathbf{b})$.

Let x_i^* be the welfare optimal allocation. Since the bid $b_i^*=rac{v_i}{2}$ guarantees non-negative utility,

$$u_i(b_i^*, \mathbf{b}_{-i}; v_i) \ge \left(\frac{1}{2}v_i - \mathcal{R}(\mathbf{b})\right) \cdot x_i^*(\mathbf{v}).$$

Smoothness of First-Price Auction

For each agent i, one possible strategy is to bid $b_i^* = \frac{v_i}{2}$ regardless of the opponents' strategy.

$$u_i(b_i^*, \mathbf{b}_{-i}; v_i) \ge \frac{1}{2}v_i - \mathcal{R}(\mathbf{b}).$$

since the bidder either wins and obtains utility $v_i - b_i^* = v_i - \frac{1}{2}v_i = \frac{1}{2}v_i \geq \frac{1}{2}v_i - \mathcal{R}(\mathbf{b})$, or loses and obtains utility $0 \geq \frac{1}{2}v_i - \mathcal{R}(\mathbf{b})$.

Let x_i^* be the welfare optimal allocation. Since the bid $b_i^* = \frac{v_i}{2}$ guarantees non-negative utility,

$$u_i(b_i^*, \mathbf{b}_{-i}; v_i) \ge \left(\frac{1}{2}v_i - \mathcal{R}(\mathbf{b})\right) \cdot x_i^*(\mathbf{v}).$$

Summing this inequality over all bidders i, we obtain

$$\sum_{i=1}^{n} u_i(b_i^*, \mathbf{b}_{-i}; v_i) \ge \sum_{i=1}^{n} \left(\frac{1}{2}v_i - \mathcal{R}(\mathbf{b})\right) \cdot x_i^*(\mathbf{v}) = \frac{1}{2} \text{Wel}(\mathbf{v}) - \mathcal{R}(\mathbf{b}),$$

for every valuation profile v and bid profile b.

Theorem

For any $\lambda \leq 1, \mu \geq 1$, if an auction M is (λ, μ) -smooth, then for every product distribution F, every Bayes-Nash equilibrium of the auction has expected welfare at least $\frac{\lambda}{\mu} \cdot \operatorname{Wel}(F)$.

Theorem

For any $\lambda \leq 1, \mu \geq 1$, if an auction M is (λ, μ) -smooth, then for every product distribution F, every Bayes-Nash equilibrium of the auction has expected welfare at least $\frac{\lambda}{\mu} \cdot \operatorname{Wel}(F)$.

Intuition: utility in BNE \geq utility given bidding strategy $D^* \geq \mathbf{E}[\lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(b)].$

Theorem

For any $\lambda \leq 1, \mu \geq 1$, if an auction M is (λ, μ) -smooth, then for every product distribution F, every Bayes-Nash equilibrium of the auction has expected welfare at least $\frac{\lambda}{\mu} \cdot \operatorname{Wel}(F)$.

Intuition: utility in BNE \geq utility given bidding strategy $D^* \geq \mathbf{E}[\lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(b)].$

welfare in BNE = utility in BNE + revenue $\geq \frac{\lambda}{\mu} \cdot \text{Wel}(F)$.

Theorem

For any $\lambda \leq 1, \mu \geq 1$, if an auction M is (λ, μ) -smooth, then for every product distribution F, every Bayes-Nash equilibrium of the auction has expected welfare at least $\frac{\lambda}{\mu} \cdot \operatorname{Wel}(F)$.

Intuition: utility in BNE \geq utility given bidding strategy $D^* \geq \mathbf{E}[\lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(b)].$

welfare in BNE = utility in BNE + revenue $\geq \frac{\lambda}{\mu} \cdot \operatorname{Wel}(F)$.

Corollary

The price of anarchy for first-price auction is at most 2.

- The PoA for first-price auction can be improved to $\frac{e^2}{e^2-1} \approx 1.16$ [Jin and Lu '22].
- ullet Techniques can be applied to other auction formats: all-pay auction is $(\frac{1}{2},1)$ -smooth.

Proof of the Smooth Auction Theorem (1/2)

Extension Theorem: We construct a valid randomized deviation $D'_i(v_i)$.

- Agent i samples fictitious valuations $\mathbf{v}'_{-i} \sim F_{-i}$.
- ② Agent i plays a bid b_i^* drawn from $D_i^*(v_i, \mathbf{v}'_{-i})$.

Proof of the Smooth Auction Theorem (1/2)

Extension Theorem: We construct a valid randomized deviation $D'_i(v_i)$.

- **1** Agent i samples fictitious valuations $\mathbf{v}'_{-i} \sim F_{-i}$.
- 2 Agent i plays a bid b_i^* drawn from $D_i^*(v_i, \mathbf{v}'_{-i})$.

Let s be a BNE strategy profile, and let \mathcal{G} be the resulting distribution of bids when $\mathbf{v} \sim F$. By the BNE condition, the expected utility of $s_i(v_i)$ is at least that of the deviation $D_i'(v_i)$.

$$\mathbb{E}_{BNE}[u_i|v_i] \geq \mathbb{E}_{b_i^* \sim D_i'(v_i), \mathbf{b}_{-i} \sim \mathcal{G}_{-i}}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)]$$

$$= \mathbb{E}_{\mathbf{v}'_{-i} \sim F_{-i}} \left[\mathbb{E}_{b_i^* \sim D_i^*(v_i, \mathbf{v}'_{-i}), \mathbf{b}_{-i} \sim \mathcal{G}_{-i}}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)] \right].$$

We take expectation over v_i and sum over i. Let U^{BNE} be the total expected utility.

$$U^{BNE} \ge \sum_{i} \mathbb{E}_{v_i, \mathbf{v}'_{-i}} \left[\mathbb{E}_{b_i^* \sim D_i^*(v_i, \mathbf{v}'_{-i}), \mathbf{b}_{-i} \sim \mathcal{G}_{-i}} [u_i(b_i^*, \mathbf{b}_{-i}; v_i)] \right].$$

Proof of the Smooth Auction Theorem (1/2)

Extension Theorem: We construct a valid randomized deviation $D'_i(v_i)$.

- **1** Agent i samples fictitious valuations $\mathbf{v}'_{-i} \sim F_{-i}$.
- ② Agent i plays a bid b_i^* drawn from $D_i^*(v_i, \mathbf{v}'_{-i})$.

Let s be a BNE strategy profile, and let \mathcal{G} be the resulting distribution of bids when $\mathbf{v} \sim F$. By the BNE condition, the expected utility of $s_i(v_i)$ is at least that of the deviation $D_i'(v_i)$.

$$\mathbb{E}_{BNE}[u_i|v_i] \ge \mathbb{E}_{b_i^* \sim D_i'(v_i), \mathbf{b}_{-i} \sim \mathcal{G}_{-i}}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)]$$

$$= \mathbb{E}_{\mathbf{v}'_{-i} \sim F_{-i}}\left[\mathbb{E}_{b_i^* \sim D_i^*(v_i, \mathbf{v}'_{-i}), \mathbf{b}_{-i} \sim \mathcal{G}_{-i}}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)]\right].$$

We take expectation over v_i and sum over i. Let U^{BNE} be the total expected utility.

$$U^{BNE} \ge \sum_{i} \mathbb{E}_{v_i, \mathbf{v}'_{-i}} \left[\mathbb{E}_{b_i^* \sim D_i^*(v_i, \mathbf{v}'_{-i}), \mathbf{b}_{-i} \sim \mathcal{G}_{-i}} [u_i(b_i^*, \mathbf{b}_{-i}; v_i)] \right].$$

Let $\mathbf{v} = (v_i, \mathbf{v}'_{-i})$. We rewrite the inequality using linearity of expectation:

$$U^{BNE} \ge \mathbb{E}_{\mathbf{v} \sim F, \mathbf{b} \sim \mathcal{G}} \left[\sum_{i} \mathbb{E}_{b_{i}^{*} \sim D_{i}^{*}(\mathbf{v})} [u_{i}(b_{i}^{*}, \mathbf{b}_{-i}; v_{i})] \right].$$

Crucially, v (from sampling) and b (from equilibrium) are independent here.

Proof of the Smooth Auction Theorem (2/2)

We apply the smoothness definition inside the expectation:

$$\sum_{i} \mathbb{E}_{b_i^*}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)] \ge \lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}).$$

Proof of the Smooth Auction Theorem (2/2)

We apply the smoothness definition inside the expectation:

$$\sum_{i} \mathbb{E}_{b_i^*}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)] \ge \lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}).$$

Substituting this bound:

$$\begin{split} U^{BNE} &\geq \mathbb{E}_{\mathbf{v} \sim F, \mathbf{b} \sim \mathcal{G}} \left[\lambda \mathrm{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}) \right] \\ &= \lambda \mathbb{E}_{\mathbf{v}} [\mathrm{Wel}(\mathbf{v})] - \mu \mathbb{E}_{\mathbf{b}} [\mathcal{R}(\mathbf{b})] \\ &= \lambda \mathrm{Wel}(F) - \mu R^{BNE}. \end{split} \tag{By independence}$$

Proof of the Smooth Auction Theorem (2/2)

We apply the smoothness definition inside the expectation:

$$\sum_{i} \mathbb{E}_{b_i^*}[u_i(b_i^*, \mathbf{b}_{-i}; v_i)] \ge \lambda \text{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}).$$

Substituting this bound:

$$\begin{split} U^{BNE} &\geq \mathbb{E}_{\mathbf{v} \sim F, \mathbf{b} \sim \mathcal{G}} \left[\lambda \mathrm{Wel}(\mathbf{v}) - \mu \mathcal{R}(\mathbf{b}) \right] \\ &= \lambda \mathbb{E}_{\mathbf{v}} [\mathrm{Wel}(\mathbf{v})] - \mu \mathbb{E}_{\mathbf{b}} [\mathcal{R}(\mathbf{b})] \\ &= \lambda \mathrm{Wel}(F) - \mu R^{BNE}. \end{split}$$
 (By independence)

Since $M(F)=U^{BNE}+R^{BNE}$, we have $M(F)+(\mu-1)R^{BNE}\geq \lambda \mathrm{Wel}(F)$. Since $\mu\geq 1$ and $M(F)\geq R^{BNE}$:

$$\mu \cdot M(F) \ge M(F) + (\mu - 1)R^{BNE} \ge \lambda \text{Wel}(F).$$

In practice, many auctions run in parallel.

• E.g., different sellers auction their products in parallel using first-price auctions.

In practice, many auctions run in parallel.

• E.g., different sellers auction their products in parallel using first-price auctions.

Is the equilibrium outcome approximately efficient in aggregation?

In practice, many auctions run in parallel.

• E.g., different sellers auction their products in parallel using first-price auctions.

Is the equilibrium outcome approximately efficient in aggregation?

Definition

A utility function is complement-free if there exists m additive valuations f_1,\ldots,f_m such that for any set S, $f(S) = \max_{k \le m} f_k(S).$

Theorem (Composition Theorem)

If players have complement-free utility functions, then the simultaneous composition of (λ, μ) -smooth auctions is again a (λ, μ) -smooth auction.

In practice, many auctions run in parallel.

• E.g., different sellers auction their products in parallel using first-price auctions.

Is the equilibrium outcome approximately efficient in aggregation?

Definition

A utility function is complement-free if there exists m additive valuations f_1, \ldots, f_m such that for any set S, $f(S) = \max_{k \le m} f_k(S).$

Theorem (Composition Theorem)

If players have complement-free utility functions, then the simultaneous composition of (λ,μ) -smooth auctions is again a (λ,μ) -smooth auction.

Corollary: PoA of the simultaneous composition of (λ, μ) -smooth auctions is at most $\frac{\mu}{\lambda}$.

Theorem (Composition Theorem)

If players have submodular utility functions, then the simultaneous composition of (λ,μ) -smooth auctions is again a (λ,μ) -smooth auction.

Theorem (Composition Theorem)

If players have submodular utility functions, then the simultaneous composition of (λ,μ) -smooth auctions is again a (λ,μ) -smooth auction.

Illustration for unit-demand auction and simultaneous first-price auction.

- given valuation profile v, find optimal allocation x(v);
- consider strategy profile where each agent i only bids $\frac{v_{ij}}{2}$ in auction j where $x_{ij}(v)=1$.

Theorem (Composition Theorem)

If players have submodular utility functions, then the simultaneous composition of (λ,μ) -smooth auctions is again a (λ,μ) -smooth auction.

Illustration for unit-demand auction and simultaneous first-price auction.

- given valuation profile v, find optimal allocation x(v);
- consider strategy profile where each agent i only bids $\frac{v_{ij}}{2}$ in auction j where $x_{ij}(v)=1$.

Reference: Roughgarden, T., Syrgkanis, V., & Tardos, E. (2017). The price of anarchy in auctions. Journal of Artificial Intelligence Research, 59, 59-101.

Reduction from Algorithms to Mechanisms

VCG Mechanisms

VCG mechanism: mechanism that implements efficient allocation in general environments.

allocation: chooses outcome

$$\omega^* = \underset{\omega \in \Omega}{\operatorname{argmax}} \sum_i v_i(\omega, \theta_i).$$

• payment: each agent i pays his externality on the welfare

$$p_i(\theta) = \max_{\omega \in \Omega} \sum_{j \neq i} v_j(\omega, \theta_j) - \sum_{j \neq i} v_j(\omega^*, \theta_j) \ge 0.$$

VCG Mechanisms

VCG mechanism: mechanism that implements efficient allocation in general environments.

allocation: chooses outcome

$$\omega^* = \operatorname*{argmax}_{\omega \in \Omega} \sum_{i} v_i(\omega, \theta_i).$$

• payment: each agent i pays his externality on the welfare

$$p_i(\theta) = \max_{\omega \in \Omega} \sum_{j \neq i} v_j(\omega, \theta_j) - \sum_{j \neq i} v_j(\omega^*, \theta_j) \ge 0.$$

VCG mechanism may not be implementable in polynomial time.

Welfare Maximization

Implementing the VCG mechanism requires solving the optimal allocation problem:

$$\omega^* = \operatorname*{argmax}_{\omega \in \Omega} \sum_i v_i(\omega, \theta_i).$$

Welfare Maximization

Implementing the VCG mechanism requires solving the optimal allocation problem:

$$\omega^* = \underset{\omega \in \Omega}{\operatorname{argmax}} \sum_i v_i(\omega, \theta_i).$$

Is this tractable in practice? NO!

Welfare Maximization

Implementing the VCG mechanism requires solving the optimal allocation problem:

$$\omega^* = \underset{\omega \in \Omega}{\operatorname{argmax}} \sum_i v_i(\omega, \theta_i).$$

Is this tractable in practice? NO!

Example: (Knapsack problem) consider the allocation problem of servicing agents, where $\Omega \subset 2^N$.

- each agent has private value θ_i for being serviced;
- servicing each agent i requires a resource of r_i ;
- there is a total budget of B on resource;
- allocation ω is feasible if and only if $\sum_{i \in \omega} r_i \leq B$.

Welfare Maximization

Implementing the VCG mechanism requires solving the optimal allocation problem:

$$\omega^* = \underset{\omega \in \Omega}{\operatorname{argmax}} \sum_i v_i(\omega, \theta_i).$$

Is this tractable in practice? NO!

Example: (Knapsack problem) consider the allocation problem of servicing agents, where $\Omega \subset 2^N$.

- each agent has private value θ_i for being serviced;
- servicing each agent i requires a resource of r_i ;
- there is a total budget of B on resource;
- allocation ω is feasible if and only if $\sum_{i \in \omega} r_i \leq B$.

How to find the optimal allocation? Trying all combination requires time exponential in |N|. Not practical if n=|N| is large!

An algorithm is a polynomial-time algorithm if there exists $c \in (0, \infty)$ such that its running time f(n) satisfies $f(n) = O(n^c)$.

An algorithm is a polynomial-time algorithm if there exists $c \in (0, \infty)$ such that its running time f(n) satisfies $f(n) = O(n^c)$.

Enumerating all subsets of N is not a polynomial-time algorithm: $2^n = \omega(n^c)$ for any $c < \infty$.

An algorithm is a polynomial-time algorithm if there exists $c \in (0, \infty)$ such that its running time f(n) satisfies $f(n) = O(n^c)$.

Enumerating all subsets of N is not a polynomial-time algorithm: $2^n = \omega(n^c)$ for any $c < \infty$.

Under the assumption that $P\neq NP$, the knapsack problem does not have any polynomial-time algorithm.

• There exist polynomial-time algorithms for approximating the optimal solutions.

An algorithm is a polynomial-time algorithm if there exists $c \in (0, \infty)$ such that its running time f(n) satisfies $f(n) = O(n^c)$.

Enumerating all subsets of N is not a polynomial-time algorithm: $2^n = \omega(n^c)$ for any $c < \infty$.

Under the assumption that $P\neq NP$, the knapsack problem does not have any polynomial-time algorithm.

• There exist polynomial-time algorithms for approximating the optimal solutions.

Theorem

The maximum of greedy algorithm and max-value is a 2-approximation to the optimal value in the knapsack problem.

Example: 3D Matching

3D Matching: serving each agent requires two types of resources. N: agents; X: resource type 1; Y: resource type 2.

- $L = \{(i, x, y)\}$: the set of feasible ways to serve the agents;
- find the maximum number of agents that can be served simultaneously.

Example: 3D Matching

3D Matching: serving each agent requires two types of resources. N: agents; X: resource type 1; Y: resource type 2.

- $L = \{(i, x, y)\}$: the set of feasible ways to serve the agents;
- find the maximum number of agents that can be served simultaneously.

Under the assumption that $P\neq NP$, the 3D matching problem does not have any polynomial-time algorithm.

Example: 3D Matching

3D Matching: serving each agent requires two types of resources. N: agents; X: resource type 1; Y: resource type 2.

- $L = \{(i, x, y)\}$: the set of feasible ways to serve the agents;
- find the maximum number of agents that can be served simultaneously.

Under the assumption that $P \neq NP$, the 3D matching problem does not have any polynomial-time algorithm.

Theorem

The greedy algorithm for finding the maximal matching is a 3-approximation to the optimal.

Intuition: in the greedy algorithm, when an agent is served, it will exclude at most two additional agents from the optimal matching.

The design of approximation algorithms does not take incentives of agents into consideration.

The design of approximation algorithms does not take incentives of agents into consideration.

Question: does there exist polynomial-time mechanism that guarantees good welfare approximations?

• A tricky issue: given an approximately optimal algorithm, there may not exist any mechanism that implements its allocation.

The design of approximation algorithms does not take incentives of agents into consideration.

Question: does there exist polynomial-time mechanism that guarantees good welfare approximations?

• A tricky issue: given an approximately optimal algorithm, there may not exist any mechanism that implements its allocation.

Theorem

For any $\beta \geq 1$ and given any polynomial time algorithm with approximation ratio β , there exists a polynomial time mechanism that achieves a β -approximation to the optimal welfare.

The design of approximation algorithms does not take incentives of agents into consideration.

Question: does there exist polynomial-time mechanism that guarantees good welfare approximations?

• A tricky issue: given an approximately optimal algorithm, there may not exist any mechanism that implements its allocation.

Theorem

For any $\beta \geq 1$ and given any polynomial time algorithm with approximation ratio β , there exists a polynomial time mechanism that achieves a β -approximation to the optimal welfare.

Idea: efficiency by matching.

- see illustration on board;
- apply efficiency in general equilibrium models to prove the reduction.

The design of approximation algorithms does not take incentives of agents into consideration.

Question: does there exist polynomial-time mechanism that guarantees good welfare approximations?

• A tricky issue: given an approximately optimal algorithm, there may not exist any mechanism that implements its allocation.

Theorem

For any $\beta \geq 1$ and given any polynomial time algorithm with approximation ratio β , there exists a polynomial time mechanism that achieves a β -approximation to the optimal welfare.

Idea: efficiency by matching.

- see illustration on board;
- apply efficiency in general equilibrium models to prove the reduction.

Reference: Hartline, J. D., Kleinberg, R., & Malekian, A. (2015). Bayesian incentive compatibility via matchings. Games and Economic Behavior, 92, 401-429.

General Equilibrium

Consider a market with n agents and n items.

- ullet each agent i has unit value v_{ij} for item j;
- ullet each agent i has demand at most f_i ;
- each item j has supply at most g_j ;
- $\sum_{i} f_{i} = \sum_{j} g_{j} = 1$.

General Equilibrium

Consider a market with n agents and n items.

- ullet each agent i has unit value v_{ij} for item j;
- each agent i has demand at most f_i ;
- each item j has supply at most g_j ;
- $\sum_{i} f_{i} = \sum_{j} g_{j} = 1$.

Theorem

There exists a price p_j on each item j such that when each agent purchases their favorite consumption bundle,

- the allocation is efficient:
- supply meets the demand, i.e., all items are sold out and all agents purchase up to their demand.

General Equilibrium

Consider a market with n agents and n items.

- ullet each agent i has unit value v_{ij} for item j;
- ullet each agent i has demand at most f_i ;
- each item j has supply at most g_j ;
- $\sum_{i} f_{i} = \sum_{j} g_{j} = 1$.

Theorem

There exists a price p_j on each item j such that when each agent purchases their favorite consumption bundle,

- the allocation is efficient:
- supply meets the demand, i.e., all items are sold out and all agents purchase up to their demand.

Intuition: use tâtonnement rule to adjust the price

• gradually increase the price of the item with excessive demand.

Revenue Approximations

Posted Pricing

Posted pricing mechanisms: offer price p_i to agent i. The item is sold to the first agent who is willing to purchase.

• this is in fact order oblivious posted pricing where the seller cannot control the order of the agents.

Posted Pricing

Posted pricing mechanisms: offer price p_i to agent i. The item is sold to the first agent who is willing to purchase.

• this is in fact order oblivious posted pricing where the seller cannot control the order of the agents.

Question: is posted pricing mechanisms also approximately optimal for revenue maximization?

Recap: Prophet Inequality

Online Selection Problem: n items arriving online.

- item i has value $v_i \sim F_i$;
- the agent knows F_1, \ldots, F_n at time 0.
- at time $i \leq n$, the agent observes value v_i and decides whether to select item i (if the selection has not been made).

Theorem

There exists a threshold policy that achieves a 2-approximation, i.e., it achieves expected value at least $\frac{1}{2}\mathbf{E}[\max_i v_i]$.

Connection to Revenue Maximization

Prophet inequality: n items

- value distributions $F = F_1 \times \cdots \times F_n$;
- threshold τ for each item;
- arrival order π .

Posted pricing mechanism: n agents

- marginal revenues $F = F_1 \times \cdots \times F_n$;
- threshold τ for each agent i;
- tie breaking rule π .

Any threshold au in the marginal revenue space corresponds to a price p_i in the value space.

Connection to Revenue Maximization

Prophet inequality: *n* items

- value distributions $F = F_1 \times \cdots \times F_n$;
- threshold τ for each item;
- arrival order π .

Posted pricing mechanism: n agents

- marginal revenues $F = F_1 \times \cdots \times F_n$;
- threshold τ for each agent i;
- tie breaking rule π .

Any threshold τ in the marginal revenue space corresponds to a price p_i in the value space.

Given any valuation profile $v=(v_1,\ldots,v_n)$, the selected value and the optimal value in both problems are the same.

• Posted pricing mechanism has a 2-approximation to the expected maximum marginal revenue.

Connection to Revenue Maximization

Prophet inequality: *n* items

- value distributions $F = F_1 \times \cdots \times F_n$;
- threshold τ for each item;
- arrival order π .

Posted pricing mechanism: n agents

- marginal revenues $F = F_1 \times \cdots \times F_n$;
- threshold τ for each agent i;
- tie breaking rule π .

Any threshold au in the marginal revenue space corresponds to a price p_i in the value space.

Given any valuation profile $v=(v_1,\ldots,v_n)$, the selected value and the optimal value in both problems are the same.

• Posted pricing mechanism has a 2-approximation to the expected maximum marginal revenue.

expected marginal revenue = expected revenue

 \Rightarrow posted pricing mechanism has a 2-approximation to the expected revenue.

Third-degree Price Discrimination

Can we approximate the optimal revenue without relying on third-degree price discrimination, i.e., posting an anonymous price?

Third-degree Price Discrimination

Can we approximate the optimal revenue without relying on third-degree price discrimination, i.e., posting an anonymous price?

Example: n agents. For agent $i \leq n$, $v_i = 2^i$ w.p $\frac{1}{2^{i+1}}$, and $v_i = 0$ w.p $1 - \frac{1}{2^{i+1}}$.

- optimal revenue $\Theta(n)$;
- anonymous pricing O(1).

Third-degree Price Discrimination

Can we approximate the optimal revenue without relying on third-degree price discrimination, i.e., posting an anonymous price?

Example: n agents. For agent $i \leq n$, $v_i = 2^i$ w.p $\frac{1}{2^{i+1}}$, and $v_i = 0$ w.p $1 - \frac{1}{2^{i+1}}$.

- optimal revenue $\Theta(n)$;
- anonymous pricing O(1).

Third-degree price discrimination is crucial for revenue maximization.

• competition and simultaneous implementation is not.

Sequential Posted Pricing

Can we do better if the seller can control the order of approaching the agents? Yes!

Sequential Posted Pricing

Can we do better if the seller can control the order of approaching the agents? Yes!

Sequential posted pricing: design an order π and prices p_i for each agent i.

- Agents arrive according to order π ;
- The item is sold to the first agent who is willing to purchase.

Sequential Posted Pricing

Can we do better if the seller can control the order of approaching the agents? Yes!

Sequential posted pricing: design an order π and prices p_i for each agent i.

- Agents arrive according to order π ;
- The item is sold to the first agent who is willing to purchase.

Theorem (Yan '11)

Sequential posted pricing mechanism has an $\frac{e}{e-1}$ -approximation to the expected revenue.

A non-negative real-valued set function f over subsets S of an n element ground set $N = \{1, \dots, n\}$ and a distribution over subsets given by \mathcal{D} .

- \hat{q}_i : ex ante probability that element i is in the random set $S \sim \mathcal{D}$
- \mathcal{D}^I : distribution over subsets induced by independently adding each element i to the set with probability equal to its ex ante probability \hat{q}_i .

A non-negative real-valued set function f over subsets S of an n element ground set $N=\{1,\ldots,n\}$ and a distribution over subsets given by $\mathcal{D}.$

- ullet \hat{q}_i : ex ante probability that element i is in the random set $S \sim \mathcal{D}$
- \mathcal{D}^I : distribution over subsets induced by independently adding each element i to the set with probability equal to its ex ante probability \hat{q}_i .

The correlation gap is the ratio of the expected value of the set function for the (correlated) distribution \mathcal{D} to that with independent distribution \mathcal{D}^I , i.e.,

$$\frac{E_{S \sim \mathcal{D}}[f(S)]}{E_{S \sim \mathcal{D}^I}[f(S)]}.$$

Definition

A set function $f: 2^S \to \mathbb{R}$ defined on the subsets of a finite set S is called submodular if for all $A \subseteq B \subseteq S$ and $x \notin B$, the following inequality holds:

$$f(A \cup \{x\}) - f(A) \ge f(B \cup \{x\}) - f(B).$$

Submodular functions captures decreasing marginal return.

Definition

A set function $f: 2^S \to \mathbb{R}$ defined on the subsets of a finite set S is called submodular if for all $A \subseteq B \subseteq S$ and $x \notin B$, the following inequality holds:

$$f(A \cup \{x\}) - f(A) \ge f(B \cup \{x\}) - f(B).$$

Submodular functions captures decreasing marginal return.

Theorem

If the set function f is submodular, the correlation gap for function f is at most $\frac{e}{e-1}$.

Example: submodular function $f(S) = \mathbf{1} (S \neq \emptyset)$.

• ex ante probability $\hat{q}_i = \frac{1}{n}$ for all i.

Correlation Gap

Example: submodular function $f(S) = \mathbf{1} (S \neq \emptyset)$.

• ex ante probability $\hat{q}_i = \frac{1}{n}$ for all i.

Under correlated distribution \mathcal{D} , it is optimal that only one item is selected in each realization:

$$E_{S \sim \mathcal{D}}[f(S)] = 1.$$

Correlation Gap

Example: submodular function $f(S) = \mathbf{1} (S \neq \emptyset)$.

• ex ante probability $\hat{q}_i = \frac{1}{n}$ for all i.

Under correlated distribution \mathcal{D} , it is optimal that only one item is selected in each realization:

$$E_{S \sim \mathcal{D}}[f(S)] = 1.$$

Under independent distribution \mathcal{D}^I :

$$E_{S \sim \mathcal{D}^I}[f(S)] = 1 - (1 - \frac{1}{n})^n \xrightarrow{n \to \infty} 1 - \frac{1}{e}.$$

Correlation Gap

Example: submodular function $f(S) = \mathbf{1} (S \neq \emptyset)$.

• ex ante probability $\hat{q}_i = \frac{1}{n}$ for all i.

Under correlated distribution \mathcal{D} , it is optimal that only one item is selected in each realization:

$$E_{S \sim \mathcal{D}}[f(S)] = 1.$$

Under independent distribution \mathcal{D}^I :

$$E_{S \sim \mathcal{D}^I}[f(S)] = 1 - (1 - \frac{1}{n})^n \xrightarrow{n \to \infty} 1 - \frac{1}{e}.$$

As
$$n \to \infty$$
, $\frac{E_{S \sim \mathcal{D}}[f(S)]}{E_{S \sim \mathcal{D}^I}[f(S)]} = \frac{e}{e-1}$.

Ex ante relaxation: consider the relaxed problem where the sum of ex ante probabilities of receiving an item is at most 1.

$$EAR = \sum_{i} R_i(q_i) \quad \text{s.t.} \quad \sum_{i} q_i \le 1.$$

EAR is an upper bound on the optimal revenue.

Ex ante relaxation: consider the relaxed problem where the sum of ex ante probabilities of receiving an item is at most 1.

$$EAR = \sum_{i} R_i(q_i) \quad \text{s.t.} \quad \sum_{i} q_i \le 1.$$

EAR is an upper bound on the optimal revenue.

Let \hat{q}_i be the optimal solution for EAR.

Ex ante relaxation: consider the relaxed problem where the sum of ex ante probabilities of receiving an item is at most 1.

$$EAR = \sum_{i} R_i(q_i) \quad \text{s.t.} \quad \sum_{i} q_i \le 1.$$

EAR is an upper bound on the optimal revenue.

Let \hat{q}_i be the optimal solution for EAR.

Sequential posted pricing: offer price \hat{p}_i that sells with probability \hat{q}_i to agent i;

• agents are approached in decreasing order of \hat{p}_i .

Let function $f(S) = \max_{i \in S} \hat{p}_i$. f is submodular.

Let function $f(S) = \max_{i \in S} \hat{p}_i$. f is submodular.

Given marginal probability \hat{q}_i on each i:

- $E_{S \sim \mathcal{D}^I}[f(S)]$: expected revenue from sequential posted pricing;
- $E_{S \sim \mathcal{D}}[f(S)]$: EAR.

Let function $f(S) = \max_{i \in S} \hat{p}_i$. f is submodular.

Given marginal probability \hat{q}_i on each i:

- $E_{S \sim \mathcal{D}^I}[f(S)]$: expected revenue from sequential posted pricing;
- $E_{S \sim \mathcal{D}}[f(S)]$: EAR.

Correlation gap implies that

$$\frac{E_{S \sim \mathcal{D}}[f(S)]}{E_{S \sim \mathcal{D}^I}[f(S)]} \le \frac{e}{e - 1}.$$

Extension of Approximations Under Non-linear Utilities

Two options, which one would you choose:

- get \$10M;
- draw a lottery, with probability $\frac{1}{2}$, get \$20M, and get nothing otherwise.

In practice, buyers have non-linear utilities: e.g., risk aversion, budget constraints, and etc.

In practice, buyers have non-linear utilities: e.g., risk aversion, budget constraints, and etc.

Risk aversion: $t_i = (v_i, \varphi_i)$ where $v_i \in \mathbb{R}_+$, φ_i is an increasing concave function, and

$$u_i(t_i, x_i, p_i) = \varphi_i(v_i x_i - p_i).$$

Private budgets: $t_i = (v_i, B_i)$ where $v_i, B_i \in \mathbb{R}_+$, and

$$u_i(t_i, x_i, p_i) = \begin{cases} v_i x_i - p_i & p_i \le B_i \\ -\infty & p_i > B_i. \end{cases}$$

In practice, buyers have non-linear utilities: e.g., risk aversion, budget constraints, and etc.

Risk aversion: $t_i = (v_i, \varphi_i)$ where $v_i \in \mathbb{R}_+$, φ_i is an increasing concave function, and

$$u_i(t_i, x_i, p_i) = \varphi_i(v_i x_i - p_i).$$

Private budgets: $t_i = (v_i, B_i)$ where $v_i, B_i \in \mathbb{R}_+$, and

$$u_i(t_i, x_i, p_i) = \begin{cases} v_i x_i - p_i & p_i \le B_i \\ -\infty & p_i > B_i. \end{cases}$$

Are simple mechanisms approximately optimal for non-linear utilities?

In single-agent environments, a mechanism is posting a per-unit price p if the agent can purchase any lottery x with price $x \cdot p$ for any $x \in [0,1]$.

• agent pays price $x \cdot p$ even if the realized allocation is 0.

In single-agent environments, a mechanism is posting a per-unit price p if the agent can purchase any lottery x with price $x \cdot p$ for any $x \in [0,1]$.

• agent pays price $x \cdot p$ even if the realized allocation is 0.

Definition (Demands)

The demand of the agent $d^u(t,p)$ is the optimal lottery of the agent given per-unit price p.

In single-agent environments, a mechanism is posting a per-unit price p if the agent can purchase any lottery x with price $x \cdot p$ for any $x \in [0,1]$.

• agent pays price $x \cdot p$ even if the realized allocation is 0.

Definition (Demands)

The demand of the agent $d^u(t,p)$ is the optimal lottery of the agent given per-unit price p.

Remark: We denote $D^u(p) = \mathbf{E}_{t \sim F}[d^u(t,p)]$ as the expected demand, and omit u in notations when it is clear from context.

In single-agent environments, a mechanism is posting a per-unit price p if the agent can purchase any lottery x with price $x \cdot p$ for any $x \in [0,1]$.

• agent pays price $x \cdot p$ even if the realized allocation is 0.

Definition (Demands)

The demand of the agent $d^u(t,p)$ is the optimal lottery of the agent given per-unit price p.

Remark: We denote $D^u(p) = \mathbf{E}_{t \sim F}[d^u(t,p)]$ as the expected demand, and omit u in notations when it is clear from context.

Assumption (Ordinary Goods)

 $d^u(t,p)$ is non-increasing in p for all $t \in T$.

Excludes Giffen goods or Veblen goods.

Recall for linear utilities: let $q = \mathbf{Pr}_{t' \sim F}[t' \geq t]$ be the quantile for type t.

• lower quantile \Leftrightarrow higher willingness to pay.

Recall for linear utilities: let $q = \mathbf{Pr}_{t' \sim F}[t' \geq t]$ be the quantile for type t.

• lower quantile ⇔ higher willingness to pay.

Type space for non-linear agents can be multi-dimensional.

Recall for linear utilities: let $q = \mathbf{Pr}_{t' \sim F}[t' \geq t]$ be the quantile for type t.

• lower quantile ⇔ higher willingness to pay.

Type space for non-linear agents can be multi-dimensional.

Demand of an agent with a private budget given per-unit pricing:

	$t_1: (v=5, B=1)$	$t_2: (v=2, B=2)$
$p_1 = 4$	0.25	0
$p_2 = 2$	0.5	1

Recall for linear utilities: let $q = \mathbf{Pr}_{t' \sim F}[t' \geq t]$ be the quantile for type t.

• lower quantile ⇔ higher willingness to pay.

Type space for non-linear agents can be multi-dimensional.

Demand of an agent with a private budget given per-unit pricing:

	$t_1: (v=5, B=1)$	$t_2: (v=2, B=2)$
$p_1 = 4$	0.25	0
$p_2 = 2$	0.5	1

There isn't a simple deterministic and consistent way of ordering types for a non-linear agent.

Solution: a random mapping from types to quantiles based on demand functions.

Solution: a random mapping from types to quantiles based on demand functions.

Intuition: given any threshold \hat{q} and any type t, the following two quantities should coincide

- **1** probability the quantile of t is below \hat{q} ;
- 2 the demand of t given market clearing price $p^{\hat{q}}$.

Solution: a random mapping from types to quantiles based on demand functions.


Intuition: given any threshold \hat{q} and any type t, the following two quantities should coincide

- probability the quantile of t is below \hat{q} ;
- 2 the demand of t given market clearing price $p^{\hat{q}}$.

Definition (Quantiles for Non-linear Agents)

The randomized quantile q for type $t \in T$ is drawn from distribution with CDF $d(t, p^q)$.

- $d(t, p^0) = 0$ and $d(t, p^1) = 1$;
- $d(t, p^q)$ is weakly increasing in q for all type t (ordinary good assumption).



Solution: a random mapping from types to quantiles based on demand functions.

Intuition: given any threshold \hat{q} and any type t, the following two quantities should coincide


- probability the quantile of t is below \hat{q} ;
- 2 the demand of t given market clearing price $p^{\hat{q}}$.

Definition (Quantiles for Non-linear Agents)

The randomized quantile q for type $t \in T$ is drawn from distribution with CDF $d(t, p^q)$.

- $d(t, p^0) = 0$ and $d(t, p^1) = 1$;
- $d(t, p^q)$ is weakly increasing in q for all type t (ordinary good assumption).

Remark: $q \sim U[0,1]$: $\Pr[z \leq q] = \mathbf{E}_{t \sim F}[d(t, p^q)] = q$.



Pricing-based Mechanisms in Quantile Space

Definition (Pricing-based Mechanisms in Quantile Space)

Given any profile of feasible thresholds $\{Q_i\}_{i\in[n]}$,

- **1** Map type t_i to quantile q_i according to $d(t, p^q)$, and calculate threshold as $\hat{q}_i = Q_i(q_{-i})$.
- 2 The allocation of agent i is $x_i = 1$ if and only if $q_i \leq \hat{q}_i$. The payment of agent i is $p_i = p^{\hat{q}_i} \cdot d(t_i, p^{\hat{q}_i})$ regardless of the allocation.

Pricing-based Mechanisms in Quantile Space

Definition (Pricing-based Mechanisms in Quantile Space)

Given any profile of feasible thresholds $\{Q_i\}_{i\in[n]}$,

- ① Map type t_i to quantile q_i according to $d(t, p^q)$, and calculate threshold as $\hat{q}_i = Q_i(q_{-i})$.
- ② The allocation of agent i is $x_i=1$ if and only if $q_i \leq \hat{q}_i$. The payment of agent i is $p_i=p^{\hat{q}_i}\cdot d(t_i,p^{\hat{q}_i})$ regardless of the allocation.

Price-posting Equivalence Interpretation: Fixing any \hat{q}_i , from perspective of agent i

- wins the item with probability $d(t_i, p^{\hat{q}_i})$;
- deterministically pays $p^{\hat{q}_i} \cdot d(t_i, p^{\hat{q}_i})$.

Equivalent to posting per-unit price $p^{\hat{q}_i}$ to agent i.

Pricing-based Mechanisms in Quantile Space

Definition (Pricing-based Mechanisms in Quantile Space)

Given any profile of feasible thresholds $\{Q_i\}_{i\in[n]}$,

- ① Map type t_i to quantile q_i according to $d(t, p^q)$, and calculate threshold as $\hat{q}_i = Q_i(q_{-i})$.
- ② The allocation of agent i is $x_i=1$ if and only if $q_i \leq \hat{q}_i$. The payment of agent i is $p_i=p^{\hat{q}_i}\cdot d(t_i,p^{\hat{q}_i})$ regardless of the allocation.

Price-posting Equivalence Interpretation: Fixing any \hat{q}_i , from perspective of agent i

- wins the item with probability $d(t_i, p^{\hat{q}_i})$;
- deterministically pays $p^{\hat{q}_i} \cdot d(t_i, p^{\hat{q}_i})$.

Equivalent to posting per-unit price $p^{\hat{q}_i}$ to agent i.

Expected revenue: from any pricing-based mechanisms M for non-linear agents,

$$M(P) = \sum_{i} \mathbf{E}_{\forall j \neq i, q_{j} \sim U[0,1]} [P_{i}(Q_{i}(q_{-i}))].$$

For linear agents: $R = \bar{P}$ [Bulow and Robert '89].

For non-linear agents: pricing-based mechanisms in general are not optimal, i.e., $R \neq \bar{P}$.

For linear agents: $R = \bar{P}$ [Bulow and Robert '89].

For non-linear agents: pricing-based mechanisms in general are not optimal, i.e., $R \neq \bar{P}$.

Example: an agent with private budget:

- with probability $\frac{1}{2}$, $t_1 : (v = 2, B = 1)$;
- with probability $\frac{1}{2}$, $t_2 : (v = 10, B = 3)$.

For linear agents: $R = \bar{P}$ [Bulow and Robert '89].

For non-linear agents: pricing-based mechanisms in general are not optimal, i.e., $R \neq \bar{P}$.

Example: an agent with private budget:

- with probability $\frac{1}{2}$, $t_1 : (v = 2, B = 1)$;
- with probability $\frac{1}{2}$, $t_2 : (v = 10, B = 3)$.

Revenue from posted pricing:

- if p > 2: only sold to $t_2 \Rightarrow$ revenue is at most $\frac{3}{2}$;
- if $p \leq 2$: payment from t_2 is at most $2 \Rightarrow$ revenue is at most $\frac{3}{2}$.

For linear agents: $R = \bar{P}$ [Bulow and Robert '89].

For non-linear agents: pricing-based mechanisms in general are not optimal, i.e., $R \neq \bar{P}$.

Example: an agent with private budget:

- with probability $\frac{1}{2}$, $t_1 : (v = 2, B = 1)$;
- with probability $\frac{1}{2}$, $t_2 : (v = 10, B = 3)$.

Revenue from posted pricing:

- if p > 2: only sold to $t_2 \Rightarrow$ revenue is at most $\frac{3}{2}$;
- if $p \leq 2$: payment from t_2 is at most $2 \Rightarrow$ revenue is at most $\frac{3}{2}$.


Lottery mechanism:

- offer menu of lotteries $(x_1 = \frac{1}{2}, p_1 = 1)$ and $(x_2 = 1, p_2 = 3)$;
- expected revenue equals 2.

Resemblance: Approximations in Single-agent Settings

Definition (ζ -resemblance)


A non-linear agent is ζ -resemblant to a linear agent if given any supply constraint $q \in [0,1]$, there exists a posted pricing mechanism with expected demand $q^\dagger \leq q$ such that $\bar{P}(q^\dagger) \geq \frac{1}{\zeta} R(q)$.



Resemblance: Approximations in Single-agent Settings

Definition (ζ -resemblance)


A non-linear agent is ζ -resemblant to a linear agent if given any supply constraint $q \in [0,1]$, there exists a posted pricing mechanism with expected demand $q^\dagger \leq q$ such that $\bar{P}(q^\dagger) \geq \frac{1}{\zeta} R(q)$.



Resemblance: Approximations in Single-agent Settings

Definition (ζ -resemblance)

A non-linear agent is ζ -resemblant to a linear agent if given any supply constraint $q \in [0,1]$, there exists a posted pricing mechanism with expected demand $q^\dagger \leq q$ such that $\bar{P}(q^\dagger) \geq \frac{1}{\zeta} R(q)$.



Reduction from Non-linear to Linear Agents

Theorem

For non-linear agents that are ζ -resemblant to linear agents, pricing-based mechanism M is a γ -approximation to ex ante relaxation for linear agents $\Rightarrow M$ is a $\zeta \gamma$ -approximation to ex ante relaxation for non-linear agents.

- expected revenue only depends on the price-posting revenue curves

Reduction from Non-linear to Linear Agents

Theorem

For non-linear agents that are ζ -resemblant to linear agents, pricing-based mechanism M is a γ -approximation to ex ante relaxation for linear agents $\Rightarrow M$ is a $\zeta\gamma$ -approximation to ex ante relaxation for non-linear agents.

- expected revenue only depends on the price-posting revenue curves

Reduction from Non-linear to Linear Agents

Theorem

For non-linear agents that are ζ -resemblant to linear agents, pricing-based mechanism M is a γ -approximation to ex ante relaxation for linear agents $\Rightarrow M$ is a $\zeta\gamma$ -approximation to ex ante relaxation for non-linear agents.

- expected revenue only depends on the price-posting revenue curves

Non-linearities are often details that can be dispensed from the model without affecting main economic conclusions.

Economic conclusions for linear agents \Rightarrow economic conclusions for non-linear agents.

ζ -resemblance for Non-linear Agents

	independent private budget	risk averse
revenue	3	е
welfare	2	1

Table: Summary of results for ζ -resemblance, assuming regularity for budgeted utility and MHR for risk averse utility under revenue objective.

ζ -resemblance for Non-linear Agents

	independent private budget	risk averse
revenue	3	е
welfare	2	1

Table: Summary of results for ζ -resemblance, assuming regularity for budgeted utility and MHR for risk averse utility under revenue objective.

Corollary

For risk averse agents, sequential posted pricing is an e/(e-1)-approximation to the optimal welfare.