

Note for Scattering Amplitude Computation

Su Yingze

March 21, 2025

1 4-point case

For the four point case $\mathcal{A}(V_2 \Phi^\dagger V_1 \Phi)$, we can construct the color-ordered amplitude from the residue. First, we consider the $(+, -)$ helicity configuration. There are two feynman diagrams contributing to the color-ordered amplitude.




Figure 1: 4pt.

For the first diagram, the residue equals to

$$\mathcal{R}es|_{s_{12}=0} = \frac{[3I][23]}{[I2]} \times \frac{\langle I4 \rangle \langle 41 \rangle}{\langle 1I \rangle} = \frac{\langle 24 \rangle [31] \langle 41 \rangle [23]}{[42] \langle 24 \rangle}$$

Similarly, the second one is

$$\mathcal{R}es|_{s_{13}=0} = \frac{\langle 4I \rangle \langle 24 \rangle}{\langle I2 \rangle} \times \frac{[31][I3]}{[1I]} = \frac{\langle 24 \rangle [31] \langle 41 \rangle [23]}{\langle 32 \rangle [23]}$$

Then we can conclude that the four-point color-ordered amplitude $A[1, 2, 3^+, 4^-]$ equals to

$$A[1, 2, 3^+, 4^-] = \frac{\langle 24 \rangle [31] \langle 41 \rangle [23]}{\langle 32 \rangle [23][42] \langle 24 \rangle} = \frac{\langle 24 \rangle \langle 14 \rangle}{\langle 13 \rangle \langle 23 \rangle}$$

★Bonus

It is still necessary to prove the color-ordered amplitude $A[1, 2, 3^+, 4^+]$ equals to 0. Here we can use the color ordered Feynman rules to show the result.

$$A[1, 2, 3^+, 4^+] \propto \frac{(\epsilon_3 \cdot p_2)(\epsilon_4 \cdot p_1)}{s_{23}} + \frac{(\epsilon_4 \cdot p_2)(\epsilon_3 \cdot p_1)}{s_{24}}$$

Here we can utilize the spinor-helicity variable to express polarization vector

$$\epsilon_2^{+\mu} = \frac{\langle r_1 | \gamma^\mu | 3 \rangle}{\sqrt{2} \langle r_1 3 \rangle}, \quad \epsilon_4^{+\mu} = \frac{\langle r_2 | \gamma^\mu | 4 \rangle}{\sqrt{2} \langle r_2 4 \rangle}$$

here r_1 and r_2 represent the reference spinor.

We can freely choose $r_1 = r_2 = 1$ or 2 , then $\langle r_1 2 \rangle, \langle r_2 1 \rangle, \langle r_1 1 \rangle, \langle r_2 2 \rangle$, two of them equal to 0, so we can conclude that

$$A[1, 2, 3^+, 4^+] = 0$$

2 5-point case

For the 5-point case, we can utilize the BCFW recursion relation which can help us generate higher point amplitude from lower point on-shell subamplitudes. Here, we always consider the MHV (Maximal helicity violation) amplitude.

Let us begin with the simplest case $A[1, 2, 3_1^+, 4_1^+, 5_2^-]$, where the subscript represent which gauge group the particle belongs to. Because of the property of this kind of gauge theory, the color structure is invariant under the OPP (Order Preserving Permutation), in this case, for example,

$$(3_1^+, 4_1^+, 5_2^-) \quad (3_1^+, 5_2^-, 4_1^+) \quad (5_2^-, 3_1^+, 4_1^+)$$

give us the same color factor. So in the process of BCFW recursion, these three order offer the same amplitude.

The amplitude equals to

$$\begin{aligned} A[1, 2, 3_1^+, 4_1^+, 5_2^-] &= A[1, 2, \hat{I}^+, 5^-] \times \frac{1}{s_{34}} \times A[\hat{3}^+, 4^+, \hat{I}^-] \\ &= \frac{\langle 15 \rangle \langle 25 \rangle}{\langle 1\hat{I} \rangle \langle 2\hat{I} \rangle} \times \frac{1}{s_{34}} \times \frac{[\hat{3}4]^3}{[4\hat{I}][\hat{I}3]} \\ &= \frac{\langle 15 \rangle \langle 25 \rangle [\hat{3}4]^3}{\langle 14 \rangle \langle 23 \rangle \langle 43 \rangle [\hat{4}3][\hat{4}3][\hat{3}4]} \\ &= \frac{\langle 15 \rangle \langle 25 \rangle}{\langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \\ &= \frac{(-1) \langle 2\hat{5} \rangle^2 \langle 1\hat{5} \rangle^2}{\langle 23 \rangle \langle 34 \rangle \langle 41 \rangle \langle 2\hat{5} \rangle \langle 1\hat{5} \rangle} \end{aligned}$$

where we use the fact $|\hat{3}\rangle = |\hat{3}\rangle$ and following identities

$$\langle 1\hat{I} \rangle [\hat{I}3] = \langle 14 \rangle [43], \quad \langle 2\hat{I} \rangle [\hat{I}3] = \langle 24 \rangle [43]$$

and also

$$\begin{aligned} \frac{[\hat{I}3]}{[4\hat{I}]} &= -\frac{[3\hat{I}]\langle \hat{I}2 \rangle}{[4\hat{I}]\langle \hat{I}2 \rangle} = -\frac{[34]\langle 42 \rangle}{[43]\langle \hat{3}2 \rangle}, \quad (\langle \hat{3}2 \rangle = \langle 32 \rangle + z \langle 22 \rangle = \langle 32 \rangle) \\ &= \frac{\langle 42 \rangle}{\langle 32 \rangle} \end{aligned}$$

here green refers to the particle with (-) helicity, red refers to particles belong to gauge group 1, red refers to particles belong to gauge group 2.

Similarly, it is very easy to obtain another color-ordered amplitude $A[1, 2, 3_1^+, 4_1^-, 5_2^+]$

$$A[1, 2, 3_1^+, 4_1^-, 5_2^+] = \frac{(-1) \langle 2\textcolor{green}{4} \rangle^2 \langle 1\textcolor{green}{4} \rangle^2}{\langle \textcolor{blue}{23} \rangle \langle \textcolor{blue}{34} \rangle \langle \textcolor{blue}{41} \rangle \langle \textcolor{red}{25} \rangle \langle \textcolor{red}{51} \rangle}$$

and also $A[1, 2, 3_1^-, 4_1^+, 5_2^+]$ equals to

$$A[1, 2, 3_1^-, 4_1^+, 5_2^+] = \frac{(-1) \langle 2\textcolor{green}{3} \rangle^2 \langle 1\textcolor{green}{3} \rangle^2}{\langle \textcolor{blue}{23} \rangle \langle \textcolor{blue}{34} \rangle \langle \textcolor{blue}{41} \rangle \langle \textcolor{red}{25} \rangle \langle \textcolor{red}{51} \rangle}$$

But here we need to emphasize that it is necessary to choose another BCFW shift, like $[1, 5]$, as $[2, 3^-]$ is not a valid shift.

3 6-point case

Here we consider $(V_2 V_2 \Phi^\dagger V_1 V_1 \Phi)$ case, the corresponding color-ordered amplitude is $A[1, 2, 3_1^+, 4_1^+, 5_2^+, 6_2^-]$