

Some Details

Su Yingze

November 26, 2025

1 Normalization of Polarization Vectors

First, we give the definition of polarization vectors which is written in terms of adjoint representation

$$\varepsilon_a^\mu = \frac{\langle p_I | (r \cdot \Gamma) \Gamma^\mu | p^J \rangle (\sigma_a)^I_J}{4(p \cdot r)} \quad (1)$$

here a runs from 1 to 3. We can use antisymmetric tensor to rewrite this equation, like

$$\varepsilon_a^\mu = \frac{\langle p_I | (r \cdot \Gamma) \Gamma^\mu | p_K \rangle \epsilon^{JK} (\sigma_a)^I_J}{4(p \cdot r)} \quad (2)$$

here we choose the left-multiplication convention.

Then, a 2×2 symmetric matrix can be defined like

$$(\sigma_a)^{IJ} = \epsilon^{JK} (\sigma_a)^I_K. \quad (3)$$

This means that the original polarization vector can be understood as

$$\begin{aligned} \varepsilon_a^\mu &= -\frac{\langle p_I | (r \cdot \Gamma) \Gamma^\mu | p_J \rangle}{4(p \cdot r)} (\sigma_a)^{IJ} \\ &= -\varepsilon_{IJ}^\mu (\sigma_a)^{IJ} \end{aligned} \quad (4)$$

here I, J are fundamental little group indices, runs from 1 to 2.

And we know that the reasonable polarization vector for physical state should be symmetric with little group indices, it means that we need to symmetrize the indices I and J as $(IJ) = \frac{1}{2}(IJ + JI)$. Because of the symmetry of $(\sigma_a)^{IJ}$, so it can be obtained that

$$\varepsilon_a^\mu = -(\varepsilon_{(IJ)}^\mu + \varepsilon_{[IJ]}^\mu) (\sigma_a)^{IJ} = -\varepsilon_{(IJ)}^\mu (\sigma_a)^{IJ} \quad (5)$$

which means we only need to consider the symmetric part of ε_{IJ}^μ .

Now, we want to compute the normalization for ε_{IJ}^μ , and after multiplying the symmetric matrix, it should go back to the normalization

$$\varepsilon_a^\mu \varepsilon_{b,\mu} = -\delta_{ab} \quad (6)$$

The method to compute the normalization is just multiplying two polarization vectors directly, like

$$\varepsilon_{(IJ)\varepsilon_{(KL),\mu}}^\mu = \frac{\langle p_I | (r \cdot \Gamma) \Gamma^\mu | p_J \rangle}{4(p \cdot r)} \times \frac{\langle p_K | (r \cdot \Gamma) \Gamma_\mu | p_L \rangle}{4(p \cdot r)} \quad (7)$$

here we need to use a special 5d equality

$$(\Gamma^\mu)_A^B (\Gamma_\mu)_C^D = -2\Omega_{AC}\Omega^{BD} + 2\delta_A^D \delta_C^B - \delta_A^B \delta_C^D \quad (8)$$

and the dirac operator can be decomposed to

$$\not{p} = p \cdot \Gamma = p_A^B = |p_I\rangle_A \langle p^I|^B \quad (9)$$

which help us to rewrite the equation (7) to be

$$\begin{aligned} \varepsilon_{IJ\varepsilon_{KL,\mu}}^\mu &= \frac{\langle p_I |^A | r_M \rangle_A \langle r^M |^B (\Gamma^\mu)_B^C | p_J \rangle_C \langle p_K |^D | r_N \rangle_D \langle r^N |^E (\Gamma_\mu)_E^F | p_L \rangle_F}{16(p \cdot r)^2} \\ &= \frac{\langle p_I | r_M \rangle \langle p_K | r_N \rangle}{16(p \cdot r)^2} \times \langle r^M |^B (\Gamma^\mu)_B^C | p_J \rangle_C \langle r^N |^E (\Gamma_\mu)_E^F | p_L \rangle_F \end{aligned} \quad (10)$$

After plugging the equation (8) into the second part of (10), it becomes

$$\begin{aligned} &\langle r^M |^B | p_J \rangle_C \langle r^N |^E | p_L \rangle_F (-2\Omega_{BE}\Omega^{CF} + 2\delta_B^F \delta_E^C - \delta_B^C \delta_E^F) \\ &= -2\langle r^M | r^N \rangle \langle p_L | p_J \rangle + 2\langle r^M | p_L \rangle \langle r^N | p_J \rangle - \langle r^M | p_J \rangle \langle r^N | p_L \rangle \end{aligned} \quad (11)$$

The first term of (11) equals to 0 because the equality

$$\langle p_I | p_J \rangle = 0 \quad (12)$$

and the remaining two term can be contracted with the first part of (10), which helps to obtain the following two terms

$$2\langle p_I | r_M \rangle \langle r^M | p_L \rangle \langle p_K | r_N \rangle \langle r^N | p_J \rangle - \langle p_I | r_M \rangle \langle r^M | p_J \rangle \langle p_K | r_N \rangle \langle r^N | p_L \rangle \quad (13)$$

It is known that

$$\langle p_I | (r \cdot \Gamma) | p_J \rangle = \text{Tr}[|p_J\rangle \langle p_I| (r \cdot \Gamma)] = \frac{1}{2}\epsilon_{IJ}\text{Tr}[(p \cdot \Gamma)(r \cdot \Gamma)] = 2\epsilon_{IJ}(p \cdot r) \quad (14)$$

so the formal equation can be rewritten like

$$8(p \cdot r)^2 \epsilon_{IL} \epsilon_{KJ} - 4(p \cdot r)^2 \epsilon_{IJ} \epsilon_{KL} \quad (15)$$

Finally, we need to symmetrize the indices (IJ) and (KL) , so that we obtain

$$\varepsilon_{IJ\varepsilon_{KL,\mu}}^\mu = \frac{1}{8(p \cdot r)^2} (p \cdot r)^2 (\epsilon_{IL}\epsilon_{KJ} + \epsilon_{JL}\epsilon_{KI} + \epsilon_{IK}\epsilon_{LJ} + \epsilon_{JK}\epsilon_{LI}) - \frac{1}{16(p \cdot r)^2} (p \cdot r)^2 (\epsilon_{(IJ)}\epsilon_{(KL)}) \quad (16)$$

because ϵ itself is an antisymmetric tensor so it will become zero under the symmetrization, so that the second term is vanishing.

After some rearrangement of indices, we can obtain

$$\varepsilon_{IJ}^\mu \varepsilon_{KL,\mu} = \frac{1}{4}(\epsilon_{IL}\epsilon_{KJ} + \epsilon_{IK}\epsilon_{LJ}) \quad (17)$$

Unfortunately, when we want to reproduce the original normalization relation, we will obtain something like

$$\begin{aligned} \varepsilon_a^\mu \varepsilon_{b,\mu} &= (-\varepsilon_{(IJ)}^\mu (\sigma_a)^{IJ})(-\varepsilon_{(KL)}^\mu (\sigma_b)^{KL}) \\ &= \frac{1}{4}(\epsilon_{IL}\epsilon_{KJ} + \epsilon_{IK}\epsilon_{LJ})(\sigma_a)^{IJ}(\sigma_b)^{KL} \\ &= \frac{1}{4}\epsilon_{IL}\epsilon_{KJ}\epsilon^{JM}(\sigma_a)^I{}_M\epsilon^{LN}(\sigma_b)^K{}_N + \frac{1}{4}\epsilon_{IK}\epsilon_{LJ}\epsilon^{IM}(\sigma_a)^J{}_M\epsilon^{LN}(\sigma_b)^K{}_N \\ &= \frac{1}{4}\delta_I^N\delta_K^M(\sigma_a)^I{}_M(\sigma_b)^K{}_N + \frac{1}{4}\delta_K^M\delta_J^N(\sigma_a)^J{}_M(\sigma_b)^K{}_N \\ &= \frac{1}{4}2\text{tr}(\sigma_a\sigma_b) \\ &= \delta_{ab} \end{aligned} \quad (18)$$

2 Completeness Relation of Polarization Vectors

For this purpose, we need to compute the following formula

$$\varepsilon_a^\mu \varepsilon^{a,\nu} = ? \quad (19)$$

If we multiply the momentum p_μ , it should become vanishing so that the completeness relation should look like

$$\varepsilon_a^\mu \varepsilon^{a,\nu} = k(-\eta^{\mu\nu} + \frac{p^\mu r^\nu + p^\nu r^\mu}{(p \cdot r)^2}) \quad (20)$$

here k is just a overall coefficient.

Here, we choose to compute the completeness relation for polarization vectors with fundamental indices

$$\varepsilon_{IJ}^\mu \varepsilon^{IJ,\nu} = a(-\eta^{\mu\nu} + \frac{p^\mu r^\nu + p^\nu r^\mu}{(p \cdot r)^2}) \quad (21)$$

with a different coefficient.

For symplisity, we start with the convention from paper arxiv:2202.08257, in which the polarization vector looks like

$$\epsilon_{IJ}^\mu = \frac{\langle p_I | \Gamma^\mu | r_J \rangle}{\sqrt{2}} \quad (22)$$

with the normalization

$$\langle p^I | r^J \rangle = \epsilon^{IJ} \quad (23)$$

which means the following condition

$$2(p \cdot r) = 1 \quad (24)$$

The method to compute this completeness relation is still just multiplying them together like

$$\varepsilon_{(IJ)}^\mu \varepsilon^{(IJ),\nu} = \frac{\langle p_I | \Gamma^\mu | r_J \rangle}{\sqrt{2}} \times \frac{\langle p^I | \Gamma^\nu | r^J \rangle}{\sqrt{2}} \quad (25)$$

It can be decomposed into 4 pices IJ^IJ , $IJJI$, $JIIJ$, $JIII$, and the first one can be computed like

$$\begin{aligned} \langle p_I | \Gamma^\mu | r_J \rangle \times \langle p^I | \Gamma^\nu | r^J \rangle &= -\langle p_I | \Gamma^\mu | r_J \rangle \langle r^J | \Gamma^\nu | p^I \rangle \\ &= -\text{Tr}[[p^I] \langle p_I | \Gamma^\mu | r_J \rangle \langle r^J | \Gamma^\nu] \\ &= \text{Tr}[(p \cdot \Gamma) \Gamma^\mu (r \cdot \Gamma) \Gamma^\nu] \\ &= 4(p^\mu r^\nu + p^\nu r^\mu - (p \cdot r) \eta^{\mu\nu}) \end{aligned} \quad (26)$$

where we use a nontrivial fact that

$$\langle p^I | \Gamma^\nu | r^J \rangle = -\langle r^J | \Gamma^\nu | p^I \rangle \quad (\text{obtained from } C^T \Gamma^\mu C = -(\Gamma^\mu)^T) \quad (27)$$

The fourth piece is nearly the same as the first one

$$\langle p_J | \Gamma^\mu | r_I \rangle \times \langle p^J | \Gamma^\nu | r^I \rangle = 4(p^\mu r^\nu + p^\nu r^\mu - (p \cdot r) \eta^{\mu\nu}) \quad (28)$$

The remaining two need some extra efforts, here I compute it explicitly

$$\langle p_I | \Gamma^\mu | r_J \rangle \times \langle p^J | \Gamma^\nu | r^I \rangle \quad (29)$$

In terms of two massless spinors that satisfy (23), the $USp(2, 2)$ identity operator can be written as

$$|r_a\rangle_A \langle p^a|^B + |p_a\rangle_A \langle r^a|^B = \delta_A^B \quad (30)$$

It can be used to rewrite the equation (29) like

$$\begin{aligned} &\langle p_I |^A (\Gamma^\mu)_A^B | r_J \rangle_B \langle p^J |^C (\Gamma^\nu)_C^D | r^I \rangle_D \\ &= \langle p_I |^A (\Gamma^\mu)_A^B (\delta_B^C - |p_J\rangle_B \langle r^J|^C) (\Gamma^\nu)_C^D | r^I \rangle_D \\ &= \langle p_I |^A (\Gamma^\mu)_A^B (\Gamma^\nu)_B^D | r^I \rangle_D - \langle p_I | \Gamma^\mu | p_J \rangle \langle r^J | \Gamma^\nu | r^I \rangle \end{aligned} \quad (31)$$

And similarly for

$$\begin{aligned} \langle p_J | \Gamma^\mu | r_I \rangle \times \langle p^I | \Gamma^\nu | r^J \rangle &= \langle p^I | \Gamma^\nu | r^J \rangle \times \langle p_J | \Gamma^\mu | r_I \rangle \\ &= \langle p^I |^A (\Gamma^\nu)_A^B (-|p^J\rangle_B \langle r_J|^C - \delta_B^C) (\Gamma^\mu)_C^D | r_I \rangle_D \\ &= -\langle p^I |^A (\Gamma^\nu)_A^B (\Gamma^\mu)_B^D | r_I \rangle_D - \langle p^I | \Gamma^\nu | p^J \rangle \langle r_J | \Gamma^\mu | r_I \rangle \end{aligned} \quad (32)$$

Now we can combine these two terms

$$\begin{aligned} &\langle p_I |^A (\Gamma^\mu)_A^B (\Gamma^\nu)_B^D | r^I \rangle_D - \langle p_I | \Gamma^\mu | p_J \rangle \langle r^J | \Gamma^\nu | r^I \rangle - \langle p^I |^A (\Gamma^\nu)_A^B (\Gamma^\mu)_B^D | r_I \rangle_D - \langle p^I | \Gamma^\nu | p^J \rangle \langle r_J | \Gamma^\mu | r_I \rangle \\ &= \langle p_I | (\Gamma^\mu \Gamma^\nu + \Gamma^\nu \Gamma^\mu) | r^I \rangle - \langle p_I | \Gamma^\mu | p_J \rangle \langle r^J | \Gamma^\nu | r^I \rangle - \langle p^I | \Gamma^\nu | p^J \rangle \langle r_J | \Gamma^\mu | r_I \rangle \\ &= 2\eta^{\mu\nu} \langle p_I | r^I \rangle - (2p^\mu \epsilon_{IJ})(-2r^\nu \epsilon^{JI}) - (-2p^\nu \epsilon^{IJ})(2r^\mu \epsilon_{JI}) \\ &= 4\eta^{\mu\nu} + 8p^\mu r^\nu + 8p^\nu r^\mu \end{aligned} \quad (33)$$

Above all, we can conclude that

$$\begin{aligned}\varepsilon_{(IJ)}^\mu \varepsilon^{(IJ),\nu} &= \frac{1}{2} \times \frac{1}{8} [8(p^\mu r^\nu + p^\nu r^\mu - (p \cdot r) \eta^{\mu\nu}) + 4\eta^{\mu\nu} + 8p^\mu r^\nu + 8p^\nu r^\mu] \\ &\stackrel{?}{=} 2p^{(\mu} r^{\nu)}\end{aligned}\tag{34}$$

3 Completeness Relation Review

Under the following definition of polarization vector

$$\varepsilon_a^\mu = \frac{\langle p_I | (r \cdot \Gamma) \Gamma^\mu | p_J \rangle}{4(p \cdot r)} (\sigma_a)^I_K \epsilon^{KJ}\tag{35}$$

and same for

$$\varepsilon_a^\mu = \frac{\langle p^I | (r \cdot \Gamma) \Gamma^\mu | p^J \rangle}{4(p \cdot r)} (\sigma_a)^K_J \epsilon_{KI},\tag{36}$$

the polarization vector living in the adjoint representation can be defined as

$$\varepsilon_{IJ}^\mu(p; r) := \frac{\langle p_I | (r \cdot \Gamma) \Gamma^\mu | p_J \rangle}{4(p \cdot r)}\tag{37}$$

$$\varepsilon^{IJ,\mu}(p; r) := \frac{\langle p^I | (r \cdot \Gamma) \Gamma^\mu | p^J \rangle}{4(p \cdot r)}.\tag{38}$$

The corresponding complex conjugate can be obtained

$$(\varepsilon_{IJ}^\mu(p; r))^* = -\frac{\langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle}{4(p \cdot r)}\tag{39}$$

and similarly

$$(\varepsilon^{IJ,\mu}(p; r))^* = -\frac{\langle p_I | \Gamma^\mu(r \cdot \Gamma) | p_J \rangle}{4(p \cdot r)}.\tag{40}$$

For the consistency of index placement with the summation convention, we can define the following quantities

$$\varepsilon_{IJ}^{*\mu}(p; r) := (\varepsilon^{IJ,\mu}(p; r))^* = -\frac{\langle p_I | \Gamma^\mu(r \cdot \Gamma) | p_J \rangle}{4(p \cdot r)},\tag{41}$$

$$\varepsilon^{*IJ,\mu}(p; r) := (\varepsilon_{IJ}^\mu(p; r))^* = -\frac{\langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle}{4(p \cdot r)}\tag{42}$$

Then the completeness relation can be computed like

$$\varepsilon^{*IJ,\mu} \varepsilon_{IJ}^\nu = -\frac{\langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle}{4(p \cdot r)} \frac{\langle p_I | (r \cdot \Gamma) \Gamma^\nu | p_J \rangle}{4(p \cdot r)}\tag{43}$$

It includes 4 terms and can be grouped into two part, the first part can be computed like

$$\begin{aligned}
& \langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle \langle p_I | (r \cdot \Gamma) \Gamma^\nu | p_J \rangle \\
&= -\langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle \langle p_J | \Gamma^\nu(r \cdot \Gamma) | p_I \rangle \\
&= -\text{tr}[\langle p_I | \langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle \langle p_J | \Gamma^\nu(r \cdot \Gamma)] \\
&= \text{tr}[(p \cdot \Gamma) \Gamma^\mu(r \cdot \Gamma) (p \cdot \Gamma) \Gamma^\nu(r \cdot \Gamma)]
\end{aligned} \tag{44}$$

Here we need to compute trace for the product of 6 gamma matrices, and I just give the answer

$$\begin{aligned}
\text{Tr}(\Gamma^a \Gamma^b \Gamma^c \Gamma^d \Gamma^e \Gamma^f) = 4 \big(& \eta^{ab} \eta^{cd} \eta^{ef} - \eta^{ab} \eta^{ce} \eta^{df} + \eta^{ab} \eta^{cf} \eta^{de} \\
& - \eta^{ac} \eta^{bd} \eta^{ef} + \eta^{ac} \eta^{be} \eta^{df} - \eta^{ac} \eta^{bf} \eta^{de} \\
& + \eta^{ad} \eta^{bc} \eta^{ef} - \eta^{ad} \eta^{be} \eta^{cf} + \eta^{ad} \eta^{bf} \eta^{ce} \\
& - \eta^{ae} \eta^{bc} \eta^{df} + \eta^{ae} \eta^{bd} \eta^{cf} - \eta^{ae} \eta^{bf} \eta^{cd} \\
& + \eta^{af} \eta^{bc} \eta^{de} - \eta^{af} \eta^{bd} \eta^{ce} + \eta^{af} \eta^{be} \eta^{cd} \big).
\end{aligned} \tag{45}$$

A compact way to write the result is

$$\text{Tr}(\Gamma^{a_1} \dots \Gamma^{a_6}) = 4 \sum_{\text{pairings}} (-1)^\sigma \eta^{a_{i_1} a_{j_1}} \eta^{a_{i_2} a_{j_2}} \eta^{a_{i_3} a_{j_3}}.$$

Here the sum runs over all pairwise partitions (pairings) of the index set $\{1, \dots, 6\}$ into three disjoint pairs $\{(i_1, j_1), (i_2, j_2), (i_3, j_3)\}$. Here $(-1)^\sigma$ denotes the sign (parity) of the permutation needed to reorder the sequence so that the members of each pair become adjacent.

Using this formula, we can rewrite the equation (44)

$$\begin{aligned}
(44) = & p^\mu r^\nu (p \cdot r) - p^\mu r^\nu (p \cdot r) + p^\mu p^\nu r^2 \\
& - p^\mu r^\nu (p \cdot r) + \eta^{\mu\nu} (p \cdot r)^2 - p^\nu r^\mu (p \cdot r) \\
& + r^\mu r^\nu p^2 - \eta^{\mu\nu} p^2 r^2 + r^\mu r^\nu p^2 \\
& - p^\nu r^\mu (p \cdot r) + p^\nu p^\mu r^2 - p^\nu r^\mu (p \cdot r) \\
& + p^\nu r^\mu (p \cdot r) - p^\mu r^\nu (p \cdot r) + \eta^{\mu\nu} (p \cdot r)^2 \\
& = 4(-2p^\mu r^\nu (p \cdot r) - 2p^\nu r^\mu (p \cdot r) + \eta^{\mu\nu} (p \cdot r)^2)
\end{aligned} \tag{46}$$

in which many terms cancel with each other.

Similarly, the second part can be computed like

$$\begin{aligned}
& \langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle \langle p_J | (r \cdot \Gamma) \Gamma^\nu | p_I \rangle \\
&= \text{tr}[\langle p_I | \langle p^I | \Gamma^\mu(r \cdot \Gamma) | p^J \rangle \langle p_J | (r \cdot \Gamma) \Gamma^\nu] \\
&= -\text{tr}[(p \cdot \Gamma) \Gamma^\mu(r \cdot \Gamma) (p \cdot \Gamma) (r \cdot \Gamma) \Gamma^\nu] \\
&= -\text{tr}[(p \cdot \Gamma) \Gamma^\mu(r \cdot \Gamma) (p \cdot \Gamma) \{(r \cdot \Gamma), \Gamma^\nu\}] + \text{tr}[(p \cdot \Gamma) \Gamma^\mu(r \cdot \Gamma) (p \cdot \Gamma) \Gamma^\nu(r \cdot \Gamma)]
\end{aligned} \tag{47}$$

It can be noticed that the first term equals to 0 and the second term is just the same as the result for first part. In short, the two part give us the same result.

Above all, the completeness relation can be written as

$$\begin{aligned}
\varepsilon^{*IJ,\mu} \varepsilon_{IJ}^\nu &= -\frac{1}{16(p \cdot r)^2} \times \frac{1}{4} \times [4 \times 4(-2p^\mu r^\nu(p \cdot r) - 2p^\nu r^\mu(p \cdot r) + \eta^{\mu\nu}(p \cdot r)^2)] \\
&= -\frac{1}{2} \left[\frac{-p^\mu r^\nu - p^\nu r^\mu}{(p \cdot r)} + \eta^{\mu\nu} \right] \\
&= -\frac{1}{2} \eta^{\mu\nu} + \frac{p^{(\mu} r^{\nu)}}{(p \cdot r)}
\end{aligned} \tag{48}$$

We can quickly check its correctness by multiplying a momentum

$$p_\mu \varepsilon^{*IJ,\mu} \varepsilon_{IJ}^\nu = -\frac{1}{2} p^\nu + \frac{\frac{1}{2} p^2 r^\nu + \frac{1}{2} p^\nu (p \cdot r)}{(p \cdot r)} = 0 \tag{49}$$

4 All for 5 dimension

4.1 Spinor-Helicity in 5D

The Lorentz group in 5D is $\text{Spin}(5) \cong \text{USp}(4) \cong \text{Sp}(4, \mathbb{C}) \cap \text{SU}(4)$, we can also consider the non-compact version $\text{SO}(1, 4)$, which has an isomorphism $\text{SO}(1, 4) \cong \text{USp}(2, 2)$. For a 5-momentum vector p^μ , we can identify it with a 4×4 antisymmetric matrix

$$p_{AB} = p_\mu \gamma_{AB}^\mu \equiv -p_\mu (\gamma^\mu)_A^C \Omega_{CB} = -(\not{p} \Omega)_{AB} \tag{50}$$

Here we choose a symplectic form to be the metric

$$\Omega_{AB} = \begin{pmatrix} \epsilon_{\alpha\beta} & 0 \\ 0 & -\epsilon^{\dot{\alpha}\dot{\beta}} \end{pmatrix} = \Omega^{BA} \tag{51}$$

where we use the fact that fundamental indices A, B for $\text{USp}(2, 2)$ can be decomposed to $\text{SL}(2, \mathbb{C})$ indices like $A = \alpha \oplus \dot{\alpha}$, and the normalization for 2 dimension Levi-Civita tensor is $\epsilon^{12} = \epsilon_{21} = 1$.

We use metric Ω to lower and raise the indices like

$$X^A = \Omega^{AB} X_B, \quad X_A = \Omega_{AB} X^B \tag{52}$$

where X are spinors belongs to $\text{USp}(2, 2)$.

Also, from the definition (50), we know the gamma matrices with lower indices can be defined like

$$\gamma_{AB} = \Omega_{BC} \gamma_A^C \tag{53}$$

similary the gamma matrix with upper indices can be obtained by

$$\gamma^{AB} = \Omega^{AC} \gamma_C^B. \tag{54}$$

The gamma matrices with lower indices are antisymmetric and Ω -traceless

$$\gamma_{AB} = \gamma_{BA}, \quad \gamma_{AB} \Omega^{AB} = 0 \tag{55}$$

The determinant of p_{AB} evaluates to $\text{Det}(p_{AB}) = \frac{1}{4}(p_{AB}p^{AB})^2 = (p^2)^2$. For 5D massless particle, $p^2 = 0 \Rightarrow \text{Det}(p_{AB}) = 0$, so the rank of this matrix equals to 2. Because the little group of 5D massless particle is $\text{SO}(3) \cong \text{SU}(2)$, so the matrix p_{AB} can be decomposed to be the outer product of two 4×2 matrix, here specifically, the $\text{USp}(2, 2)$ spinors with $\text{SU}(2)$ little group indices

$$p_{AB} = \lambda_A^a \lambda_{Ba} = |p^a\rangle_A |p_a\rangle_B, \quad p_A^B = \lambda_A^a \lambda_a^B = |p^a\rangle_A \langle p_a|^B \quad (56)$$

with a, b runs over 1, 2. A little group transformation that preserves p_{AB} behaves like $|p^a\rangle_A \rightarrow t^a_b |p^b\rangle_A$, here t^a_b refers to the $\text{SU}(2)$ little group transformation.

Since the little groups of massive 4D momentum and massless 5D momentum are identified, there exist a consistent embedding of the former into the latter.

$$\lambda_A^a = \begin{pmatrix} \lambda_\alpha^a \\ \tilde{\lambda}^{\dot{\alpha}a} \end{pmatrix}, \quad \Omega_{AB} = \begin{pmatrix} -\epsilon_{\alpha\beta} & 0 \\ 0 & \epsilon^{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \quad (57)$$

where we can think of 5D Lorentz indices breaking up into 4D Lorentz indices $A \mapsto (\alpha, \dot{\alpha})$ once a particular direction is chosen, means we manifestly break the 5D Lorentz symmetry.

The antisymmetry of p_{AB} implies the important identity that

$$\Omega^{AB} \lambda_A^a \lambda_B^b = 0, \quad (58)$$

so no Lorentz-invariant little-group tensors can be formed from just considering a single particle alone.

4.2 5D 3-particle Amplitude

Special Kinematics in 5D

Scattering amplitudes, are Lorentz invariant and transform covariantly under the little group. So like the massive 4D case, the amplitudes here are $\text{SU}(2)$ covariants, means it carries $\text{SU}(2)$ little group indices and no free Lorentz indices. We want keep the consistency with 4D so here the spin- s particle still transformed in the $(2s+1)$ -dimension symmetric representation of $2s$ product of $\text{SU}(2)$. In short, a spin- s particle should carry $2s$ $\text{SU}(2)$ little group indices.

Then, a 3-particle amplitude of spin s_1, s_2, s_3 is a tensor carrying $(2s_1 + 2s_2 + 2s_3)$ indices like

$$\mathcal{M}_{(a_1 \dots a_{2s_1})(b_1 \dots b_{2s_2})(c_1 \dots c_{2s_3})} \quad (59)$$

for a_i, b_j, c_k indices of $\text{SU}(2)$ little group of particle 1, 2, 3 respectively.

In order to construct amplitudes which is a tensor carrying little group indices, it is necessary to figure out what tensors structure are possible here. The spinors that make up an on-shell 3-point amplitude without any Lorentz indices can be possibly listed as follows

$$y^{12}_{ab} := (\lambda_1)^A_a (\lambda_2)_{Ab} \quad (60)$$

$$y^{23}_{bc} := (\lambda_2)^A_b (\lambda_3)_{Ac} \quad (61)$$

$$y^{31}_{ca} := (\lambda_3)^A_c (\lambda_1)_{Aa} \quad (62)$$

Because of momentum conservation, the determinant of this y^{ij} vanishes

$$\text{Det}(y^{12}) = \frac{1}{2}\epsilon^{ab}\epsilon^{cd}y^{12}_{ac}y^{12}_{bd} = \frac{1}{2}y^{12}_{ac}y^{12ac} = \frac{1}{2}p_1^{AB}p_{2AB} = 2(p_1 \cdot p_2) = 0 \quad (63)$$

This means we can decompose each of them into a outer product of SU(2) spinors:

$$y^{12}_{ab} = y^1_a y^2_b \quad (64)$$

$$y^{23}_{bc} = y^2_b y^3_c \quad (65)$$

$$y^{31}_{ca} = y^3_c y^1_a \quad (66)$$

In the following, we use the notation: $1_a := y^1_a$ and $1^A_a := (\lambda_1)^A_a$ for massless 5D spinors.

First, we can try to write down a 3-point amplitude using these variables, e.g. the Yukawa interaction in 5D is precisely $y^1 2_{ab}$:

$$\mathcal{M}_{a,b} = \text{Corresponding Feynman diagram} = g 1_a 2_b \quad (67)$$

Connecting to 4D x-factor

If there exist a similar "x" factor in 5D, it would be of the form $\chi^c_{c'}$ with no Lorentz indices but little group indices. It can be understood as

$$p_2^{AB} 3_B^c = \chi^c_{c'} 3^{Ac'} \quad (68)$$

There is no inverse of χ because of $\chi^2 = 0$, $\text{tr}\chi = 0$ and $\text{Det}\chi = 0$. In the following, it can be known that

$$\chi^c_{c'} = 3^c 3_{c'} \quad (69)$$

so the special kinematics SU(2) spinors $y^3_c = 3_c$ is a sort of "square root" of χ , the x-factor in 5D. Similarly, for other two particles, we can obtain the corresponding x-factor

$$p_3^{AB} 1_B^a = \chi^a_{a'} 1^{Aa'} = (1_a 1^{a'}) 1^{Aa'} \quad (70)$$

$$p_1^{AB} 2_B^b = \chi^b_{b'} 2^{Ab'} = (2_b 2^{b'}) 2^{Ab'} \quad (71)$$

Then, in scalar-QED in 5D, the scalar-scalar-photon amplitude can be directly written as

$$\mathcal{M}_{cc'} = (\text{Corresponding Feynman diagram}) = g 3_c 3_{c'} \quad (72)$$

If one compactifies to 4D, all 4D massive amplitudes should satisfy $m_1 \pm m_2 \pm' m_3 = 0$. If one particle 3 is massless in 4D and other 2 are massive with same mass m , the 4D x factor appears in the form as

$$3_c 3_{c'} = \begin{pmatrix} -mx^{-1} & \mp im \\ \mp im & mx \end{pmatrix}_{cc'} \quad (73)$$

So we obtain the corresponding amplitude after compactifying, where the diagonal components refer to the massless 4-vector of particle 3, and the off-diagonal components represent the scalar sector. Either $\mathcal{M} \sim x$ or x^{-1} depending on its helicity.

5 Detailed Construction

We start from the convention for 5D gamma matrix and momentum

$$p^\mu = (p_0, \vec{p}, p^4), \quad \mu = \bar{\mu} = (0, 1, 2, 3) \quad \mu = \hat{\mu} = 4 \quad (74)$$

$$p_\mu = (p^0, -\vec{p}, -p^4) \quad (75)$$

and

$$\Gamma^{\bar{\mu}} = \begin{pmatrix} 0 & \sigma^{\bar{\mu}} \\ \bar{\sigma}^{\bar{\mu}} & 0 \end{pmatrix} \quad (76)$$

$$\Gamma^4 = \begin{pmatrix} -i & & & \\ & -i & & \\ & & i & \\ & & & i \end{pmatrix} \quad (77)$$

where $\sigma^{\bar{\mu}} = (\mathbf{1}, \vec{\sigma})$ and $\bar{\sigma}^{\bar{\mu}} = (\mathbf{1}, -\vec{\sigma})$.

Then the Dirac operator can be written as

$$\begin{aligned} \not{p} &= p_\mu \Gamma^\mu = p_{\bar{\mu}} \Gamma^{\bar{\mu}} - p^4 \Gamma^4 \\ &= \begin{pmatrix} 0 & 0 & p^0 - p^3 & -p^1 + ip^2 \\ 0 & 0 & -p^1 - i - p^2 & p^0 + p^3 \\ p^0 + p^3 & p^1 - ip^2 & 0 & 0 \\ p^1 + ip^2 & p^0 - p^3 & 0 & 0 \end{pmatrix} - \begin{pmatrix} -ip^4 & & & \\ & -ip^4 & & \\ & & ip^4 & \\ & & & ip^4 \end{pmatrix} \\ &= \begin{pmatrix} ip^4 & 0 & p^0 - p^3 & -p^1 + ip^2 \\ 0 & ip^4 & -p^1 - i - p^2 & p^0 + p^3 \\ p^0 + p^3 & p^1 - ip^2 & -ip^4 & 0 \\ p^1 + ip^2 & p^0 - p^3 & 0 & -ip^4 \end{pmatrix} \end{aligned} \quad (78)$$

The determinant of this matrix equals to zero, which means it can be decomposed to the outer product of two 4-component spinor with $\text{Sp}(2, \mathbb{C})$ little group indices

$$\not{p} = |p^a\rangle_A \langle p_a|^B \quad (79)$$

and they satisfy the massless Dirac equation

$$\not{p} |p^a\rangle_A = 0. \quad (80)$$

A specific construction for the spinor-helicity variable is

$$|p^a\rangle_A = \begin{pmatrix} p^0 - p^3 & 0 \\ -p^1 - ip^2 & \frac{ip^4}{p^0 - p^3} \\ -ip^4 & \frac{p^1 - ip^2}{p^0 - p^3} \\ 0 & 1 \end{pmatrix} \quad (81)$$

and the bra can be obtained by introducing a symplectic form

$$\Omega_{AB} = \begin{pmatrix} 0 & -1 & & \\ 1 & 0 & & \\ & & 0 & -1 \\ & & 1 & 0 \end{pmatrix} = \Omega^{BA}. \quad (82)$$

Then the corresponding bra variable can be written like

$$\begin{aligned} \langle p_a |^A &= \epsilon_{ab} (\Omega^{AB} |p^b\rangle_B)^T \\ &= \begin{pmatrix} \frac{-ip^4}{p^0-p^3} & 0 & -1 & \frac{p^1-ip^2}{p^0-p^3} \\ -p^1-ip^2 & -p^0+p^3 & 0 & p^4 \end{pmatrix} \end{aligned} \quad (83)$$

Also, we have

$$\begin{aligned} |p_a\rangle_A &= \epsilon_{ab} |p^b\rangle_A \\ &= \begin{pmatrix} p^0-p^3 & 0 \\ -p^1-ip^2 & \frac{ip^4}{p^0-p^3} \\ -ip^4 & \frac{p^1-ip^2}{p^0-p^3} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & p^0-p^3 \\ \frac{-ip^4}{p^0-p^3} & -p^1-ip^2 \\ \frac{-p^1+ip^2}{p^0-p^3} & -ip^4 \\ -1 & 0 \end{pmatrix} \end{aligned} \quad (84)$$

Similar for 4d massless case, we can also construct the corresponding spinor-helicity variable from

$$p_{\alpha\dot{\alpha}} = \begin{pmatrix} p^0-p^3 & -p^1+ip^2 \\ -p^1-ip^2 & p^0+p^3 \end{pmatrix} = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}} \quad (85)$$

where the λ and $\tilde{\lambda}$ are represented by

$$\lambda_\alpha = |p\rangle_\alpha = \begin{pmatrix} p^0-p^3 \\ -p^1-ip^2 \end{pmatrix}, \quad \tilde{\lambda}_{\dot{\alpha}} = [p]_{\dot{\alpha}} = \begin{pmatrix} 1 \\ \frac{-p^1+ip^2}{p^0-p^3} \end{pmatrix} \quad (86)$$

Also, we can obtain the corresponding "bra" and "ket" variable by contracting with antisymmetric tensor ϵ

$$\langle p |^\alpha = \epsilon^{\alpha\beta} |p\rangle_\beta = \begin{pmatrix} -p^1-ip^2 \\ -p^0+p^3 \end{pmatrix}, \quad |p]^{\dot{\alpha}} = \epsilon^{\dot{\alpha}\dot{\beta}} [p]_{\dot{\beta}} = \begin{pmatrix} \frac{-p^1+ip^2}{p^0-p^3} \\ -1 \end{pmatrix} \quad (87)$$

Then it can found that if we choose the limit in which the fifth component $p^4 = 0$, the 5D spinor-helicity variable reduces to

$$\langle p_a |^A |_{p^4=0} = \begin{pmatrix} 0 & \langle p |^\alpha \\ -[p]_{\dot{\alpha}} & 0 \end{pmatrix}^T \quad (88)$$

and

$$|p_a\rangle_A|_{p^4=0} = \begin{pmatrix} 0 & |p\rangle_\alpha \\ |p]_{\dot{\alpha}} & 0 \end{pmatrix} \quad (89)$$

It is straightforward now to compute y^{ij}_{ab} , here we choose y^{12}_{ab} as an example

$$\langle 1_a | 2_b \rangle = \begin{pmatrix} [21] & 0 \\ 0 & \langle 12 \rangle \end{pmatrix} = 1_a 2_b \quad (90)$$

From the 3 particle momentum conservation, we have known that 4D massless spinor-helicity variable should satisfy

$$|1\rangle \propto |2\rangle \propto |3\rangle \quad \text{or} \quad |1] \propto |2] \propto |3] \quad (91)$$

which means one of $[21]$ and $\langle 12 \rangle$ will become vanishing.

First we consider non-vanishing $[21]$ case, in which we can choose a particular decomposition

$$y^{12} = 1_a 2_b = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} [21] \\ 0 \end{pmatrix} \quad (92)$$

$$y^{23} = \Upsilon_{2b} 3_c = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} [32] \\ 0 \end{pmatrix} \quad (93)$$

$$y^{31} = \Upsilon_{3c} \Upsilon_{1a} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} [13] \\ 0 \end{pmatrix} \quad (94)$$

from which we can read off

$$\Upsilon_{1a} = [13] 1_a, \quad \Upsilon_{2b} = \frac{1}{[21]} 2_b, \quad \Upsilon_{3c} = \frac{1}{[32]} 3_c \quad (95)$$

We denote these coefficients as A , B and C . We want to use the consistent decomposition for these y variables, means the same 2_b appearing both in y^{12} and y^{23} , like

$$y^{12} = 1'_a 2'_b \quad (96)$$

$$y^{23} = 2'_b 3'_c \quad (97)$$

$$y^{31} = 3'_c 1'_a \quad (98)$$

Because of the ambiguity in the decomposition, we can rescale like

$$1'_a = \alpha_1 1_a, \quad 2'_b = \alpha_2 2_b, \quad 3'_c = \alpha_3 3_c \quad (99)$$

which means the following equation should satisfy

$$y^{12} = \alpha_1 1_a \alpha_2 2_b = 1_a 2_b \quad (100)$$

$$y^{23} = \alpha_2 2_b \alpha_3 3_c = 2_b 3_c \quad (101)$$

$$y^{31} = \alpha_3 3_c \alpha_1 1_a = 3_c 1_a \quad (102)$$

So we have a group of equations

$$\alpha_1 \alpha_2 = 1 \quad (103)$$

$$\alpha_2 \alpha_3 = BC \quad (104)$$

$$\alpha_3 \alpha_1 = CA \quad (105)$$

and the solutions are given as following

$$\alpha_1 = \sqrt{\frac{AC}{B}}, \quad \alpha_2 = \sqrt{\frac{B}{AC}}, \quad \alpha_3 = \sqrt{ABC} \quad (106)$$

After inserting these solutions, we obtain the final decomposition

$$1_a = \sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad 2_b = \sqrt{\frac{[21][32]}{[13]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad 3_c = \sqrt{\frac{[13][32]}{[21]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (107)$$

In order to give the correct Yang-Mills amplitude, we need the "inverse" of these two-component spinor, defined as following

$$\hat{i}^a i_a = 1. \quad (108)$$

From (107), we can explicitly solve the inverse spinor

$$\hat{1}^a = \sqrt{\frac{[32]}{[13][21]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \alpha_1 1^a, \quad (109)$$

$$\hat{2}^b = \sqrt{\frac{[13]}{[21][32]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \alpha_2 2^b, \quad (110)$$

$$\hat{3}^c = \sqrt{\frac{[21]}{[13][32]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \alpha_3 3^c \quad (111)$$

here the second means the shift ambiguity. (α are arbitrary complex number here)

Because of the physical constraint

$$\Lambda \cdot z = 0 \quad (112)$$

we need

$$\hat{\delta}_{1A} + \hat{\delta}_{2A} + \hat{\delta}_{3A} + (\alpha_1 + \alpha_2 + \alpha_3)\delta_A = 0 \quad (113)$$

where $\hat{\delta}_{iA} = \hat{i}^a i_{Aa}$ and $\delta_A = i^a i_{Aa}$.

We also know that

$$\hat{\delta}_{1A} + \hat{\delta}_{2A} + \hat{\delta}_{3A} = \hat{K}\delta_A \quad (114)$$

so in order to satisfy the constraint, we just need

$$\hat{K} + (\alpha_1 + \alpha_2 + \alpha_3) = 0 \quad (115)$$

Here, we can try to give the explicit form of $\hat{\delta}_{1A}$

$$\hat{\delta}_{1A} = \hat{1}^a 1_{Aa} = \begin{pmatrix} 0 & p^0 - p^3 \\ 0 & -p^1 - ip^2 \\ \frac{-p^1 + ip^2}{p^0 - p^3} & 0 \\ -1 & 0 \end{pmatrix} \sqrt{\frac{[32]}{[13][21]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \sqrt{\frac{[32]}{[13][21]}} \begin{pmatrix} 0 \\ 0 \\ \frac{-p^1 + ip^2}{p^0 - p^3} \\ -1 \end{pmatrix} = \sqrt{\frac{[32]}{[13][21]}} \begin{pmatrix} 0 \\ |1]^{\dot{\alpha}} \end{pmatrix} \quad (116)$$

The non-vanishing component can be written as

$$\sqrt{\frac{[32]}{[13][21]}}|1]^{\dot{\alpha}} = \sqrt{\frac{1}{[32][21][13]}}[32]|1]^{\dot{\alpha}} \quad (117)$$

From the Schouten identity

$$|i]\langle jk\rangle + |k]\langle ij\rangle + |j]\langle ki\rangle = 0 \quad (118)$$

we can obtain

$$[32]|1] = -([13]|2] + [21]|3]). \quad (119)$$

Surprisingly, we can find that

$$\sqrt{\frac{1}{[32][21][13]}}[13]|2] \quad \text{and} \quad \sqrt{\frac{1}{[32][21][13]}}[21]|3] \quad (120)$$

are just the non-vanishing components of $\hat{\delta}_{2A}$ and $\hat{\delta}_{3A}$, in other words,

$$\hat{\delta}_{1A} = -\hat{\delta}_{2A} - \hat{\delta}_{3A} \quad (121)$$

so that

$$\hat{\delta}_{1A} + \hat{\delta}_{2A} + \hat{\delta}_{3A} = 0 \quad (122)$$

which means the constraint is equivalent to

$$(\alpha_1 + \alpha_2 + \alpha_3)\delta_A = 0 \quad \Leftrightarrow \quad \alpha_1 + \alpha_2 + \alpha_3 = 0 \quad (123)$$

Here, we can simply choose them to be 0. Also, we can confirm $\delta_A = 1^a 1_{Aa} = 2^b 2_{Ab} = 3^c 3_{Ac}$ through these explicit forms

$$1^a 1_{Aa} = \begin{pmatrix} 0 & p^0 - p^3 \\ 0 & -p^1 - ip^2 \\ \frac{-p^1 + ip^2}{p^0 - p^3} & 0 \\ -1 & 0 \end{pmatrix} \sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} 0 \\ -1 \end{pmatrix} = \sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} -(p^0 - p^3) \\ -(-p^1 - ip^2) \\ 0 \\ 0 \end{pmatrix} = \sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} -|1\rangle_{\alpha} \\ 0 \end{pmatrix} \quad (124)$$

From the momentum conservation

$$|1\rangle[1] + |2\rangle[2] + |3\rangle[3] = 0 \quad (125)$$

by contracting a $|3]$ in both sides, we can obtain

$$|1\rangle = -\frac{[23]}{[13]}|2\rangle \quad (126)$$

and this gives us the following result

$$\sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} \frac{[23]}{[13]}|2\rangle_{\alpha} \\ 0 \end{pmatrix} = \sqrt{\frac{[21][32]}{[13]}} \begin{pmatrix} -|2\rangle_{\alpha} \\ 0 \end{pmatrix} = 2^b 2_{Ab} \quad (127)$$

So we confirm the fact

$$\delta_A = 1^a 1_{Aa} = 2^b 2_{Ab} \quad (128)$$

the remaining two can be confirmed in similar manner.

Suppose we want to compute the amplitude with two spin $\frac{1}{2}$ fermions and spin 1 gauge boson in 5D, the amplitude can be directly written as

$$\mathcal{M}_{ab(cc')} = g(1_a 2_b \hat{3}_c + 1_a \hat{2}_b 3_c + \hat{1}_a 2_b 3_c) 3_{c'}. \quad (129)$$

By using the explicit form of these building blocks, it can be written as

$$\begin{aligned} \mathcal{M}_{ab(cc')} = & \left(\sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_a \sqrt{\frac{[21][32]}{[13]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_b \sqrt{\frac{[21]}{[13][32]}} \begin{pmatrix} 0 \\ 1 \end{pmatrix}_c + \right. \\ & \sqrt{\frac{[13][21]}{[32]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_a \sqrt{\frac{[13]}{[21][32]}} \begin{pmatrix} 0 \\ 1 \end{pmatrix}_b \sqrt{\frac{[13][32]}{[21]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_c + \\ & \left. \sqrt{\frac{[32]}{[13][21]}} \begin{pmatrix} 0 \\ 1 \end{pmatrix}_a \sqrt{\frac{[21][32]}{[13]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_b \sqrt{\frac{[13][32]}{[21]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_c \right) \sqrt{\frac{[13][32]}{[21]}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_{c'} \quad (130) \end{aligned}$$

By picking the different indices, we can reproduce the different helicity amplitudes in 4D, like

$$a, b, (c, c') = 1, 2, (1, 1) \Rightarrow \sqrt{\frac{[13][21]}{[32]}} \sqrt{\frac{[13]}{[21][32]}} \frac{[13][32]}{[21]} = \frac{[13]^2}{[21]} \quad (131)$$

$$a, b, (c, c') = 1, 1, (1, 2) = \sqrt{\frac{[13][21]}{[32]}} \sqrt{\frac{[32][21]}{[13]}} \frac{1}{2}(1+1) = [21] \quad (132)$$

$$a, b, (c, c') = 2, 1, (1, 1) = \sqrt{\frac{[32]}{[13][21]}} \sqrt{\frac{[21][32]}{[13]}} \frac{[13][32]}{[21]} = \frac{[32]^2}{[21]} \quad (133)$$

From the known 4D results

$$M^{h_1 h_2 h_3} = \begin{cases} \tilde{g} [12]^{h_1+h_2-h_3} [23]^{h_2+h_3-h_1} [31]^{h_3+h_1-h_2}, & \text{when } h_1 + h_2 + h_3 > 0, \\ g \langle 12 \rangle^{h_3-h_1-h_2} \langle 23 \rangle^{h_1-h_2-h_3} \langle 31 \rangle^{h_2-h_3-h_1}, & \text{when } h_1 + h_2 + h_3 < 0. \end{cases} \quad (134)$$

we know that the 3 point amplitudes with $(1/2^+, 1/2^-, +1)$ configuration equals to

$$\mathcal{M}(+\frac{1}{2}, -\frac{1}{2}, +1) = \frac{[31]^2}{[12]} \quad (135)$$

$$\mathcal{M}(-\frac{1}{2}, +\frac{1}{2}, +1) = \frac{[32]^2}{[12]} \quad (136)$$

$$\mathcal{M}(0, 0, +1) = [12] \quad (137)$$

We found that the 5D 3 point amplitudes actually reproduce the corresponding 4D amplitudes.

A Detailed explanation of special 5D spinor variables

Lemma : *Momentum consevation for 3-particle amplitude implies*

$$y^{12}_{ab} := 1^A_a 2_{Ab} = y^1_a y^2_b = 1_a 2_b \quad (138)$$

$$y^{23}_{bc} := 2^A_b 3_{Ac} = y^2_b y^3_c = 2_b 3_c \quad (139)$$

$$y^{31}_{ca} := 3^A_c 1_{Aa} = y^3_c y^1_a = 3_c 1_a \quad (140)$$

Proof. First note that $y^{(i)(i)}_{ab} = 0$, e.g.

$$\Omega_{AB} 1^A_a 2^B_b = C \epsilon_{ab} \text{(from antisymmetry of ab)} \quad \text{and} \quad C \propto \Omega_{AB} 1^A_a 1^B_b \epsilon^{ab} \propto \Omega_{AB} p_1^{AB} = 0 \quad (141)$$

Another property of y is that the determinant of this y^{ij} vanishes such that

$$\text{Det}(y^{12}) = \frac{1}{2} \epsilon^{ab} \epsilon^{cd} y^{12}_{ac} y^{12}_{bd} = \frac{1}{2} y^{12}_{ac} y^{12}_{ac} = \frac{1}{2} p_1^{AB} p_{2AB} = 2(p_1 \cdot p_2) = 0 \quad (142)$$

Thus it is a rank 1 quantity, which can be decomposed into the outer product of 2-spinors:

$$y^{12}_{ab} = y^1_a y^2_b \quad (143)$$

$$y^{23}_{bc} = \Upsilon^2_b \Upsilon^3_c \quad (144)$$

$$y^{31}_{ca} = \Upsilon^3_c \Upsilon^1_a \quad (145)$$

Note that not all of there $\text{SU}(2)$ spinors are independent, indeed it is only a 2-dimension vector space. By momentum conservation

$$y^{12}_a y^{23}_b y^{31}_c = 1^A_a 2^B_b 3^C_c = 1^A_a p_{aA}^B 3_B^C = 1^A_a (-p_{1A}^B - p_{3A}^B) 3_B^C = 0 \quad (146)$$

Thus $\Upsilon^{2b} y^2_b = 0$, so $\Upsilon^2_b \propto y^2_b$ (note the metric we are using now is antisymmetric tensor ϵ_{ab}). Similarly,

$$y^{23}_b y^{31}_c y^{12}_a = 0 \Rightarrow \Upsilon^3_c \propto y^3_c \quad (147)$$

$$y^{31}_c y^{12}_a y^{23}_b = 0 \Rightarrow \Upsilon^1_a \propto y^1_a \quad (148)$$

Therefore, up to some constants A,B,C, we have

$$y^{12}_{ab} = y^1_a y^2_b \quad (149)$$

$$y^{23}_{bc} = B y^2_b y^3_c \quad (150)$$

$$y^{31}_{ca} = C y^3_c y^1_a \quad (151)$$

As long as we are working over $\mathbb{C}$, we can always rescale them to obtain

$$y^{12}_{ab} = y^1_a y^2_b \quad (152)$$

$$y^{23}_{bc} = y^2_b y^3_c \quad (153)$$

$$y^{31}_{ca} = y^3_c y^1_a \quad (154)$$

here each y are determined uniquely, up to an overall flip $(y^1, y^2, y^3) \mapsto (-y^1, -y^2, -y^3)$.

The space of 2-dimension representation of $SU(2)$ is too small to ensure the hold of following equality

$$\delta_A := 1^a 1_{Aa} = 2^b 2_{Ab} = 3^c 3_{Ac} \quad (155)$$

δ_A is same for all particles.

Proof. From the 3-particle momentum conservation

$$0 = 1_A^a 1_{Ba} + 2_A^b 2_{Bb} + 3_A^c 3_{Bc} \quad (156)$$

$$0 = \underbrace{1_A^a (1_A^{a'} 1_{Ba'} + 2_A^b 2_{Bb} + 3_A^c 3_{Bc})}_{=0} \quad (157)$$

$$0 = (1_a 2^b) 2_{Bb} + (-1_a 3^c) 3_{Bc} \quad (158)$$

$$0 = 1_a (2^b 2_{Bb} - 3^c 3_{Bc}) \quad (159)$$

$$\Rightarrow 2^b 2_{Bb} = 3^c 3_{Bc} \quad (160)$$

Cyclically premutating the indices, we can obtain $1^a 1_{Ba} = 2^b 2_{Bb} = 3^c 3_{Bc}$.

A useful equality can be obtained

$$\delta_A \lambda(i)^A_b \propto y^{ia} y^i_b = 0 \quad (161)$$

B χ in 5D

Since $\delta_A = 1^a 1_{Aa} = 2^b 2_{Ab} = 3^c 3_{Ac}$, we can evaluate χ_3 by

$$p_2^{AB} 3_B^c = 2^{Ab} 2_B^B 3_B^c = 2^{Ab} 2_b 3^c = -\delta^A 3^c = 3^{Ac'} 3_{c'} 3^c = \underbrace{(3^c 3_{c'})}_{\chi_3} 3^{Ac'} \quad (162)$$

Cycling the particle labels, we can obtain the χ for each particle

$$p_1^{AB} 2_B^b = (2^b 2_{b'}) 2^{Ab'} \quad (163)$$

$$p_3^{AB} 1_B^a = (1^a 1_{a'}) 1^{Aa'} \quad (164)$$

Other combinations can be simply obtained by momentum conservation, like $p_1^{AB} 3_B^c = -p_2^{AB} 3_B^c = -\chi_{3c'}^c 3^{Ac'}$.