

**Problem 1**

Uniform distribution. A random variable that is equally likely to take any value in a finite set  $S$  is said to have the uniform distribution on  $S$ . If  $U$  is such a random variable and  $\emptyset \neq R \subseteq S$ , show that the distribution of  $U$  conditional on  $\{U \in R\}$  is uniform on  $R$ .

*Proof.* Since  $U$  is uniformly distributed on set  $S$ , for any  $s \in S$ , we have:

$$P(U = s) = \frac{1}{|S|}$$

For any element  $r \in R$ , we need to compute the conditional probability  $P(U = r \mid U \in R)$ .

By the definition of conditional probability:

$$P(U = r \mid U \in R) = \frac{P(U = r \text{ and } U \in R)}{P(U \in R)}$$

Since  $r \in R$ , the event  $\{U = r \text{ and } U \in R\}$  is equivalent to  $\{U = r\}$ . Therefore:

$$P(U = r \mid U \in R) = \frac{P(U = r)}{P(U \in R)}$$

We know that  $P(U = r) = \frac{1}{|S|}$ .

For  $P(U \in R)$ , we use the fact that  $U$  has a uniform distribution on  $S$ :

$$P(U \in R) = \sum_{t \in R} P(U = t) = \sum_{t \in R} \frac{1}{|S|} = \frac{|R|}{|S|}$$

Substituting these values:

$$P(U = r \mid U \in R) = \frac{\frac{1}{|S|}}{\frac{|R|}{|S|}} = \frac{1}{|R|}$$

Since this probability equals  $\frac{1}{|R|}$  for every  $r \in R$ , the conditional distribution assigns equal probability to all elements in  $R$ . This is precisely the definition of a uniform distribution on  $R$ .

Therefore, the distribution of  $U$  conditional on  $\{U \in R\}$  is uniform on  $R$ .  $\square$

**Problem 2**

- Let  $X$  be a random variable with distribution function

$$\mathbb{P}\{X \leq x\} = \begin{cases} 0 & \text{if } x \leq 0, \\ x & \text{if } 0 < x \leq 1, \\ 1 & \text{if } x > 1. \end{cases}$$

Let  $F$  be a distribution function which is continuous and strictly increasing. Show that  $Y = F^{-1}(X)$  is a random variable having distribution function  $F$ .

Is it necessary that  $F$  be continuous and/or strictly increasing?

*Proof.* We need to show that if  $X$  has the given distribution function and  $F$  is continuous and strictly increasing, then  $Y = F^{-1}(X)$  has distribution function  $F$ .

For any real number  $y$ , we have:

$$\begin{aligned} \mathbb{P}\{Y \leq y\} &= \mathbb{P}\{F^{-1}(X) \leq y\} \\ &= \mathbb{P}\{X \leq F(y)\} \end{aligned}$$

This equivalence holds because  $F$  is strictly increasing, so the inequality is preserved when applying  $F$  to both sides.

Since  $X$  has the distribution function:

$$\mathbb{P}\{X \leq x\} = \begin{cases} 0 & \text{if } x \leq 0 \\ x & \text{if } 0 < x \leq 1 \\ 1 & \text{if } x > 1 \end{cases}$$

Therefore:

$$\mathbb{P}\{X \leq F(y)\} = \begin{cases} 0 & \text{if } F(y) \leq 0 \\ F(y) & \text{if } 0 < F(y) \leq 1 \\ 1 & \text{if } F(y) > 1 \end{cases}$$

Since  $F$  is a distribution function, we know  $0 \leq F(y) \leq 1$  for all  $y$ , which means we're always in the middle case where  $\mathbb{P}\{X \leq F(y)\} = F(y)$ . Thus,  $\mathbb{P}\{Y \leq y\} = F(y)$ , which proves that  $Y$  has distribution function  $F$ .

Regarding the necessity of the conditions:

**Strict monotonicity:** If  $F$  is not strictly increasing but merely non-decreasing, then for some values  $a < b$  with  $F(a) = F(b)$ , the inverse function  $F^{-1}$  is not uniquely defined at  $F(a)$ . This makes  $Y = F^{-1}(X)$  ambiguous.

**Continuity:** If  $F$  has jump discontinuities, then there exist values  $c$  in the range of  $F$  for which  $F^{-1}(c)$  is not defined. If  $X$  takes such values with positive probability, then  $Y = F^{-1}(X)$  is not properly defined as a random variable.

Both issues can be addressed by using the generalized inverse function:

$$F^{-1}(u) = \inf\{y : F(y) \geq u\}$$

With this definition,  $Y = F^{-1}(X)$  will have distribution function  $F$  even when  $F$  is not continuous or strictly increasing, provided that  $F$  is at least right-continuous and non-decreasing.  $\square$

### Problem 3

- (a) Show that any discrete random variable may be written as a linear combination of indicator variables.
- (b) Show that any random variable may be expressed as the limit of an increasing sequence of discrete random variables.
- (c) Show that the limit of any increasing convergent sequence of random variables is a random variable.

*Proof.* (a)

Let  $X$  be a discrete random variable taking values in the countable set  $\{x_1, x_2, x_3, \dots\}$ . We can express  $X$  as:

$$X = \sum_{i=1}^{\infty} x_i \mathbf{1}_{\{X=x_i\}}$$

where  $\mathbf{1}_{\{X=x_i\}}$  is the indicator variable for the event  $\{X = x_i\}$ , defined as:

$$\mathbf{1}_{\{X=x_i\}}(\omega) = \begin{cases} 1 & \text{if } X(\omega) = x_i \\ 0 & \text{otherwise} \end{cases}$$

To verify this representation, consider any outcome  $\omega$  in the sample space. If  $X(\omega) = x_j$  for some  $j$ , then  $\mathbf{1}_{\{X=x_i\}}(\omega) = 0$  for all  $i \neq j$  and  $\mathbf{1}_{\{X=x_j\}}(\omega) = 1$ . Therefore:

$$\sum_{i=1}^{\infty} x_i \mathbf{1}_{\{X=x_i\}}(\omega) = x_j \cdot 1 + \sum_{i \neq j} x_i \cdot 0 = x_j = X(\omega)$$

Thus,  $X$  is indeed a linear combination of indicator variables.

(b)

Let  $X$  be any random variable. For each positive integer  $m$ , we partition the real line into intervals of the form  $[k2^{-m}, (k+1)2^{-m})$  for  $k \in \mathbb{Z}$  (i.e., all integers from  $-\infty$  to  $\infty$ ).

We define the discrete random variable  $X_m$  as:

$$X_m = \sum_{k=-\infty}^{\infty} k2^{-m} I_{k,m}$$

where  $I_{k,m}$  is the indicator function of the event  $\{k2^{-m} \leq X < (k+1)2^{-m}\}$ .

This construction has the following properties:

1.  $X_m$  is a discrete random variable because it takes values in the countable set  $\{k2^{-m} : k \in \mathbb{Z}\}$ .
2. For any outcome  $\omega$ , if  $X(\omega) \in [k2^{-m}, (k+1)2^{-m})$  for some integer  $k$ , then  $X_m(\omega) = k2^{-m}$ , which is the left endpoint of the interval containing  $X(\omega)$ .
3. For each fixed  $\omega$ ,  $X_m(\omega) \leq X(\omega)$  since  $X_m(\omega) = k2^{-m}$  when  $X(\omega) \in [k2^{-m}, (k+1)2^{-m})$ .
4. As  $m$  increases, the partition becomes finer and  $X_m(\omega)$  becomes a better approximation of  $X(\omega)$ . Specifically, if  $X(\omega) \in [k2^{-m}, (k+1)2^{-m})$ , then:

$$0 \leq X(\omega) - X_m(\omega) < (k+1)2^{-m} - k2^{-m} = 2^{-m}$$

5. The sequence  $\{X_m(\omega)\}_{m=1}^{\infty}$  is increasing for each fixed  $\omega$ . This is because as  $m$  increases, the left endpoints of the intervals in the partition either stay the same or increase to a value closer to  $X(\omega)$ .

Since  $2^{-m} \rightarrow 0$  as  $m \rightarrow \infty$ , we have  $X_m(\omega) \uparrow X(\omega)$  as  $m \rightarrow \infty$  for all  $\omega$ .

Therefore, any random variable  $X$  can be expressed as the limit of an increasing sequence of discrete random variables  $\{X_m\}_{m=1}^{\infty}$ .

(c)

Let  $\{Y_n\}$  be an increasing sequence of random variables converging to  $Y$ . We need to show that  $Y$  is a random variable, which means we need to prove that for any Borel set  $B$ , the preimage  $Y^{-1}(B) = \{\omega : Y(\omega) \in B\}$  is measurable.

Consider any  $a \in \mathbb{R}$ . We need to show that the set  $\{\omega : Y(\omega) \leq a\}$  is measurable.

Since  $\{Y_n\}$  is increasing and converges to  $Y$ , we have:

$$\{\omega : Y(\omega) \leq a\} = \{\omega : \lim_{n \rightarrow \infty} Y_n(\omega) \leq a\}$$

For an increasing sequence,  $\lim_{n \rightarrow \infty} Y_n(\omega) \leq a$  if and only if  $Y_n(\omega) \leq a$  for all  $n$ . Therefore:

$$\{\omega : Y(\omega) \leq a\} = \bigcap_{n=1}^{\infty} \{\omega : Y_n(\omega) \leq a\}$$

Since each  $Y_n$  is a random variable, each set  $\{\omega : Y_n(\omega) \leq a\}$  is measurable. The countable intersection of measurable sets is measurable, so  $\{\omega : Y(\omega) \leq a\}$  is measurable.

Since the preimage of any interval  $(-\infty, a]$  under  $Y$  is measurable, and the Borel  $\sigma$ -algebra is generated by such intervals,  $Y$  is a random variable.  $\square$

#### Problem 4

Express the distribution functions of

$$X^+ = \max\{0, X\}, \quad X^- = -\min\{0, X\}, \quad |X| = X^+ + X^-, \quad -X,$$

in terms of the distribution function  $F$  of the random variable  $X$ .

*Proof.* Let  $X$  be a random variable with distribution function  $F(x) = \mathbb{P}(X \leq x)$ . We'll express the distribution functions of the derived random variables in terms of  $F$ .

**Distribution function of  $X^+ = \max\{0, X\}$ :**

For  $x < 0$ , since  $\max\{0, X\} \geq 0 > x$  for all values of  $X$ :

$$\mathbb{P}(X^+ \leq x) = 0$$

For  $x \geq 0$ :

$$\begin{aligned} \mathbb{P}(X^+ \leq x) &= \mathbb{P}(\max\{0, X\} \leq x) \\ &= \mathbb{P}(X \leq x) \\ &= F(x) \end{aligned}$$

Therefore:

$$\mathbb{P}(X^+ \leq x) = \begin{cases} 0 & \text{if } x < 0, \\ F(x) & \text{if } x \geq 0. \end{cases}$$

**Distribution function of  $X^- = -\min\{0, X\}$ :**

For  $x < 0$ , since  $X^- \geq 0$  by definition:

$$\mathbb{P}(X^- \leq x) = 0$$

For  $x \geq 0$ , noting that  $X^- = 0$  when  $X \geq 0$  and  $X^- = -X$  when  $X < 0$ :

$$\begin{aligned} \mathbb{P}(X^- \leq x) &= \mathbb{P}((X \geq 0) \cup (X < 0 \text{ and } -X \leq x)) \\ &= \mathbb{P}(X \geq 0) + \mathbb{P}(-x \leq X < 0) \end{aligned}$$

$$\begin{aligned}
 &= 1 - \lim_{y \uparrow 0} F(y) + F(0) - F(-x) \\
 &= 1 - \lim_{y \uparrow -x} F(y)
 \end{aligned}$$

Therefore:

$$\mathbb{P}(X^- \leq x) = \begin{cases} 0 & \text{if } x < 0, \\ 1 - \lim_{y \uparrow -x} F(y) & \text{if } x \geq 0. \end{cases}$$

**Distribution function of  $|X| = X^+ + X^-$ :**

For  $x < 0$ , since  $|X| \geq 0$ :

$$\mathbb{P}(|X| \leq x) = 0$$

For  $x \geq 0$ , using  $|X| = X$  when  $X \geq 0$  and  $|X| = -X$  when  $X < 0$ :

$$\begin{aligned}
 \mathbb{P}(|X| \leq x) &= \mathbb{P}(0 \leq X \leq x) + \mathbb{P}(-x \leq X < 0) \\
 &= F(x) - F(0) + F(0) - F(-x) \\
 &= F(x) - \lim_{y \uparrow -x} F(y)
 \end{aligned}$$

Therefore:

$$\mathbb{P}(|X| \leq x) = \begin{cases} 0 & \text{if } x < 0, \\ F(x) - \lim_{y \uparrow -x} F(y) & \text{if } x \geq 0. \end{cases}$$

**Distribution function of  $-X$ :**

For any real  $x$ :

$$\begin{aligned}
 \mathbb{P}(-X \leq x) &= \mathbb{P}(X \geq -x) \\
 &= 1 - \mathbb{P}(X < -x) \\
 &= 1 - \lim_{y \uparrow -x} F(y)
 \end{aligned}$$

□

## Problem 5

The real number  $m$  is called a median of the distribution function  $F$  whenever  $\lim_{y \rightarrow m} F(y) \leq \frac{1}{2} \leq F(m)$ .

(a) Show that every distribution function  $F$  has at least one median, and that the set of medians of  $F$  is a closed interval of  $\mathbb{R}$ .

(b) Show, if  $F$  is continuous, that  $F(m) = \frac{1}{2}$  for any median  $m$ .

*Proof.* **(a)**

Define  $m = \sup\{x : F(x) < \frac{1}{2}\}$ . We will show that  $m$  is a median.

For any  $y < m$ , by the definition of supremum, we have  $F(y) < \frac{1}{2}$ . This implies that  $\lim_{y \uparrow m} F(y) \leq \frac{1}{2}$ .

Now we need to show that  $F(m) \geq \frac{1}{2}$ . Suppose, for contradiction, that  $F(m) < \frac{1}{2}$ . Since  $F$  is right-continuous, there exists some  $\delta > 0$  such that  $F(m + \epsilon) < \frac{1}{2}$  for all  $0 < \epsilon < \delta$ . But this means that  $m + \frac{\delta}{2}$  belongs to the set  $\{x : F(x) < \frac{1}{2}\}$ , which contradicts  $m$  being the supremum of this set. Therefore,  $F(m) \geq \frac{1}{2}$ .

Thus,  $m$  satisfies  $\lim_{y \uparrow m} F(y) \leq \frac{1}{2} \leq F(m)$ , making it a median.

Similarly, define  $M = \sup\{x : F(x) \leq \frac{1}{2}\}$ . Using a similar argument, we can show that  $M$  is also a median.

For any  $x \in [m, M]$ , we have:

$$\lim_{y \uparrow x} F(y) \leq \lim_{y \uparrow m} F(y) \leq \frac{1}{2}$$

because  $F$  is non-decreasing. Also:

$$F(x) \geq \frac{1}{2}$$

because  $x \geq m$  and  $F$  is non-decreasing. Therefore, every  $x \in [m, M]$  is a median.

Conversely, if  $x < m$ , then  $F(x) < \frac{1}{2}$ , so  $x$  cannot be a median. If  $x > M$ , then by definition of  $M$ , we have  $F(x) > \frac{1}{2}$ , and by right-continuity of  $F$ , we get  $\lim_{y \uparrow x} F(y) > \frac{1}{2}$ , so  $x$  cannot be a median.

Therefore, the set of all medians is exactly the closed interval  $[m, M]$ .

**(b)**

Let  $m$  be any median of the continuous distribution function  $F$ . By definition of median:

$$\lim_{y \uparrow m} F(y) \leq \frac{1}{2} \leq F(m)$$

Since  $F$  is continuous at  $m$ , we have  $\lim_{y \uparrow m} F(y) = F(m)$ . The inequality above becomes:

$$F(m) \leq \frac{1}{2} \leq F(m)$$

This can only be satisfied when  $F(m) = \frac{1}{2}$ .

Therefore, when  $F$  is continuous, every median  $m$  satisfies  $F(m) = \frac{1}{2}$ . □

**Problem 6**

Let  $(\Omega, \mathcal{F})$  be a measurable space,  $A_1, \dots, A_n \in \mathcal{F}$  and  $b_1, \dots, b_n \in \mathbb{R}$ . Set  $X = b_1 1_{A_1} + \dots + b_n 1_{A_n}$ . Suppose that for all  $i \neq j$ ,  $A_i \cap A_j = \emptyset$  and  $b_i \neq b_j$ . Show that  $\sigma(X) = \sigma(\{A_1, \dots, A_n\})$ .

*Proof.* Let  $(\Omega, \mathcal{F})$  be a measurable space,  $A_1, \dots, A_n \in \mathcal{F}$  be disjoint sets and  $b_1, \dots, b_n \in \mathbb{R}$  be distinct real numbers. We define  $X = \sum_{i=1}^n b_i 1_{A_i}$ .

We want to show that  $\sigma(X) = \sigma(\{A_1, \dots, A_n\})$ .

$\Rightarrow \sigma(X) \subseteq \sigma(\{A_1, \dots, A_n\})$ :

For any Borel set  $B \in \mathcal{B}(\mathbb{R})$ , we need to show that  $X^{-1}(B) \in \sigma(\{A_1, \dots, A_n\})$ .

Note that  $X$  takes only the values from the set  $\{b_1, \dots, b_n, 0\}$ , where 0 is taken when  $\omega \notin \bigcup_{i=1}^n A_i$ . Let's define  $A_0 = \Omega \setminus \bigcup_{i=1}^n A_i$  and  $b_0 = 0$  for convenience.

For any  $\omega \in \Omega$ , there is a unique index  $j \in \{0, 1, \dots, n\}$  such that  $\omega \in A_j$ , and thus  $X(\omega) = b_j$ .

Therefore,

$$\begin{aligned} X^{-1}(B) &= \{\omega \in \Omega : X(\omega) \in B\} \\ &= \{\omega \in \Omega : \exists j \in \{0, 1, \dots, n\} \text{ such that } \omega \in A_j \text{ and } b_j \in B\} \\ &= \bigcup_{j=0}^n \{A_j : b_j \in B\} \end{aligned}$$

Since each  $A_j \in \sigma(\{A_1, \dots, A_n\})$  (note that  $A_0$  can be expressed as  $\Omega \setminus \bigcup_{i=1}^n A_i$ , which is in  $\sigma(\{A_1, \dots, A_n\})$ ), we have  $X^{-1}(B) \in \sigma(\{A_1, \dots, A_n\})$ .

Thus,  $\sigma(X) \subseteq \sigma(\{A_1, \dots, A_n\})$ .

$\Leftarrow \sigma(\{A_1, \dots, A_n\}) \subseteq \sigma(X)$ :

We need to show that for each  $i \in \{1, \dots, n\}$ ,  $A_i \in \sigma(X)$ .

Since the values  $b_1, \dots, b_n$  are distinct, for each  $i \in \{1, \dots, n\}$ , the set  $\{b_i\}$  is a Borel set in  $\mathbb{R}$ . We have:

$$\begin{aligned} X^{-1}(\{b_i\}) &= \{\omega \in \Omega : X(\omega) = b_i\} \\ &= \{\omega \in \Omega : \sum_{j=1}^n b_j 1_{A_j}(\omega) = b_i\} \end{aligned}$$

For any  $\omega \in A_i$ , we have  $X(\omega) = b_i$  because  $A_i \cap A_j = \emptyset$  for  $i \neq j$ . For any  $\omega \notin A_i$ , we have  $X(\omega) \neq b_i$  because the values  $b_1, \dots, b_n$  are distinct and the sets  $A_1, \dots, A_n$  are disjoint.

Therefore,  $X^{-1}(\{b_i\}) = A_i$ , which implies  $A_i \in \sigma(X)$  for each  $i \in \{1, \dots, n\}$ .



Hence,  $\sigma(\{A_1, \dots, A_n\}) \subseteq \sigma(X)$ .

So we conclude that  $\sigma(X) = \sigma(\{A_1, \dots, A_n\})$ . □

### Problem 7

Consider the random walk on  $\{0, 1, \dots, N\}$  that reflects at 0, introduced in the class. More precisely, let  $S_0 = k, S_{j+1} = S_j + X_j$ , where  $X_j = 1$  if  $S_j = 0$ , otherwise  $\mathbb{P}[X_j = 1] = p$  and  $\mathbb{P}[X_j = -1] = q$ . Compute the expected number of steps to first reach state  $N$ , as a function of  $k$ . Distinguish the cases  $p = q$  and  $p \neq q$ .

*Proof.* Random walk on  $\{0, 1, \dots, N\}$  with:

$$\begin{aligned}\mathbb{P}[X_j = 1 \mid S_j = 0] &= 1 \text{ (reflection at 0)} \\ \mathbb{P}[X_j = 1 \mid S_j \neq 0] &= p \\ \mathbb{P}[X_j = -1 \mid S_j \neq 0] &= q = 1 - p\end{aligned}$$

Let  $h_k$  be the expected number of steps to first reach state  $N$  from initial state  $k$ .

**Recursive equations:**

$$\begin{aligned}h_k &= 1 + ph_{k+1} + qh_{k-1}, \quad 0 < k < N \\ h_0 &= 1 + h_1 \\ h_N &= 0\end{aligned}$$

**Case 1:** ( $p = q = 1/2$ )

For  $0 < k < N$ , the recursive equation becomes:

$$\begin{aligned}h_k &= 1 + \frac{1}{2}h_{k+1} + \frac{1}{2}h_{k-1} \\ \Rightarrow h_{k+1} - 2h_k + h_{k-1} &= -2\end{aligned}$$

This second-order linear difference equation has general solution:

$$h_k = A + Bk + k(N - k)$$

From boundary conditions:

$$\begin{aligned}h_N = 0 &\Rightarrow A + BN = 0 \\ h_0 = 1 + h_1 &\Rightarrow B = -1 \Rightarrow A = N\end{aligned}$$

Therefore:

$$h_k = (N - k)(k + 1)$$

**Case 2:** ( $p \neq q$ )

Rearranging the recursive equation:

$$ph_{k+1} - h_k + qh_{k-1} = -1$$

Let  $\rho = \frac{q}{p}$ . The general solution has form:

$$h_k = A + B\rho^k + \frac{k}{q-p} + \frac{q}{q-p}$$

Applying boundary conditions:

$$\begin{aligned} h_N &= 0 \\ h_0 &= 1 + h_1 \end{aligned}$$

Solving for constants  $A$  and  $B$ :

$$\begin{aligned} A &= \frac{1}{p-q} \left( \frac{1-\rho^N}{1-\rho} - N \right) \\ B &= \frac{1}{p-q} \cdot \frac{1}{1-\rho^N} \end{aligned}$$

Final solution when  $p \neq q$ :

$$h_k = \begin{cases} \frac{N-k}{p-q} & \text{if } p > q \\ \frac{1}{p-q} \left[ \frac{1-\rho^N}{1-\rho} - N - \frac{\rho^k - \rho^N}{1-\rho^N} \right] & \text{if } p < q \end{cases}$$

□

**Problem 8**

If  $F : \mathbb{R} \rightarrow [0, 1]$  satisfies monotone increasing, right continuity,  $\lim_{x \rightarrow -\infty} F(x) = 0$  and  $\lim_{x \rightarrow +\infty} F(x) = 1$ , show that  $F$  is the distribution function of some random variable on some probability space. (Hint: you may want to revisit Section 2.3, exercise 3)

*Proof.* Let  $F : \mathbb{R} \rightarrow [0, 1]$  satisfy:

- Monotone increasing
- Right continuous
- $\lim_{x \rightarrow -\infty} F(x) = 0$
- $\lim_{x \rightarrow +\infty} F(x) = 1$

We construct a random variable with distribution function  $F$  using the hint that the limit of any increasing sequence of random variables is a random variable.

Let  $(\Omega, \mathcal{F}, \mathbb{P}) = ((0, 1], \mathcal{B}((0, 1]), \lambda)$  where  $\lambda$  is Lebesgue measure.

For each  $n \in \mathbb{N}$ , define:

$$X_n(\omega) = \inf\{x = k2^{-n} : F(x) \geq \omega, k \in \mathbb{Z}\}$$

The sequence  $\{X_n\}$  is decreasing:  $X_n(\omega) \geq X_{n+1}(\omega)$  for all  $\omega \in \Omega$ .

Define  $X(\omega) = \lim_{n \rightarrow \infty} X_n(\omega)$ . By the hint,  $X$  is a random variable since it's the limit of a monotone sequence of random variables.

From the construction and right continuity of  $F$ :

$$X(\omega) = \inf\{x \in \mathbb{R} : F(x) \geq \omega\}$$

For any  $a \in \mathbb{R}$ :

$$\begin{aligned} \mathbb{P}(X \leq a) &= \mathbb{P}(\{\omega \in \Omega : F(a) \geq \omega\}) \\ &= F(a) \end{aligned}$$

Therefore,  $F$  is the distribution function of the random variable  $X$ . □