Paper Title

Abstract

Replace this paragraph with the abstract of your paper. State the problem, the assumptions, the main result and its scope. The examples below demonstrate the reference colors and can be replaced directly.

1 Introduction

Introduce the problem and explain the main contribution. A literature citation such as (Robbins and Monro, 1951) uses author–year labels in deep blue. Internal references such as Assumption 3.1 and Theorem 4.1, (1), and Algorithm 1, Figure 1, and Table 1 have dark-red numbers. The text of the paper, displayed equations and theorem headings remain black.

2 Related works

Discuss the relevant literature and distinguish the assumptions and conclusions. For a citation used as part of a sentence, write, for example, Robbins and Monro (1951).

3 Problem setup

Specify the objective, the domain and the algorithm. The Euclidean norm is denoted by $\|\cdot\|$. The following assumption and algorithm are illustrative examples rather than requirements of this template.

Assumption 3.1 (L-smoothness)**.** The function $f : \mathbb{R}^d \to \mathbb{R}$ is differentiable, and there is $L > 0$ such that $\|\nabla f(x) - \nabla f(y)\| \le L\|x - y\|$ for all $x, y \in \mathbb{R}^d$.

Algorithm 1 Gradient descent: an illustrative algorithm

Require: Initial point $x_0 \in \mathbb{R}^d$, stepsize $\eta > 0$, iterations $T \ge 1$

1: **for** $t = 0, \dots, T-1$ **do**
2: $\quad x_{t+1} \leftarrow x_t - \eta \nabla f(x_t)$
3: **end for**
4: **return** x_T

4 Main results

Theorem 4.1 (Illustrative descent result)**.** *Under Assumption 3.1, if $0 < \eta \le 1/L$, the iterates of Algorithm 1 satisfy, for every $0 \le t < T$,*

$$f(x_{t+1}) \le f(x_t) - \frac{\eta}{2}\|\nabla f(x_t)\|^2. \tag{1}$$

The proof is in Appendix A. Use `\cref{...}` for a named reference and `\eqref{...}` for an equation number in parentheses. Use `\citep{...}` or `\citet{...}` for literature.

5 Experiments

Describe the experimental setup and report the findings. The examples in Figure 1 and Table 1 demonstrate captions and references.




Figure 1: A self-contained figure placeholder. Figure captions remain black.

Table 1: The reference palette used throughout the template.

Element	Color
Internal cross-references	Dark red
Literature citations and URLs	Deep blue
Body text, equations and captions	Black

6 Conclusion

Summarize the results and their limitations. Replace the illustrative content with your own paper while retaining the preamble and appendix contents commands as needed.

References

H. Robbins and S. Monro. A stochastic approximation method. *The Annals of Mathematical Statistics*, 22(3):400–407, 1951. https://doi.org/10.1214/aoms/1177729586.

Contents of the Appendices

A Proofs **5**
A.1 Proof of the illustrative theorem 5

B Additional details **5**
B.1 Supplementary material 5

A Proofs

A.1 Proof of the illustrative theorem

Proof of Theorem 4.1. The smoothness assumption gives

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{L}{2}\|y - x\|^2. \tag{A.1}$$

Substituting $x = x_t$ and $y = x_t - \eta \nabla f(x_t)$ into (A.1), we obtain

$$f(x_{t+1}) \leq f(x_t) - \eta \left(1 - \frac{L\eta}{2}\right) \|\nabla f(x_t)\|^2.$$

Since $L\eta \leq 1$, this proves (1). □

B Additional details

B.1 Supplementary material

Place further proofs, implementation details or additional results here. The appendix contents update automatically as sections are added or removed. Compile twice, or use `latexmk`, to refresh page numbers.