

VE230 Homework 3

2020 Summer

P.3-22 The polarization in a dielectric cube of side L centered at the origin is given by $\mathbf{P} = P_0(\mathbf{a}_x x + \mathbf{a}_y y + \mathbf{a}_z z)$.

- Determine the surface and volume bound-charge densities
- Show that the total bound charge is zero.

P.3-23 Determine the electric field intensity at the center of a small spherical cavity cut out of a large block of dielectric in which a polarization \mathbf{P} exists.

P.3-25 Assume that the $z=0$ plane separates two lossless dielectric regions with $\epsilon_{r1} = 2$ and $\epsilon_{r2} = 3$. If we know that \mathbf{E}_1 in region 1 is $\mathbf{a}_x 2y - \mathbf{a}_y 3x + \mathbf{a}_z (5 + z)$, what do we also know about \mathbf{E}_2 and \mathbf{D}_2 in region 2? Can we determine \mathbf{E}_2 and \mathbf{D}_2 at any point in region 2? Explain.

P.3-28 Dielectric lenses can be used to collimate electromagnetic fields. In Fig.1 the left surface of the lens is that of a circular cylinder, and the right surface is a plane. If \mathbf{E}_1 at point $P(r_0, 45^\circ, z)$ in region 1 is $\mathbf{a}_r 5 - \mathbf{a}_\phi 3$, what must be the dielectric constant of the lens in order that \mathbf{E}_3 in region 3 is parallel to the x -axis?




Figure 1: A dielectric lens

P.3-32 The radius of the core and the inner radius of the outer conductor of a very long coaxial transmission line are r_i and r_o , respectively. The space between the conductors is filled with two coaxial layers of dielectrics. The dielectric constants of the dielectrics are ϵ_{r1} for $r_i < r < b$ and ϵ_{r2} for $b < r < r_o$. Determine its capacitance per unit length.