

Special examples

Yashas.N

1 Sequences and Series

1. Riemann-Zeta function $\zeta : \mathbb{C} \rightarrow \mathbb{C} + \infty$ defined by

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

- (a) if $1 < \operatorname{Re}(s) < \infty$ then the series converges uniformly and absolutely
(b) clearly ζ is analytic for $\operatorname{Re}(s) > 1$
(c) **Euler's Product formula:**

$$\zeta(z) = \prod \left(1 - \frac{1}{p^s}\right)^{-1}$$

where the product ranges over all primes p which implies $\zeta(s) \neq 0$ if $\operatorname{Re}(s) > 1$. More generally we have

$$\zeta(s)(1 - 2^{-s})(1 - 3^{-s}) \dots (1 - p_N^{-s}) = \sum m^s = 1 + p_{N+1}^{-s} \dots$$

where the R.H.S ranges for all +ve integers that contain none of prime factors $2, 3, \dots, p_N$

- (d) now if $1 < s < \infty$ (i.e. s is real > 1) then

$$\zeta(s) = s \int_1^{\infty} \frac{[x]}{x^{s+1}} dx$$

where $[x]$ is greatest integer $\leq x$

- (e) more generally if $\operatorname{Re}(s) > 1$ then

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{x^s - 1}{e^x - 1} dx$$

where $\Gamma(z)$ is defined by product representation for complex numbers.

2.
$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$$

- (a) is convergent to $\ln(2)$
(b) does not converge absolutely.

2 Functions in \mathbb{R}

1. Dirichlet Function $\delta(x)$

$$\delta(x) = \begin{cases} 1 & \text{if } x \text{ is rational} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$

- (a) δ is not Riemann Integrable in any interval $[a, b]$
 - (b) δ is Lebesgue Integrable in \mathbb{R} and has 0 integral value with usual lebesgue measure as the set for which δ is not zero is countable.
2. $f : \mathbb{R} \rightarrow \mathbb{R}$ such that for every rational $r = m/n$ where $n > 0$ and $m, n \in \mathbb{Z}$ with out any common divisors then $f(r) = f(m/n) = 1/n$, $x = 0$ take $n = 1$ i.e. $f(0) = 1$ and $f(x) = 0$ if x is irrational i.e.

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is irrational} \\ \frac{1}{n} & \text{if } x = \frac{m}{n} \end{cases}$$

- (a) $f(x)$ is continuous at every irrational number
 - (b) $f(x)$ is discontinuous at rational with simple discontinuities
3. $\sin(x)$

- (a) can be defined without geometric interpretation as $\sin x = (e^{ix} - e^{-ix})/2$ for real x

- (b) \sin is continuous one-one function in domain $[-\pi, \pi]$ onto $[-1, 1]$ hence inverse \sin^{-1} is defined in this area.

- (c) we see that $\frac{2}{\pi}x \leq \sin x \leq x$ holds $\forall x \in [0, \pi/2]$

