

Hodge theory and derived categories of cubic fourfolds

Nicolas Addington, Richard P. Thomas

arXiv:1211.3758v3; DOI:10.1215/00127094-2738639

<http://arxiv.org/abs/1211.3758v3>

Two K3 shadows of a cubic fourfold

Let $X \subset \mathbb{P}^5$ be a smooth complex cubic fourfold. Removing the exceptional collection from its derived category leaves the K3 category

$$\mathcal{A}_X = \langle \mathcal{O}_X, \mathcal{O}_X(1), \mathcal{O}_X(2) \rangle^\perp.$$

It is called *geometric* if $\mathcal{A}_X \simeq D(S)$ for a projective K3 surface S . Write h for the hyperplane class. On the Hodge side, special cubics form irreducible divisors $\mathcal{C}_d$: there is a primitive rank-two lattice $\langle h^2, T \rangle \subset H^{2,2}(X, \mathbb{Z})$ of discriminant d . Such a divisor is nonempty precisely when $d > 6$ and $d \equiv 0$ or $2 \pmod{6}$. Call d *admissible* when, in addition, it is divisible by neither 4, nor 9, nor any odd prime $p \equiv 2 \pmod{3}$.

Main theorem (Addington–Thomas, Theorem 1.1). *If $\mathcal{A}_X$ is geometric, then $X \in \mathcal{C}_d$ for an admissible d . Conversely, for every admissible d , the cubics $X \in \mathcal{C}_d$ for which $\mathcal{A}_X$ is geometric form a nonempty Zariski-open dense subset.*

Thus Hassett’s Hodge-theoretic K3 condition and Kuznetsov’s categorical one agree generically along every admissible Noether–Lefschetz divisor.

Ingredient I: an integral Mukai-lattice bridge

The authors define the Mukai lattice of $\mathcal{A}_X$ using topological K -theory. Two projected line classes λ_1, λ_2 span a primitive copy of $-A_2$ in $K_{\text{num}}(\mathcal{A}_X)$, and Proposition 2.3 gives an integral isometry

$$\langle \lambda_1, \lambda_2 \rangle^\perp \cong H_{\text{prim}}^4(X, \mathbb{Z}). \quad (1)$$

The second Chern class also identifies the corresponding quotient lattices; primitivity, signatures, and discriminants can therefore be transferred between algebraic K -classes and $H^{2,2}(X, \mathbb{Z})$.

Theorem 3.1 packages the arithmetic condition on d into the categorical criterion

$$X \in \mathcal{C}_d \text{ for some admissible } d \iff K_{\text{num}}(\mathcal{A}_X) \text{ contains } U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (2)$$

If $\mathcal{A}_X \simeq D(S)$, the classes of a point sheaf and its ideal sheaf span precisely this hyperbolic plane, proving the first direction of the main theorem. The converse in (2) uses lattice embeddings and, in the rank-four case, a binary quadratic form whose represented primes control the forbidden factors of d .

Ingredient II: one geometric seed on each divisor

A cubic in $\mathcal{C}_8$ contains a plane P . Blowing up P gives a quadric-surface fibration over $\mathbb{P}^2$. If its discriminant sextic is smooth, the branched double cover is a K3 surface S , and $\mathcal{A}_X \simeq D(S, \alpha)$ for a Brauer class α . Put $Q = h^2 - P$, the class of a smooth quadric fibre. Theorem 4.1 constructs, for every nonempty $\mathcal{C}_d$, a point $X_0 \in \mathcal{C}_d \cap \mathcal{C}_8$ with an extra class T satisfying $T \cdot Q = 1$. A lattice and period-map argument also proves that this seed’s sextic is smooth by excluding any other plane meeting P . The intersection condition kills α , so $\mathcal{A}_{X_0} \simeq D(S_0)$.

For admissible d , Proposition 5.1 then changes this to the *right* equivalence: its cohomological transform extends the prescribed Hodge isometry

$$H_{\text{prim}}^2(S_0, \mathbb{Z})(-1) \cong \langle h^2, T \rangle^\perp. \quad (3)$$

This alignment is what lets the Fourier–Mukai kernel move away from $\mathcal{C}_8$ while X moves inside $\mathcal{C}_d$.

Ingredient III: deform the kernel, not just the Hodge class

After a finite level cover of $\mathcal{C}_d$, Proposition 5.2 supplies families of cubics $\mathcal{X}$ and K3 surfaces $\mathcal{S}$ whose period lines match under (3). Let P be the central Fourier–Mukai kernel. Variation of Hodge structure says that matching period lines make the contracted Kodaira–Spencer classes match. Proposition 6.2 translates this into Hochschild cohomology:

$$\Phi_P^{HH^*}(\kappa_X) = \kappa_S.$$

Theorem 7.1 identifies the first-order obstruction with $(\kappa_S, \kappa_X) \circ \text{At}(P) \in \text{Ext}^2(P, P)$. The two partial Atiyah classes appear with opposite signs under convolution, so the displayed Hochschild equality makes this obstruction vanish.

First order is not enough. Proposition 7.6 uses T^1 -lifting to turn compatible sideways first-order deformations into an extension from $\mathbb{C}[t]/(t^{n+1})$ to $\mathbb{C}[t]/(t^{n+2})$. Theorem 7.7 iterates this to all orders. There is no choice to accumulate:

$$\text{Ext}^1(P, P) \cong HH^1(S) = H^1(\mathcal{O}_S) \oplus H^0(T_S) = 0,$$

and rigidity also preserves full faithfulness.

Finally, the locally finitely presented stack of complexes algebraizes the formal kernel. Full faithfulness and orthogonality to $\mathcal{O}_X, \mathcal{O}_X(1), \mathcal{O}_X(2)$ persist on an open set. There $D(S_t)$ is nonzero, $\mathcal{A}_{X_t}$ is indecomposable, and both Serre functors are the shift [2]; together these facts upgrade the embedding to an equivalence. The image is a nonempty open subset of the irreducible divisor $\mathcal{C}_d$, hence dense.

Algebraic-cycle payoff

The kernel itself may fail to have a controllable limit at the boundary of that open set, but its Mukai vector is an algebraic cycle. Writing ℓ for the K3 polarization, Theorem 1.2 adds a large multiple of $(\ell + h)^3$ to make its codimension-three part effective, then uses properness of a relative Hilbert scheme. Consequently *every* $X \in \mathcal{C}_d$ with admissible d has a degree- d polarized K3 surface S and a rational algebraic cycle on $S \times X$ inducing

$$H_{\text{prim}}^2(S, \mathbb{Z})(-1) \cong \langle h^2, T \rangle^\perp.$$

Theorem 1.3 goes further: any rational Hodge class on $S \times X$ inducing an integral Hodge isometry between the transcendental lattices is algebraic.

Scope and hardest point

The theorem proves geometricity only on an open dense subset, not on all of $\mathcal{C}_d$; the unresolved issue is choosing the right limiting family of derived objects. It also does not prove the motivating rationality conjecture for cubic fourfolds. The technical bottleneck is the passage from a Hodge-compatible first-order motion to a unique all-orders perfect kernel and then to an algebraic family; the Atiyah-class cancellation, T^1 -lifting, and rigidity address these three separate steps.