

The arithmetic Hodge Index Theorem and rigidity of dynamical systems over function fields

Alexander Carney

arXiv:1810.06342

<http://arxiv.org/abs/1810.06342v2>

The rigidity question

Let $K = k(B)$ be the function field of a smooth projective curve and let X/K be projective. A polarized dynamical system (f, L, q) consists of an endomorphism $f : X \rightarrow X$, an ample line bundle L , and an isomorphism $f^*L \simeq L^{\otimes q}$ with $q > 1$. Carney equips L with an f -admissible adelic metric $\bar{L}_f$ whose height is the Call–Silverman canonical height. Write

$$Z_f = \{x \in X(\bar{K}) : h_{\bar{L}_f}(x) = 0\}.$$

Unlike over a number field, Z_f can strictly contain the preperiodic points because the constant field may destroy Northcott finiteness.

Main theorem (Carney, Theorem 4.6). *Let (f, L) and (g, M) be polarized algebraic dynamical systems on X , and let Z be the Zariski closure of $Z_f \cap Z_g$. Then*

$$Z_f \cap Z(\bar{K}) = Z_g \cap Z(\bar{K}).$$

In particular, if the common height-zero points are Zariski dense in X , then $Z_f = Z_g$.

Finite extensions and normalizations used in the proof do not change the claim. If the constant field is finite, $Z_f = \text{Prep}(f)$, so a dense intersection of two preperiodic sets forces equality (Corollary 1.3).

Ingredient I: the arithmetic Hodge index equality case

Assume first that k is algebraically closed, $K = k(B)$, and X is geometrically normal of dimension n . For an integrable adelic $\mathbf{Q}$ -line bundle $\bar{N}$ and big nef $\bar{A}_1, \dots, \bar{A}_{n-1}$, Theorem 3.1 assumes $NA_1 \cdots A_{n-1} = 0$ if $n \geq 2$, or $\deg N = 0$ if $n = 1$, and gives

$$\bar{N}^2 \bar{A}_1 \cdots \bar{A}_{n-1} \leq 0. \tag{1}$$

Under the theorem's stronger equality hypotheses—the $\bar{A}_i$ are arithmetically positive and $\bar{N}$ is $\bar{A}_i$ -bounded—equality holds exactly when

$$\bar{N} \in \pi^* \widehat{\text{Pic}}(K)_{\mathbf{Q}} + \text{Tr}_{K/k}(\text{Pic}^0(X))_{\mathbf{Q}}. \tag{2}$$

The first summand changes a height only by a constant; the second is the constant-field part of the Picard variety. Theorem 3.2 correspondingly identifies numerical triviality with an identically zero height. A general vertical pullback contributes a constant, but the theorem's subgroup has arithmetic degree zero, so that constant vanishes.

For a curve, write $\bar{N} = \bar{N}_0 + \bar{V}$ with $\bar{N}_0$ flat and $\bar{V}$ vertical. Then $\bar{N}^2 = -2h_{\text{NT}}(N) + \sum_v \bar{V}_v^2$, where every $\bar{V}_v \leq 0$ and equality forces the vertical metric to be constant; the Néron–Tate kernel is the K/k -trace. In higher dimension, Bertini sections reduce the inequality inductively. The hard part is equality: a flat bundle restricting to the trace on a hyperplane section must lift to the trace on X . Lemma 3.6 proves the required compatibility of trace with homomorphisms of abelian varieties; the local Hodge index theorem then forces the remaining vertical metric to be constant.

Ingredient II: canonical adelic metrics from a spectral gap

Modulo the trace, pullback by a polarization has two disjoint spectral scales (Lemma 4.1): absolute value $q^{1/2}$ on Pic^0 and q on the Néron–Severi group. Their minimal polynomials are coprime, so the sequence

$$0 \longrightarrow \text{Pic}_{\text{tr}}^0(X)_{\mathbf{Q}} \longrightarrow \text{Pic}_{\text{tr}}(X)_{\mathbf{Q}} \longrightarrow \text{NS}(X)_{\mathbf{Q}} \longrightarrow 0 \quad (3)$$

has a unique f^* -equivariant splitting. On metrics, if R is the combined minimal polynomial, $R(f^*)$ is invertible on continuous metric errors: each factor $(1 - f^*/\lambda)^{-1}$ is a convergent geometric series since $|\lambda| > 1$. Theorem 4.2 thereby assigns every line bundle an f -admissible adelic metric; it is flat on Pic^0 and nef for ample classes in the split Néron–Severi part.

This construction also proves that Z_f is independent of the chosen polarization. If L and M both polarize f , ampleness gives $cL - M$ ample for some $c > 0$, and positivity of canonical heights yields

$$0 \leq h_{\overline{M}_f}(x) \leq ch_{\overline{L}_f}(x). \quad (4)$$

The reverse inclusion follows symmetrically (Lemma 4.7).

Proof roadmap: common small points force equal heights

Normalize an irreducible component Y of Z and put $d = \dim Y$. Choose the f - and g -equivariant lifts L_f, L_g of the same ample class in $\text{NS}(X)_{\mathbf{Q}}$, with nef admissible metrics $\overline{L}_f, \overline{L}_g$. Their sum $\overline{A} = \overline{L}_f + \overline{L}_g$ has a dense set of height-zero points on Y . The successive-minima inequality therefore gives

$$0 = h_{\overline{A}}(Y), \quad 0 = (\overline{L}_f|_Y + \overline{L}_g|_Y)^{d+1}. \quad (5)$$

Every mixed intersection in the binomial expansion is nonnegative, so each is zero. For $\overline{D} = \overline{L}_f|_Y - \overline{L}_g|_Y$ this implies

$$\overline{D}^2 \overline{A}^{d-1} = 0, \quad D \cdot \overline{A}^{d-1} = 0. \quad (6)$$

The underlying D is numerically trivial and $\overline{D}$ is $\overline{A}$ -bounded. Although $\overline{A}$ is only nef, adding the pullback of a positive-degree adelic bundle from $\text{Spec } K$ makes it arithmetically positive without changing (6), because $\overline{D}$ is flat. Applying (1)–(2) shows that $h_{\overline{L}_f} - h_{\overline{L}_g}$ is constant on Y . It vanishes on the dense common zero set, hence the two heights coincide on Y . Equation (4) then identifies their zero sets, component by component.

Toy model, technical bottleneck, and scope

The numerical core is the elementary identity

$$0 = (a + b)^{d+1} = \sum_{i=0}^{d+1} \binom{d+1}{i} a^i b^{d+1-i} \quad (a, b \geq 0), \quad (7)$$

which forces every mixed term to vanish. Nef adelic intersections play the roles of a and b ; the real work is then the equality classification for the resulting difference class. That classification, especially lifting the constant-field trace through hyperplane sections, is the paper’s technical bottleneck.

Over a finite constant field, Northcott converts the theorem into a statement about preperiodic points. Over a general constant field the paper deliberately asserts rigidity of height-zero loci instead. For non-isotrivial rational maps on $\mathbf{P}^1$, Baker’s theorem restores $Z_f = \text{Prep}(f)$ and gives Corollary 1.4; no analogous blanket identification is claimed in higher dimension.