

Rational curves on fibered Calabi-Yau manifolds

Simone Diverio, Claudio Fontanari, Diletta Martinelli

arXiv:1607.01561

<http://arxiv.org/abs/1607.01561v3>

A rational curve forced by an elliptic fibration

Work over $\mathbb{C}$, and assume throughout that $n = \dim X > 2$. The theorem does not begin with an elliptic fibration with a section: the map need not have connected fibers, and the base need not be smooth. Instead, ellipticity is recovered from the canonical-bundle hypothesis.

Main theorem (Diverio–Fontanari–Martinelli, Theorem 1.1). *Let X be a smooth projective manifold with finite fundamental group. Suppose that there are a projective variety B , a morphism $f: X \rightarrow B$ whose general fiber has dimension one, and a line bundle L on B such that*

$$K_X \simeq f^*L. \quad (1)$$

Then X contains a rational curve.

In particular, with the paper’s convention that a Calabi–Yau manifold is smooth and projective with $K_X \simeq \mathcal{O}_X$ and finite fundamental group, every elliptically fibered Calabi–Yau manifold of dimension greater than two contains a rational curve (Corollary 1.2). This supplies a broad fibered case of the general existence problem for rational curves on Calabi–Yau manifolds.

Ingredient I: remove monodromy and higher-dimensional fibers

Pass to the finite universal cover $\pi: \tilde{X} \rightarrow X$ and take the Stein factorization

$$\tilde{X} \xrightarrow{\phi} B' \xrightarrow{\nu} B \quad (f \circ \pi = \nu \circ \phi). \quad (2)$$

Then $\tilde{X}$ is smooth projective and simply connected, ϕ has connected fibers, and $K_{\tilde{X}} \simeq \phi^*\nu^*L$. On a smooth general fiber E , the relative tangent sequence gives $K_E \simeq K_{\tilde{X}}|_E \simeq \mathcal{O}_E$; hence E is an elliptic curve. A rational curve on $\tilde{X}$ descends to one on X , so it suffices to prove the simply connected version, Theorem 2.1.

Because $K_{\tilde{X}}$ is pulled back, $-K_{\tilde{X}}$ is ϕ -nef. Kawamata’s theorem (Theorem 2.2) says that every component of the locus where a fiber has dimension larger than one is uniruled. If that locus is nonempty, the desired curve is already present. Thus the remaining case is an equidimensional elliptic fiber space; this reduction is essential for controlling codimensions after deleting the discriminant.

Ingredient II: the discriminant must contain a divisor

Let $Z \subset B'$ be the union of the singular locus of B' and the singular values of ϕ , and set $B^0 = B' \setminus Z$, $X^0 = \phi^{-1}(B^0)$. Lemma 2.3 proves that Z has a codimension-one component. Indeed, if $\text{codim}_{B'} Z \geq 2$, equidimensionality makes $\tilde{X} \setminus X^0$ codimension at least two, so $\pi_1(X^0) = \pi_1(\tilde{X}) = 1$. Ehresmann’s theorem makes $\phi^0: X^0 \rightarrow B^0$ a differentiable fiber bundle, and its homotopy sequence then makes B^0 simply connected as well.

Consequently the local systems $R^q\phi_*^0\mathbb{Q}$ are constant. Deligne’s degeneration of the Leray spectral sequence at E_2 would make

$$H^0(B^0, R^1\phi_*^0\mathbb{Q}) \simeq H^1(E, \mathbb{Q}) \simeq \mathbb{Q}^2 \quad (3)$$

a graded piece of $H^1(X^0, \mathbb{Q}) = 0$, a contradiction. This is the proof’s hardest conceptual step: simple connectedness forces genuine codimension-one degeneration of the elliptic fibers.

Ingredient III: cut down to a controlled elliptic surface

Choose a very ample H on B' sufficiently positive that $H^{\otimes(n-2)} \otimes \nu^*L$ is globally generated. A general complete-intersection curve $C \subset B'$ cut out by $n-2$ members of $|H|$, together with its inverse-image surface $S \subset \tilde{X}$, is smooth by Bertini. Since B' is normal, a general C avoids its codimension-two singular locus but meets a divisorial component of Z , so the induced elliptic surface $\phi: S \rightarrow C$ has a singular fiber. If S is not relatively minimal, an exceptional (-1) -curve already gives the required rational curve.

In the relatively minimal case, iterated adjunction gives Lemma 2.5:

$$K_S \simeq \phi^*(\nu^*L \otimes H^{\otimes(n-2)}) \Big|_S, \quad (4)$$

so K_S is globally generated. For a threefold, this is the especially transparent picture: one general hyperplane section of the surface base produces an elliptic surface, while (4) keeps its canonical system free.

Ingredient IV: global generation rules out multiple fibers

For a relatively minimal elliptic surface whose multiple fibers are $m_i F_i$, the canonical bundle formula used in the paper is

$$K_S \simeq \phi^*G \otimes \mathcal{O}_S \left(\sum_i (m_i - 1) F_i \right) \quad (5)$$

for a line bundle G on C . If $m_i > 1$, then $\mathcal{O}_{F_i}(F_i)$ is torsion of order m_i , and (5) implies that $K_S|_{F_i}$ has no nonzero section. This contradicts the global generation from (4), proving Proposition 2.6: there are no multiple fibers.

The singular fiber forced above is therefore nonmultiple. Kodaira's classification finishes the argument. An irreducible singular fiber is a nodal or cuspidal rational curve; a reducible nonmultiple fiber has smooth rational (-2) -curves as its irreducible components. In either case S , hence $\tilde{X}$ and finally X , contains a rational curve.

Roadmap, scope, and omissions

The logical chain is short but rigid: finite fundamental group permits the simply connected cover; Kawamata disposes of nonequidimensional fibers; Lemma 2.3 forces a divisorial discriminant; a general slice turns that degeneration into a singular elliptic-surface fiber; and global generation plus the canonical bundle formula makes Kodaira's classification produce the curve.

The hypotheses are stronger than numerical relative triviality: (1) is an isomorphism of line bundles. The result asserts existence, not a count or deformation class, and the proof uses both the finite fundamental group and complex Hodge theory. The paper separately proves that a Calabi–Yau manifold fibred over a curve with general abelian fiber contains a rational multisection (Theorem 1.6 and Corollary 1.7), by a different nef/non-nef divisor dichotomy; that second argument is not developed here.