

# 1-Cycles on Fano varieties

Cristian Minoccheri, Xuanyu Pan

arXiv:1711.09987

<http://arxiv.org/abs/1711.09987v1>

## Two questions about curves on positive varieties

For a smooth complex projective variety  $X$ , let

$$\mathrm{Griff}_1(X) = \mathrm{CH}_1(X)_{\mathrm{hom}} / \mathrm{CH}_1(X)_{\mathrm{alg}}.$$

The paper asks how the positivity of a Fano variety simplifies integral 1-cycles. Its first result answers the Griffiths-group question for weighted complete intersections; its second uses the stronger 2-Fano condition—positivity of both  $c_1(X)$  and  $\mathrm{ch}_2(X)$ —to control the whole first Chow group.

**Main theorem** (Theorems 1.2 and 1.3). *Let  $X_{d_1, \dots, d_c} \subset \mathbf{P}_{\mathbf{C}}(1, 1, 1, a_3, \dots, a_n)$  be a smooth weighted complete intersection of dimension at least 3, not a standard projective space. If its index satisfies  $\iota_X > \frac{1}{2} \dim X$ , then  $\mathrm{Griff}_1(X) = 0$ .*

*Now let  $X$  be smooth, 2-Fano, and non-linear. Then  $\mathrm{CH}_1(X)$  is generated by lines in each of the following cases: (i)  $X$  is such a weighted complete intersection with  $\sum d_i \leq \sum_{j=3}^n a_j$ ; (ii)  $X \subset \mathbf{P}^n$  is conic-connected and  $\iota_X \geq (n+4)/2$ ; or (iii)  $X \subset \mathbf{P}^n$  has Picard number one and  $\iota_X \geq (3 \dim X + 5)/4$ . If its Fano variety of lines is irreducible, then  $\mathrm{CH}_1(X) \simeq \mathbf{Z}$ .*

## First mechanism: break rational curves into lines

For the weighted complete intersection, topology gives  $H_2(X, \mathbf{Z}) \simeq \mathbf{Z}\alpha$ , where  $\alpha$  is the degree-one curve class, and a previous connectedness theorem says that every geometric fiber of

$$\mathrm{ev}_1 : \overline{M}_{0,1}(X, \alpha) \longrightarrow X$$

is nonempty and connected. By the rational-curve generation theorem used in Section 2,  $\mathrm{CH}_1(X)$  is generated by rational curves. If a rational curve has class  $m\alpha$ ,  $m \geq 2$ , then the expected dimension of maps through two fixed points is

$$\chi(\mathbf{P}^1, T_X(-2)) = m \left( \sum a_i - \sum d_j \right) - (n - c) \geq 2.$$

Bend-and-break makes it algebraically equivalent to a connected union of lower-degree curves; iteration leaves a chain of lines. Connected fibers of  $\mathrm{ev}_1$  make two meeting lines algebraically equivalent. Thus homologically trivial combinations are algebraically trivial, proving Theorem 2.3 and hence Theorem 1.2. The strict index bound is precisely what supplies the two-dimensional deformation needed for bend-and-break.

## Second mechanism: broken conics form a Fano fiber

Let  $X$  be smooth, 2-Fano, non-linear and broken-conic-connected. For general  $p, q \in X$ , let  $F$  be the fiber over  $(p, q)$  of  $\mathrm{ev}_2 : \overline{M}_{0,2}(X, 2\alpha) \rightarrow X \times X$ , and let  $\Delta \subset F$  be the boundary of two lines meeting in one point. Freeness and generic smoothness make  $F$  and  $\Delta$  smooth. For the universal conic  $\pi : U \rightarrow F$  with map  $g : U \rightarrow X$ , Grothendieck–Riemann–Roch gives the decisive formula

$$-K_F = \pi_* g^* \mathrm{ch}_2(\Omega_X) + \Delta \quad (\text{Proposition 3.5}).$$

Adjunction cancels the boundary term:

$$-K_\Delta = (\pi_* g^* \mathrm{ch}_2(\Omega_X))|_\Delta.$$

Pushforward preserves ampleness here, so positivity of  $\mathrm{ch}_2(X)$  makes every connected  $\Delta$  Fano, hence rationally connected.

The formula packages the technical heart of Section 3. The identities  $c_1(\omega_{U/F}) = -\sigma_0 - \sigma_1$  and  $R^i \pi_* N_{U/X}(-\sigma_0 - \sigma_1) = 0$  for  $i > 0$  reduce GRR to degree two; the nodal locus pushes forward to  $\Delta$ , and the remaining relative-dualizing terms cancel. Adjunction then removes exactly the boundary contribution, leaving the positive  $\text{ch}_2$  term on  $\Delta$ .

Varying the second endpoint produces a family  $\phi : Y \rightarrow X$  of broken conics whose general fiber is  $\Delta$ . Given an irreducible curve  $C \subset X$  through a general point, the base change  $Y_C \rightarrow C$  has rationally connected general fiber and therefore a section by Graber–Harris–Starr. Inside the family of broken conics, the section is rationally equivalent to a sum of curves that either map to lines in  $X$  or lie in the fixed marked section and are contracted. Pushing forward gives  $[C]$  as an integral linear combination of lines (Theorem 4.4). Rational chain connectedness of the Fano scheme then identifies all line classes, giving  $\text{CH}_1(X) \simeq \mathbf{Z}$ .

In the stated applications, irreducibility of the Fano variety of lines supplies the missing connectivity input: the general pointed-line evaluation fiber is smooth and irreducible; its first Chern class is positive by the 2-Fano hypothesis. Graber–Harris–Starr then makes the pointed line space, and hence the line space, rationally connected. Thus Theorem 4.4’s rational-chain-connectedness hypothesis is actually available, rather than being an extra unstated assumption.

## Why the numerical hypotheses appear

The applications must establish both broken-conic-connectedness and connectedness of  $\Delta$ . For weighted complete intersections these follow from the low-degree inequalities and an ample boundary. In the high-index conic-connected case, one proof pulls the diagonal back through the two line-evaluation spaces and applies Fulton–Hansen connectedness when  $\iota_X \geq (n+4)/2$ . The alternative bound  $(3 \dim X + 5)/4$  uses the finite tangency map from  $F$  to the space of planes through the line  $\overline{pq}$  and rules out two disjoint boundary components; Picard number one plus this bound already implies conic-connectedness.

A useful model is a broken conic  $\ell_{pz} \cup \ell_{zq}$ : the moving node  $z$  records how two lines join the fixed endpoints. The proof needs the entire parameter space of such choices to be connected and positive, not merely the existence of one chain. That connectivity is the hardest technical bridge from individual rational curves to a statement in the integral Chow group. The paper proves generation, not a classification, unless the line space is irreducible; the 2-Fano and index assumptions are substantive and are not claimed for arbitrary Fano varieties. For example, a smooth degree- $r$  cover of  $\mathbf{P}^{n-1}$  branched along a hypersurface of degree  $ar$  satisfies line generation when  $n \geq a^2(r^2 - 1)$ ; for a general such cover the irreducibility input upgrades the answer to  $\text{CH}_1(X) \simeq \mathbf{Z}$  (Corollary 4.6).