

Zariski density of points with maximal arithmetic degree

Kaoru Sano, Takahiro Shibata

arXiv:2007.15180

<http://arxiv.org/abs/2007.15180v1>

Many maximal-growth orbits, not just one

Let X be a projective variety over a number field K , let H be ample, and let $f : X \rightarrow X$ be a surjective morphism. The first dynamical degree

$$\delta_f = \lim_{n \rightarrow \infty} ((f^n)^* H \cdot H^{\dim X - 1})^{1/n}$$

measures geometric degree growth, while the arithmetic degree $\alpha_f(x)$ measures the exponential growth of an ample height along the forward orbit $O_f(x)$. One always has $\alpha_f(x) \leq \delta_f$.

The paper asks for much more than a single point attaining equality. It calls (X, f) rich in maximal-degree points over a field L if there is a Zariski-dense $S \subset X(L)$ such that $\alpha_f(x) = \delta_f$ for every $x \in S$ and the orbits $O_f(x)$ are pairwise disjoint.

Main theorem (Sano–Shibata, Theorems 1.8, 1.11, and 1.12). *Assume $\delta_f > 1$.*

- (i) *For every projective X and surjective f , there are densely many $\overline{K}$ -points of maximal arithmetic degree with pairwise disjoint forward orbits.*
- (ii) *If X is unirational, they may be chosen in $X(K)$.*
- (iii) *If $X = A$ is an abelian variety, they may be chosen over one finite extension L/K .*

Part (i) is striking because $X(\overline{K})$ is countable: density must be manufactured by an induction that also avoids every previously chosen orbit. Parts (ii)–(iii) answer the stronger bounded-field question for two basic classes of potentially dense varieties.

Ingredient I: a nef eigenclass detects maximal growth

After normalization, Theorem 2.1 supplies a nef, numerically nontrivial $\mathbf{R}$ -Cartier divisor D with

$$f^* D \sim_{\mathbf{R}} \delta_f D. \quad (1)$$

Its canonical height is the convergent limit

$$\widehat{h}_{D,f}(x) = \lim_{n \rightarrow \infty} \frac{h_D(f^n(x))}{\delta_f^n}, \quad \widehat{h}_{D,f} \circ f = \delta_f \widehat{h}_{D,f}. \quad (2)$$

Positivity of (2) gives $\alpha_f(x) \geq \delta_f$; the general upper bound then forces equality. Thus the construction only has to find points of positive canonical height while keeping their orbits apart.

Ingredient II: large prime field degrees are dense

The arithmetic engine is Lemma 3.3. For any $A > 0$ and $B \in \mathbf{R}$, the set

$$T(D, A, B) = \{x \in X(\overline{K}) : [K(x) : K] \text{ has a prime factor } p > A, h_D(x) > B\} \quad (3)$$

is Zariski dense. To prove this, intersect with sufficiently general very ample divisors, normalize the resulting curve, and map it finitely to $\mathbf{P}^1$. These reductions preserve a large prime divisor of the field degree up to bounded factors. On $\mathbf{P}^1$, choose an element of prime degree $p > A$ over $\mathbf{Q}$ with $p > [K : \mathbf{Q}]$; infinitely many points of that degree have arbitrarily large height by Northcott finiteness.

This lemma is the proof's unusual bridge: positivity is an archimedean-looking height condition, whereas orbit separation is certified by an elementary divisibility obstruction in residue-field degrees.

The general induction

Enumerate the proper closed subsets of $X_{\overline{K}}$ as $Y_1, Y_2, \dots$. Suppose $x_1, \dots, x_k$ have already been chosen. Take A larger than every $[K(x_i) : K]$ and than the uniformly bounded degrees

$$[K(x) : K(f(x))]. \quad (4)$$

Apply Lemma 3.3 with $\widehat{h}_{D,f}$ itself as a height associated to D . It produces $x \notin Y_{k+1}$ with $\widehat{h}_{D,f}(x) > 0$ and with a prime $p > A$ dividing $[K(x) : K]$.

If $f^N(x) = f^M(x_i)$, the tower formula gives

$$[K(x) : K] = \prod_{j=0}^{N-1} [K(f^j(x)) : K(f^{j+1}(x))] [K(f^M(x_i)) : K]. \quad (5)$$

Every factor on the right is at most A , so none is divisible by p , whereas the left side is. This contradiction separates the new orbit from all old ones. Induction makes the points avoid Y_i one by one, hence makes their set Zariski dense. Equations (1)–(2) give maximal arithmetic degree.

Keeping all points over one number field

The prime-degree trick deliberately enlarges residue fields and therefore cannot prove (ii). On a unirational variety, the authors instead find, outside any prescribed closed set, a morphism $g : \mathbf{P}_K^1 \rightarrow X$ with g^*D ample. Along this curve,

$$\widehat{h}_{D,f}(g(a)) = rh(a) + O(1), \quad r > 0. \quad (6)$$

Canonical heights on each previously chosen orbit form a geometric progression with ratio δ_f . Counting K -points of bounded height on $\mathbf{P}^1$ finds a value of (6) strictly between two consecutive forbidden heights; the scaling law in (2) then keeps its whole orbit disjoint.

For an abelian variety, write $f = \tau_a \circ \phi$ and split, up to the addition isogeny $\mu : A_1 \times A_2 \rightarrow A$, so that $1 - \phi_1$ is surjective and $\delta_{\phi_2} = 1$. Write $a = a_1 + a_2$ with $a_i \in A_i$, and choose $b \in A_1$ satisfying $b - \phi_1(b) = a_1$. After one finite extension L/K , choose $x \in A_1(L)$ with $\mathbf{Z}_{>0}x$ dense and allow $y \in A_2(L)$. Points

$$z = \mu(\ell x + b, y) = \ell x + b + y \quad (7)$$

have arithmetic degree δ_f . Faltings finiteness lets multiples ℓx avoid any proper closed subset of A_1 , while Lemma 5.2 chooses ℓ so that $(\ell/\ell_i)^2 \notin \delta_f^{\mathbf{Z}}$. For a closed $Y \subsetneq A$, put $\tilde{Y} = \mu^{-1}(Y)$ and $V = \text{pr}_1(\tilde{Y})$. If $V \neq A_1$, apply Faltings to the proper translate $V - b$ and choose $\ell x \notin V - b$, equivalently $\ell x + b \notin V$; if $V = A_1$, use the density of $A_2(L)$ to choose y outside the fiber of $\tilde{Y}$ over $\ell x + b$. Applying the quadratic canonical height to a hypothetical orbit collision then gives $(\ell/\ell_i)^2 = \delta_f^{m-n}$, precisely the excluded equality. This replaces field-degree divisibility by height-value incommensurability.

Two further contribution lines

Theorem 6.1 treats a $\mathbf{P}^1$ -bundle $\pi : X \rightarrow C$ over an elliptic curve: every surjective f with $\delta_f > 1$ has the same dense, orbit-disjoint set over a finite extension. For the induced $g : C \rightarrow C$, the case $\delta_f = \delta_g$ lifts points supplied by Theorem 1.12. If $\delta_f > \delta_g$, a decomposable bundle has a nef eigenclass positive on the fibers and Theorem 4.1 applies; an indecomposable bundle becomes decomposable after finite base change, followed by finite descent (Lemma 2.3).

Appendix Theorem A.1 moves beyond morphisms: for a dominant rational $f : \mathbf{A}_K^2 \dashrightarrow \mathbf{A}_K^2$ with $\delta_f > 1$, the same finite-extension conclusion holds if some iterate is algebraically stable or if f is quadratic. Its proof constructs dense nonarchimedean regions where a local escape rate is positive and uses valuation classes to separate orbits; the quadratic case reduces, via its normal forms, either to the stable case or to analogous invariant regions.

Toy model, hard point, and scope

For $f([u : v]) = [u^d : v^d]$ on $\mathbf{P}^1$, the usual canonical height satisfies $\widehat{h}(f(x)) = d\widehat{h}(x)$. Choosing positive heights in distinct $d^{\mathbf{Z}}$ -classes is the one-dimensional shadow of (6)–(7): equal iterates would force the heights into the same class. The general theorem is harder because a nef eigenheight need not be ample or have Northcott's property; Lemma 3.3 supplies positivity without assuming it does.

The main projective-variety theorems concern morphisms with $\delta_f > 1$; Appendix A covers only the stated special affine-plane rational maps. The paper does not solve the bounded-field question for arbitrary potentially dense varieties or arbitrary dominant rational self-maps, and $\delta_f = 1$ is explicitly set aside. Its strength is the simultaneous conclusion of density, maximal growth, and pairwise orbit disjointness.