

Exponential equidistribution of periodic points for endomorphisms of projective space

Henry de Thelin, Tien-Cuong Dinh, Lucas Kaufmann

arXiv:2409.19787

<http://arxiv.org/abs/2409.19787v2>

Periodic points with an exponential error term

Let $f: \mathbb{P}^k \rightarrow \mathbb{P}^k$ be a holomorphic endomorphism of algebraic degree $d \geq 2$. Its Green current is

$$T = \lim_{m \rightarrow \infty} d^{-m} (f^m)^* \omega_{\text{FS}}.$$

It has a Hölder potential, and the equilibrium measure and small Julia set are

$$\mu = T^k, \quad J_k = \text{supp}(\mu).$$

Write $P_n = \{a : f^n(a) = a\}$; “period n ” here need not mean least period. For $0 < \gamma < 1$, let

$$P_{n,\gamma} = \left\{ a \in P_n \cap J_k : \|Df^n(a)^{-1}\| \leq d^{-(1-\gamma)n/2} \right\}. \quad (1)$$

Main theorem (de Thelin–Dinh–Kaufmann, Theorem 1.1). *For every $0 < \gamma < 1$ and $0 < \alpha \leq 1$, there are $0 < \xi < 1$ (independent of α) and $A_\alpha > 0$ such that, for every $P_{n,\gamma} \subseteq Q_n \subseteq P_n$ and every $\varphi \in C^\alpha(\mathbb{P}^k)$,*

$$\left| \left\langle d^{-kn} \sum_{a \in Q_n} \delta_a - \mu, \varphi \right\rangle \right| \leq A_\alpha \xi^{\alpha n} \|\varphi\|_{C^\alpha}. \quad (2)$$

The points may be counted with or without multiplicity.

Thus almost all of the roughly d^{kn} fixed points of f^n lie in J_k , are quantitatively repelling, and already sample μ with an exponential error. Corollary 1.2 makes the first assertion precise: the numbers of non-repelling points and of points outside J_k are both $O(\xi^n d^{kn})$ for some $\xi < 1$.

Ingredient I: deleting the postcritical danger zone

Let $PC_m = \bigcup_{j=1}^m f^j(C_f)$. Its degree grows exponentially, so mere non-charging of analytic sets by μ is insufficient. The paper proves the uniform tubular estimate (Proposition 3.1): if ν is a Monge–Ampère probability measure $\nu = (\omega_{\text{FS}} + dd^c u)^k$ with u an ω_{FS} -plurisubharmonic Hölder potential, then for some $\kappa \geq 1$ and every proper subvariety $V \subset \mathbb{P}^k$,

$$\deg V \leq \delta \implies \nu(\text{Tub}(V; \delta^{-\kappa})) \leq \delta^{-1}. \quad (3)$$

Its proof is the main potential-theoretic difficulty. A central projection turns a hypersurface away from its discriminant into finitely many bounded holomorphic graphs; local Monge–Ampère estimates control tubes around the graphs, while induction on the projection dimension controls the discriminant. Applied to $PC_{\lfloor cn \rfloor}$ for a suitably small $c > 0$, (3) discards an exponentially small μ -mass while leaving cells on which inverse branches are unramified.

Ingredient II: many small inverse branches

The authors cover projective space by translated cubical “Manhattans” of side $r = d^{-c'n}$ for another suitably small $c' > 0$. The translations are chosen so that their thin street networks have μ -mass $O(r)$ and all cell centres avoid the full postcritical set (Lemma 4.2). Quantitative equidistribution of preimages (Theorem 4.5) then compares the number of preimages entering each shrunken cell with its μ -mass.

Away from PC_ℓ , Proposition 4.4 supplies at least

$$(1 - A_3 d^{-\ell}) d^{km}$$

inverse branches of f^m on a smaller ball, with image diameter $O(d^{-(m-\ell)/2})$. The area bound on inverse images of complex lines produces the diameter estimate; Sibony–Wong extension makes it valid on the full ball. Splitting n into fixed long and short proportions, Sections 5.1–5.3 compose these branches through “nice” cells. The auxiliary proportions and exponents are chosen small relative to γ and to κ in (3). A double-counting argument shows that unsafe cells have exponentially small total μ -mass.

From inverse branches to repelling cycles

For each safe cell, Lemma 5.3 gives almost the μ -predicted number of order- n inverse branches mapping the cell strictly inside itself. The Kobayashi metric makes every such branch g a strict contraction (Lemma 5.4), hence g has a unique attracting fixed point. That point is fixed by f^n and repelling for f ; the diameter control gives (1). Starting the inverse iteration in J_k and using $f^{-1}(J_k) = J_k$ keeps the fixed point in J_k . This proves Proposition 5.5.

Summing a nonnegative Lipschitz test function over the points obtained cell by cell yields the one-sided estimate (Lemma 6.1)

$$d^{-kn} \sum_{a \in P_{n,\gamma}} \varphi(a) \geq \int \varphi d\mu - O(\eta^n) \quad (0 < \eta < 1). \quad (4)$$

Taking $\varphi = 1$ and using $\#P_n = d^{kn} + O(d^{(k-1)n})$ shows that the omitted fixed points are exponentially negligible. After normalizing a nonnegative test function so that $0 \leq \varphi \leq 1$, applying (4) to $1 - \varphi$ supplies the opposite inequality. Rescaling and decomposing a general Lipschitz test function then gives the two-sided bound; adding or deleting the negligible set handles every intermediate Q_n . Interpolation between Hölder spaces turns the Lipschitz estimate into (2).

The power-map model and the scope

For $f(z) = z^d$ on $\mathbb{P}^1$, the small Julia set is the unit circle, and the nonzero finite solutions of $f^n(z) = z$ are the $(d^n - 1)$ st roots of unity. Their normalized counting measures approach Haar measure exponentially; each root is obtained by closing a contracting inverse branch. The general proof recreates this picture without a group law, while paying for critical images, distorted branches, and uneven mass across cells.

The result is complex-analytic and concerns holomorphic endomorphisms of projective space, not rational self-maps with indeterminacy. Its exponential constant is not claimed optimal: Conjectures 6.2–6.3 predict the sharper rate λ^{-n} for every $1 < \lambda < d^{1/2}$.