

Symplectic resolutions for Higgs moduli spaces

Andrea Tirelli

arXiv:1701.07468

<http://arxiv.org/abs/1701.07468v1>

Which singular Higgs moduli spaces can be resolved?

Let X be a smooth projective complex curve of genus g , and let $\mathcal{M}_H(X, n)$ denote the moduli space of S-equivalence classes of semistable rank- n , degree-zero Higgs bundles (E, Φ) . Thus $\Phi \in H^0(X, \text{End}(E) \otimes \Omega_X^1)$, and stability is tested only on Φ -invariant subbundles. We work in the singular regime $g \geq 1$ and $n \geq 2$ implicit in the paper's classification.

A normal variety is a symplectic singularity if the symplectic form on its smooth locus extends holomorphically to every resolution. A resolution is symplectic when the extended form is nondegenerate.

Main theorem (Tirelli, Theorems 4.6, 4.7, and 4.9). *The space $\mathcal{M}_H(X, n)$ is a symplectic singularity. It admits a projective symplectic resolution exactly in the following cases:*

$$g = 1, \quad \text{or} \quad (g, n) = (2, 2). \quad (1)$$

For $g = 1$ the resolution is the Hilbert–Chow morphism

$$\text{Hilb}^n(T^*X) \longrightarrow \text{Sym}^n(T^*X) \simeq \mathcal{M}_H(X, n). \quad (2)$$

The exceptional $(2, 2)$ resolution is the one constructed earlier by Kiem–Yoo. The paper's contribution is to transfer both the general local obstruction and the symplectic-singularity property from character varieties, and to identify the elliptic construction.

Ingredient I: formal nonabelian Hodge theory

Three moduli spaces occur: the character variety $\mathcal{M}_B(X, n)$ of semisimple representations $\pi_1(X) \rightarrow \text{GL}_n(\mathbb{C})$, the de Rham moduli space $\mathcal{M}_{dR}(X, n)$ of flat bundles, and $\mathcal{M}_H(X, n)$. Nonabelian Hodge theory gives an analytic isomorphism $\mathcal{M}_B \simeq \mathcal{M}_{dR}$ and a topological homeomorphism $\mathcal{M}_{dR} \simeq \mathcal{M}_H$.

The stronger local input is Simpson's Isosingularity Theorem: at corresponding de Rham and Higgs points, the formal completions are canonically isomorphic. Analytification does not alter the completed local ring, so the analytic Betti–de Rham isomorphism yields, at corresponding points b and h ,

$$\widehat{\mathcal{M}}_{B,b} \simeq \widehat{\mathcal{M}}_{H,h}. \quad (3)$$

Artin approximation upgrades the required consequences of (3) to étale local statements. This is the decisive bridge: normality, rational Gorenstein singularities, and the presence or absence of a local symplectic resolution can be transported from the character side.

Ingredient II: prove the Higgs space is symplectic singular

Bellamy–Schedler prove that $\mathcal{M}_B(X, n)$ is a symplectic singularity. By (3), its normal and rational Gorenstein local rings transfer to $\mathcal{M}_H(X, n)$. Namikawa's criterion then reduces the remaining work to producing an everywhere nondegenerate closed holomorphic two-form on the smooth locus.

The stable Higgs locus already has its standard hyperkahler, hence holomorphic symplectic, form. Tirelli shows that its complement inside the smooth locus has codimension at least two. Indeed, a strictly polystable object is a direct sum of stable objects with ranks given by a nontrivial partition $n = n_1 + \cdots + n_k$. Since

$$\dim \mathcal{M}_H(X, r) = 2r^2(g - 1) + 2, \quad (4)$$

the direct-sum strata have dimension at most $\dim \mathcal{M}_H(X, n) - 2$. The symplectic form therefore extends across the missing codimension-two set. Namikawa's criterion proves the symplectic-singularity assertion for $g \geq 2$; the elliptic case follows from the explicit symmetric-product model below.

Ingredient III: local terminality forbids a resolution

Assume $g > 1$ and $(g, n) \neq (2, 2)$. Choose the Higgs point corresponding, under nonabelian Hodge theory, to the trivial representation. If a symplectic resolution of $\mathcal{M}_H(X, n)$ existed, restriction to an étale neighbourhood and (3) would produce one for the corresponding neighbourhood of the trivial point in $\mathcal{M}_B(X, n)$.

Bellamy–Schedler’s analysis makes that Betti neighbourhood factorial, while Namikawa supplies terminality. A symplectic resolution is crepant, whereas a factorial terminal singularity admits no crepant resolution. The assumed Higgs resolution is therefore impossible. The force of the argument is its locality: no global algebraic identification between Betti and Higgs moduli spaces is needed; their completed singularities suffice.

Ingredient IV: the two exceptional constructions

For an elliptic curve, a theorem quoted from Franco and Franco–Garcia-Prada–Newstead identifies

$$\mathcal{M}_H(X, n, d) \simeq \mathrm{Sym}^h(T^*X), \quad h = \gcd(n, d). \quad (5)$$

At degree zero, $h = n$. The cotangent bundle T^*X is a smooth symplectic surface, and for any smooth symplectic surface S the symmetric power $\mathrm{Sym}^n S$ is a symplectic singularity resolved symplectically by $\mathrm{Hilb}^n S$. This gives (2). The remaining exception $(g, n) = (2, 2)$ is not reconstructed in this paper: existence is imported from Kiem–Yoo, while the general nonexistence theorem deliberately excludes it.

A two-point model

Take $n = 2$ and $g = 1$. Formula (5) turns a polystable Higgs bundle into an unordered pair of points of the symplectic surface T^*X . Away from the diagonal the quotient $(T^*X)^2/\mathfrak{S}_2$ is smooth. On the diagonal, the two points collide and create the symmetric-product singularity. $\mathrm{Hilb}^2(T^*X)$ remembers the length-two subscheme, including its infinitesimal collision direction, and the Hilbert–Chow map resolves exactly that defect. This model explains the positive elliptic case but not the higher-genus obstruction, which is detected instead by the terminal local ring at the trivial representation.

Hard point, scope, and omissions

The conceptual core is the formal transfer (3). The Betti and Higgs spaces are not algebraically isomorphic globally, yet their completed local rings match pointwise; this is just strong enough to transport singularity type and a local obstruction to crepant resolution. The codimension estimate is then what extends the stable-locus two-form to the entire smooth locus.

The statement above is restricted to the rank- $n \geq 2$, genus- $g \geq 1$ singular setting; rank one and genus zero are smooth or degenerate cases and must not be read into (1). The deepest local factoriality input comes from Bellamy–Schedler and terminality from Namikawa, while the $(2, 2)$ construction is an input from Kiem–Yoo. The paper also notes that arbitrary-degree Higgs spaces and twisted character varieties lack the formal correspondence needed by this argument, so the same classification is not asserted there.