

Uniform bounds on periodic points of polynomials with good reduction

Isaac Rajagopal, Robin Zhang

arXiv:2510.26119

<https://arxiv.org/abs/2510.26119>

One good place forces a global bound

Let K be a number field of degree D , let $\mathfrak{p} \subset \mathcal{O}_K$ lie over the rational prime p , and put $\text{Per}_K(\phi) = \{x \in K : \phi^n(x) = x \text{ for some } n > 0\}$. For a monic degree- d polynomial $\phi(X) = \sum_{i=0}^d a_i X^i$, good reduction at $\mathfrak{p}$ means $v_{\mathfrak{p}}(a_i) \geq 0$ for every i and $v_{\mathfrak{p}}(a_d) = 0$. It is *inseparable good reduction* if, in addition,

$$a_i \in \mathfrak{p} \quad \text{whenever } p \nmid i.$$

(The definition in the paper allows a $\mathfrak{p}$ -unit leading coefficient; the polynomial is normalized to be monic.) In particular $p \mid d$.

Main theorem (Theorem 3). *If $\phi \in K[X]$ has inseparable good reduction at $\mathfrak{p}$, then*

$$\#\text{Per}_K(\phi) \leq N_{K/\mathbf{Q}}(\mathfrak{p}) \leq d^D.$$

Thus one place of this special good reduction gives an unconditional bound independent of the number of bad places. In particular it applies to $X^d + c$ whenever c is integral at a prime above some $p \mid d$.

The result is a concrete case of dynamical uniform boundedness: the general conjecture asks for a bound depending only on the ambient dimension, d , and D , and is open even for quadratic maps over $\mathbf{Q}$.

The local separation proposition

Complete at $\mathfrak{p}$. Write $F = K_{\mathfrak{p}}$, let $\mathcal{O}_F$ have maximal ideal $\mathfrak{p}_F$, and let $\mathcal{O}_F/\mathfrak{p}_F \simeq \mathbf{F}_{p^f}$. Proposition 2.1 treats every polynomial

$$\phi(X) = \sum_{i=0}^d a_i X^i \in \mathcal{O}_F[X], \quad v_F(a_d) = 0, \quad v_F(a_i) > 0 \text{ if } p \nmid i,$$

and proves $\#\text{Per}_F(\phi) \leq p^f$.

There are two complementary estimates.

- 1. Escape outside the unit ball (Lemma 2.2).** If $|\lambda|_F > 1$, the leading term strictly dominates all the others, so $|\phi(\lambda)|_F = |a_d \lambda^d|_F > |\lambda|_F$. The absolute values of successive iterates increase strictly; hence λ cannot be periodic.
- 2. Contraction inside one residue class (Lemma 2.3).** Let λ be periodic and $0 < |x - \lambda|_F < 1$. At any step, let μ be the corresponding point on the orbit of λ and let y be the corresponding iterate of x ; expand $\phi(y)$ around μ . The linear coefficient is strictly small: every summand $a_i y^{i-1}$ lies in $\mathfrak{p}_F$, using $a_i \in \mathfrak{p}_F$ when $p \nmid i$ and $i \in \mathfrak{p}_F$ when $p \mid i$. The higher Taylor coefficients need only be integral; for $k \geq 2$, the extra power $|y - \mu|_F^k < |y - \mu|_F$ supplies strictness. Consequently

$$0 < |\phi^n(x) - \phi^n(\lambda)|_F < |\phi^{n-1}(x) - \phi^{n-1}(\lambda)|_F < \cdots < |x - \lambda|_F.$$

If two iterates ever coincide, periodicity already forces $x = \lambda$; otherwise the strict decrease prevents x from being periodic.

Thus all periodic points lie in $\mathcal{O}_F$, and no two have the same image in $\mathbf{F}_{p^f}$. Reduction is injective on $\text{Per}_F(\phi)$, proving the local bound p^f .

Back to the number field

The embedding $K \hookrightarrow K_{\mathfrak{p}}$ sends K -rational periodic points to local periodic points. Hence

$$\#\mathrm{Per}_K(\phi) \leq p^f = N_{K/\mathbf{Q}}(\mathfrak{p}).$$

Since $f \leq D$ and $p \mid d$, one has $p^f \leq p^D \leq d^D$. The proof is therefore entirely local after choosing $\mathfrak{p}$, but its conclusion is uniform over all degree- D number fields.

The picture behind the argument is especially transparent for $\phi(X) = X^2 - 1$ over $\mathbf{Q}_2(\sqrt{5})$. Its four periodic points occupy the four residue classes of $\mathbf{F}_4$: nearby points contract toward the corresponding cycle, whereas points outside the integral disk escape. This models both inequalities above, although the theorem neither asserts that all residue classes contain periodic points nor classifies their exact periods.

Under the stronger prime-power hypotheses of Theorem 5, the authors do obtain exact-period restrictions by computing ϕ^n modulo a uniformizer and using Hensel lifting. That refinement is separate from the main counting argument. The hardest step in the bound is the strict Taylor contraction uniformly along an entire cycle; it is precisely where the inseparability condition is used.