

Singularities of moduli spaces of sheaves on K3 surfaces and Nakajima quiver varieties

Enrico Arbarello, Giulia Saccà

arXiv:1505.00759

<http://arxiv.org/abs/1505.00759v5>

K3-sheaf singularities are quiver singularities

Let S be a complex projective K3 surface, v a primitive Mukai vector of a pure one-dimensional sheaf, and H_0 a polarization on a v -wall. A singular point $x \in M_{H_0}(v)$ is represented by a unique H_0 -polystable sheaf

$$F = \bigoplus_{i=1}^s F_i \otimes V_i, \quad n_i = \dim V_i, \quad G = \operatorname{Aut}(F) = \prod_i \operatorname{GL}(V_i), \quad (1)$$

with pairwise nonisomorphic stable factors F_i .

Build a quiver $Q(F)$ with one vertex for each F_i , $\dim \operatorname{Ext}^1(F_i, F_j)$ edges between distinct vertices, and $\frac{1}{2} \dim \operatorname{Ext}^1(F_i, F_i)$ loops at vertex i . For the dimension vector $\mathbf{n} = (n_i)$, let $\mathfrak{M}_0 = \mu^{-1}(0) // G$ be the affine Nakajima quiver variety for the doubled quiver.

Main theorem (Arbarello–Saccà, Theorems 1.1 and 6.5). *There is an analytic isomorphism of germs*

$$(\mathfrak{M}_0, 0) \simeq (M_{H_0}(v), x). \quad (2)$$

For every chamber whose closure contains H_0 , and H in that chamber, a quiver-GIT character χ_H can be chosen so that the symplectic resolutions

$$\mathfrak{M}_{\chi_H}(\mathbf{n}) \longrightarrow \mathfrak{M}_0, \quad M_H(v) \longrightarrow M_{H_0}(v)$$

correspond locally through (2). After normalizing $H \cdot D = H_0 \cdot D$, where $D = c_1(F)$, one may take

$$\chi_H((g_i)) = \prod_{i=1}^s \det(g_i)^{D_i \cdot H - D_i \cdot H_0}, \quad D_i = c_1(F_i). \quad (3)$$

The result gives both a normal form for every such singularity and a dictionary between two wall-crossing theories: variation of sheaf polarization and variation of quiver GIT.

Ingredient I: making the Kuranishi space quadratic

The deformation space of F is cut out in $\operatorname{Ext}^1(F, F)$ by a Kuranishi map whose quadratic term is the cup product

$$\kappa_2(e) = e \smile e \in \operatorname{Ext}^2(F, F)_0. \quad (4)$$

Theorem 3.7 proves the quadraticity statement $\operatorname{Def}_F \simeq \kappa_2^{-1}(0)$, but importantly does not claim that every pure one-dimensional F is itself formal. After twisting so that F is globally generated with vanishing H^1 , its Lazarsfeld–Mukai bundle M_F is the kernel of the evaluation map. Proposition 3.5 identifies the deformation functors of F and M_F equivariantly, while Proposition 3.6 is the resulting quadraticity criterion once M_F is polystable. A Fourier–Mukai argument using Yoshioka’s results proves M_F is H_0 -polystable (Theorem 3.8), and Zhang’s positive-rank theorem makes $R\operatorname{Hom}(M_F, M_F)$ formal. Thus its Kuranishi space, and hence that of F , has only the quadratic equation (4).

Formal quadraticity alone is insufficient for taking the quotient by G . Proposition 4.4 is applied to the formal bundle M_F : it combines an equivariant formal Kuranishi family, Rim’s uniqueness, and analytic approximation to obtain a local G -equivariant analytic isomorphism with $\kappa_2^{-1}(0)$. Proposition 3.5 transports this comparison back to F . This is one of the proof’s delicate bridges.

Ingredient II: the Ext-quiver identifies the moment map

Serre duality and the decomposition (1) identify

$$\operatorname{Rep}(\overline{Q}, \mathbf{n}) \simeq \operatorname{Ext}^1(F, F), \quad \mathfrak{g}(\mathbf{n})^\vee \simeq \operatorname{Ext}^2(F, F), \quad (5)$$

and Proposition 6.1 identifies κ_2 with the quiver moment map μ . For pure one-dimensional factors, $\dim \operatorname{Ext}^1(F_i, F_j) = D_i \cdot D_j$ for $i \neq j$; up to the vertex loops, Q is the dual graph of the support. Taking the local G -quotient of (5) proves (2).

Ingredient III: the two wall arrangements coincide

Put $d_i = H_0 \cdot D_i$ and $a_i = H \cdot D_i$. On the transverse slice $\sum_i n_i a_i = H_0 \cdot D$, Lemma 6.4 sends

$$(a_i)_i \mapsto \theta = (a_i - d_i)_i \in \mathbf{n}^\perp. \quad (6)$$

The equations of the sheaf walls, obtained from possible stable-factor decompositions, become exactly the root hyperplanes defining the quiver-GIT walls. If several D_i have the same cohomology class, the image is the corresponding symmetry-invariant subspace rather than all of $\mathbf{n}^\perp$; this does not affect the chamber matching asserted by the theorem.

Ingredient IV: matching the actual resolutions

Choose an étale slice Z in a Quot scheme through the point for F . The locus Z^H parametrizing H -semistable sheaves has a cartesian quotient square with $M_H(v) \rightarrow M_{H_0}(v)$ (Proposition 7.1). The main remaining issue is to recognize Z^H as a GIT semistable locus on the quiver side.

Determinant line bundles on the universal family carry natural linearizations. Le Potier's criterion and Hilbert–Mumford weights show that Z^H is a saturated part of their semistable locus (Proposition 7.10). Lemma 7.17 computes the stabilizer character; combining the H and H_0 determinant bundles kills the scalar center and produces precisely (3), or its normalized version before rescaling. Lemmas 7.18–7.19 then transport the semistable locus through the G -equivariant analytic isomorphism. Taking quotients makes the local square of resolutions cartesian, proving the second part of the theorem.

Why purity and the wall dictionary matter

For a pure one-dimensional factor, the Mukai vector is $v_i = (0, D_i, \chi_i)$ and stability is governed by $\mu_H(F_i) = \chi_i/(D_i \cdot H)$. Thus moving H changes only the intersection numbers $D_i \cdot H$, exactly the coordinates appearing in (6) and in the determinant character (3). This is why the variation of polarization can be read directly as quiver GIT. Proposition 6.3 makes the statement more precise: decompositions of $\mathbf{n}$ into positive roots label the local polystable strata, and their sheaf-wall equations map to the corresponding root hyperplanes in $\mathbf{n}^\perp$.

Purity is also what makes the modular morphism available. For a chamber adjacent to H_0 , changing the polarization gives a regular map $M_H(v) \rightarrow M_{H_0}(v)$; in the primitive case it is the smooth symplectic resolution appearing in the theorem. The local quiver model does more than identify the singular affine quotient: the cartesian diagrams in Section 7 identify the fibers of this particular modular resolution with those of the GIT quotient. The explicit character (3) is therefore not decoration; it records which wall-crossing resolution the chosen polarization produces.

A two-factor model and the hard point

Take $F = F_1 \oplus F_2$ and suppress vertex loops to isolate the interaction. If $D_1 \cdot D_2 = m$, the local model is the doubled m -arrow Kronecker quiver with dimension vector $(1, 1)$. Under the normalization in (6), a neighboring polarization gives $\theta = (r, -r)$; the two signs select the two adjacent GIT chambers. This displays how crossing the sheaf wall becomes variation of a quiver quotient. The zero character retains the singular semisimplification, whereas a nonzero generic character selects stable representations and separates the two directions of extension between the factors. The full theorem must additionally handle loops, multiplicities n_i , multiple walls, and the restriction imposed by the ample cone.

The hardest point is not writing down the Ext-quiver, but proving that the analytic germ and its modular resolution are simultaneously compatible with the group action and the stability linearization. The conclusion is local analytic near x , not a global isomorphism of the two moduli spaces. It uses pure one-dimensional sheaves on a K3 surface and a primitive Mukai vector in the introductory form; higher rank requires different stability technology, and the paper explicitly distinguishes quadraticity from the stronger formality assertion.