

Non-density of points of small arithmetic degrees

Yohsuke Matsuzawa, Sheng Meng, Takahiro Shibata, De-Qi Zhang

arXiv:2002.10976

<http://arxiv.org/abs/2002.10976v2>

From one orbit to a bounded-degree locus

Let X be a projective variety over a number field K and $f: X \rightarrow X$ a surjective endomorphism. For an ample divisor H , the first dynamical degree and the arithmetic degree are

$$\delta_f = \lim_{n \rightarrow \infty} ((f^n)^* H \cdot H^{\dim X - 1})^{1/n}, \quad \alpha_f(x) = \lim_{n \rightarrow \infty} h_H(f^n(x))^{1/n}.$$

The latter limit exists for such morphisms and is independent of H . Write

$$X(K, d) = \{x \in X(\overline{K}) : [K(x) : K] \leq d\}, \quad Z_f = \{x \in X(\overline{K}) : \alpha_f(x) < \delta_f\}, \quad Z_f(K, d) = Z_f \cap X(K, d).$$

Kawaguchi–Silverman predicts only that each $x \in Z_f(\overline{K})$ has a non-dense orbit. The paper’s *small Arithmetic Non-Density* conjecture (sAND) asks for the stronger collective statement that $Z_f(K, d)$ is not Zariski dense for every fixed $d > 0$. The degree bound is essential: for $z \mapsto z^2$ on $\mathbf{P}^1$, all iterated preimages of 1 have arithmetic degree 1 and together form a Zariski-dense subset over $\overline{K}$.

Main theorem (Theorem 1.5). *The sAND conjecture holds for every surjective endomorphism of X if, after base change to $\overline{K}$, one of the following applies:*

- (i) X is a Mori dream space;
- (ii) $\dim X \leq 2$;
- (iii) X is hyperkähler;
- (iv) X is abelian or $\mathbf{Q}$ -abelian;
- (v) X is smooth and rationally connected and admits an int-amplified endomorphism;
- (vi) X is a smooth threefold and admits an int-amplified endomorphism.

Here “admits” does not require the endomorphism under study to be int-amplified.

Thus the theorem covers three principal building blocks suggested by equivariant birational geometry—abelian, hyperkähler, and rationally connected varieties—but not general Calabi–Yau varieties.

The model mechanism: canonical height plus Northcott

If $f^*H \sim qH$ with H ample and $q > 1$, the canonical height satisfies $\widehat{h}_H \circ f = q\widehat{h}_H$ and $Z_f = \{\widehat{h}_H = 0\} = \text{Prep}(f)$. Northcott then makes $Z_f(K, d)$ finite. For a Mori dream space, after an iterate one instead finds a nonzero base-point-free divisor D with $f^*D \sim \delta_f D$; its canonical height vanishes on Z_f , and the resulting bounded-height locus is non-dense even though D need not be ample. This is the basic pattern throughout: construct an expanding height, identify its zero locus geometrically, and use the field-degree bound to prevent that locus from spreading everywhere.

The abelian calculation

The paper makes this pattern completely explicit. For an abelian variety A and a surjective endomorphism f with $\delta_f > 1$, Theorems 5.1 and 5.6 produce a proper abelian subvariety $B \subsetneq A_{\overline{K}}$ and $p \in A(\overline{K})$ such that

$$f(B + p) = B + p, \quad Z_f = B(\overline{K}) + p + \text{Tor } A(\overline{K}),$$

and, for every $d > 0$,

$$Z_f(K, d) \subseteq \bigcup_{i=1}^r (B(\overline{K}) + p + t_i) \quad (t_i \in \text{Tor } A(\overline{K})).$$

This finite union is proper, proving sAND.

For the proof, first suppose that f is an isogeny. A symmetric nef eigen-divisor $D \neq 0$ with $f^*D \equiv \delta_f D$ has a nonnegative canonical height whose zero set is $B + \text{Tor } A$. If $\delta_{f|B} < \delta_f$, this is already the small-degree locus; if equality holds, induction on $\dim A$ shrinks B further. For a general endomorphism, write $f = \tau_a \circ \varphi$ with φ an isogeny. The isogeny decomposition $A_1 \times A_2 \rightarrow A$ separates an expanding factor from a factor of dynamical degree 1, while a translation conjugates the expanding factor back to an isogeny. Pushing the preceding zero-locus description through the finite addition map gives the displayed translate $B + p$.

Finally quotient by B . The image of $Z_f(K, d) - p$ is bounded-degree torsion in A/B , hence finite by Northcott. It is therefore killed by one integer N , so its preimage lies in the finitely many cosets indexed by $A[N]$. This last step is where the field-degree bound becomes decisive.

How the remaining cases fit

For a positive-entropy surface automorphism, Theorem 4.1 describes Z_f as finitely many periodic curves together with periodic points, which are finite in bounded degree away from those curves. On a hyperkähler variety, Theorem 4.2 instead places Z_f inside a fixed proper effective locus together with periodic points. Non-invertible surface maps are handled by classification and reduction to polarized, abelian, product, or fibred cases (Theorem 4.3).

For varieties admitting an int-amplified map, an equivariant minimal model program transfers sAND through generically finite maps and contractions. The only residual configuration (Case TIR) would yield, after a quasi-étale cover, a positive-dimensional abelian base. It is excluded when the smooth-locus algebraic fundamental group is finite, and a separate boundary analysis shows that with terminal singularities it has relative dimension at least 3; hence it cannot occur for a threefold. These are exactly the inputs for parts (v) and (vi).

The paper also proves that sAND in dimension at most n is equivalent to containment of $Z_f(K, d)$ in finitely many f -preperiodic subvarieties of smaller dynamical degree, and relates relative sAND to uniform boundedness for preperiodic and torsion points. The hard point omitted here is the detailed equivariant-MMP exclusion of Case TIR.