

Dynamical degrees of twisted rational maps

Marc Abboud, Junyi Xie

arXiv:2608.09275

<http://arxiv.org/abs/2608.09275v1>

Dynamics after a field twist

Let K be a field. A twisted rational map $f_\tau: X \dashrightarrow X$ is a rational map of schemes for which the structural maps fit over a finite map $\tau: \operatorname{Spec} K \rightarrow \operatorname{Spec} K$; equivalently it is an ordinary K -rational map $X^{(\tau)} \dashrightarrow X$, where only the K -structure of the source has changed. This is forced on the generic fibre of a semiconjugacy $q \circ f = g \circ q$: over $K(Y)$, the map induced by f is twisted by the field endomorphism induced by g .

For a dominant twisted self-map of a projective d -fold and a big and nef divisor H , set

$$\deg_{H,i}(f_\tau^n) = ((f_\tau^n)^* H)^i \cdot H^{d-i}, \quad \lambda_i(f_\tau) = \lim_{n \rightarrow \infty} \deg_{H,i}(f_\tau^n)^{1/n}. \quad (1)$$

Main theorem (Abboud–Xie, Theorems A and B). *For every $1 \leq i \leq d$, the limit in (1) exists, is independent of H , and is invariant under twisted birational conjugacy.*

More generally, suppose dominant twisted rational maps on projective varieties satisfy

$$q_\sigma \circ f_\tau = g_\omega \circ q_\sigma, \quad q_\sigma: X \dashrightarrow Y \text{ dominant.}$$

Writing f_η for the induced twisted map on the generic fibre and $e = \dim X - \dim Y$, one has, for $1 \leq i \leq \dim X$,

$$\lambda_i(f_\tau) = \max_{\substack{a+b=i \\ 0 \leq a \leq e, \ 0 \leq b \leq \dim Y}} \lambda_a(f_\eta) \lambda_b(g_\omega). \quad (2)$$

For ordinary f, g and a dominant morphism q , moreover $\lambda_a(f|q) = \lambda_a(f_\eta)$; throughout, $\lambda_0 = 1$.

Thus the growth spectrum of the total dynamics is assembled exactly from the fibre and base spectra. In terms of $\mu_i(h) = \lambda_i(h)/\lambda_{i-1}(h)$, formula (2) says that the multipliers of f_τ are the decreasing rearrangement of those of f_η and g_ω .

Ingredient I: intersection theory survives twisting

The underlying scheme of $X^{(\tau)}$ is X , so Picard groups, cycles, and numerical classes are canonically identified. The subtlety is that degrees of zero-cycles depend on the structural morphism and acquire a factor $\deg(\tau)$ (Lemma 2.8). Proposition 2.16 preserves the relevant positivity cones under twisting, while Proposition 2.20 supplies the twisted composition estimate. Consequently the degree sequence is submultiplicative up to a uniform constant; Fekete's lemma gives (1), comparison of big and nef classes removes H , and the Khovanskii–Teissier inequalities give log-concavity $\lambda_i^2 \geq \lambda_{i-1} \lambda_{i+1}$.

Ingredient II: restrict the graph to the generic fibre

For Cartier divisor classes $D_1, \dots, D_e$ on X and $L_1, \dots, L_{\dim Y}$ on Y , Proposition 3.7 proves the intersection identity

$$\deg(L_1 \cdots L_{\dim Y}) D_{1,\eta} \cdots D_{e,\eta} = D_1 \cdots D_e q^* L_1 \cdots q^* L_{\dim Y}. \quad (3)$$

Restricting the normalized graph of f^n to the generic point of Y gives the graph of f_η^n . Formula (3) therefore identifies each relative degree (take the D_j from pullbacks of a big and nef H_X) with the corresponding generic-fibre degree times the fixed number $H_Y^{\dim Y}$. Taking n th roots removes that constant and proves $\lambda_a(f \mid q) = \lambda_a(f_\eta)$.

Ingredient III: the mixed upper bound is a matrix estimate

After resolving the graph of q_σ , write $L_n = (f_\tau^n)^* L$ and $M_n = (g_\omega^n)^* M$ (with M also pulled back to X). Siu's inequality and (3) give uniform recurrences for the mixed classes $L_n^{i-b} M_n^b$. For a sufficiently large block length, these form an upper-triangular matrix inequality whose diagonal entries approach

$$\lambda_{i-b}(f_\eta)^m \lambda_b(g_\omega)^m.$$

The exponential growth of the matrix is its largest diagonal growth rate, which yields the $\leq$ direction of (2). This uniform triangular estimate is the hard point: a bound depending on the iterate would not survive the n th-root limit.

For the reverse inequality, compute $\lambda_i(f_\tau)$ with the big and nef class $L + q_\sigma^* M$. Its intersection expansion has nonnegative terms. Keeping the term with a fibre and b base factors, then applying (3), gives

$$\deg_i(f_\tau^n) \geq C_{a,b} \deg_a(f_\eta^n) \deg_b(g_\omega^n).$$

Taking roots and maximizing proves $\geq$.

The projective-space model

If $\phi_\tau: \mathbb{P}^d \rightarrow \mathbb{P}^d$ is twisted and its underlying ordinary morphism ϕ has algebraic degree r , the twist changes coefficients but fixes the hyperplane class. Hence $(\phi_\tau^n)^* H = r^n H$ and $\lambda_i(\phi_\tau) = r^i$. This is the clean model for the general construction: numerical growth ignores coefficient twisting, whereas the full proof must control rational indeterminacy and mixed intersections.

Affine algebraicity and scope

Theorem C applies a second layer of the theory. If X_0 is smooth affine of dimension d in characteristic zero, $f_\tau: X_0 \rightarrow X_0$ is dominant, and $\lambda_1^2 > \lambda_2$, then λ_1 is algebraic of degree at most $(d-1)^2$ for $d \geq 3$ and at most d for $d \leq 2$; the bound is d in every dimension when $\text{QAlb}(X_0) = 0$.

In the latter case, Theorem 5.4 supplies a rank-one spectral limit $\lambda_1^{-n}(f_\tau^n)^* \alpha \rightarrow L(\alpha)\theta^*$. Theorem 8.8 shows that the functional L , restricted to divisors at infinity, satisfies the valuation axioms and therefore determines an eigenvaluation v^* . Its rational value group has dimension at most d by Abhyankar's inequality, so λ_1 is an eigenvalue of a rational matrix of size at most d . In general, logarithmic Iitaka and quasi-Albanese fibrations reduce dimension; (2) and the spectral gap decide whether the relevant eigenvalue comes from the fibre or the quasi-abelian base. These affine conclusions require characteristic zero and the strict gap, unlike Theorems A–B; no optimality of the quadratic degree bound is claimed.