

Singular Rational Curves on Elliptic K3 Surfaces

Jonas Baltes

arXiv:2111.07808

<http://arxiv.org/abs/2111.07808v1>

Two abundance theorems for elliptic K3 surfaces

Let $p : X \rightarrow \mathbb{P}^1$ be a complex projective elliptic K3 surface. A “rational curve” below means an integral curve with rational normalization; it may be highly singular. If R is such a curve, adjunction on X gives $p_a(R) = 1 + R^2/2$, so unbounded self-intersection is precisely unbounded arithmetic genus.

For a curve $C \subset X$ with normalization $\nu : \tilde{C} \rightarrow X$, the torsion-free image of $\nu^*\Omega_X \rightarrow \Omega_{\tilde{C}}^1$ is a line bundle and determines the lift $\tilde{C} \rightarrow \mathbb{P}(\Omega_X)$, where $\mathbb{P}(\Omega_X)$ parametrizes one-dimensional quotients.

Main theorem (Baltes, Theorems 1.1 and 1.2). *Every elliptic K3 surface $X \rightarrow \mathbb{P}^1$ contains rational curves $R_i \subset X$ with $R_i^2 \rightarrow \infty$. Moreover, the union of the lifts of rational curves to $\mathbb{P}(\Omega_X)$ is Zariski-dense.*

The first conclusion strengthens mere infinitude by forcing increasing singularity complexity. The second says that even the tangent directions of rational curves are abundant; Corollary 7.1 recovers, in the elliptic case, $H^0(X, \text{Sym}^n \Omega_X) = 0$ for every $n > 0$.

The obstruction: quasi-torsion

Write $U \subset \mathbb{P}^1$ for the smooth-fiber locus. On a simply connected disc $\Delta \subset U$, a Jacobian elliptic fibration has a standard analytic form

$$(\mathbb{C} \times \Delta)/(\mathbb{Z}\tau(t) + \mathbb{Z}, t).$$

An elementary quasi-torsion curve has local branches $z = a\tau(t) + b$ for constants $a, b \in \mathbb{R}$. For a multisection $C \subset X$, form the fiberwise difference locus

$$D(C) = d(C_U \times_U C_U) \subset J^0(X), \quad (1)$$

where $J^0(X)$ is the compactified relative Jacobian. The multisection is *quasi-torsion* when $D(C)$ is a finite union of elementary quasi-torsion curves.

This definition isolates exactly what prevents multiplication from making new intersections. Proposition 3.10 supplies the local criterion: a section avoiding torsion of every order in an infinite multiplicatively closed set must have constant real coordinates (a, b) , hence is elementary quasi-torsion. Contrapositively, Corollary 3.11 says that torsion points of the prescribed orders are dense on any irreducible curve in $J^0(X)$ that is not elementary quasi-torsion. Applied to a non-elementary component of $D(C)$, this yields pairs of points of C with torsion difference.

Theorem 4.1 constructs a rational non-quasi-torsion multisection, allowing a change of elliptic fibration. For a non-isotrivial fibration, Hassett’s non-torsion rational multisection and large monodromy combine with Proposition 3.8: an algebraic elementary quasi-torsion curve in $J^0(X)$ is actually torsion. In the isotrivial case with $c(X \rightarrow \mathbb{P}^1) > 0$, suitable multiplication first removes all components of $D(C)$ that are non-trivial torsion of order at most the monodromy constant κ_X . Proposition 4.7 can then apply Hurwitz to force the geometric genus to grow with fiber degree, contradicting rationality after passage to a suitable twist. The remaining exceptional isotrivial pencils are replaced by a non-isotrivial elliptic pencil.

Multiplication manufactures singularities

Let d_0 be the index of the fibration. Compactified Jacobians are again elliptic K3 surfaces, and fiberwise tensor powers define rational maps

$$g_n : X \simeq J^1(X) \dashrightarrow J^n(X). \quad (2)$$

Start with the rational non-quasi-torsion curve R . After replacing it by a suitable $g_{k_0}(R)$, its fiber degree is minimal among the degrees obtained for $k \equiv 1 \pmod{d_0}$. This minimality makes every relevant restriction $R \dashrightarrow g_k(R)$ birational: otherwise the image would have smaller fiber degree.

Now choose r disjoint analytic opens in U . In each open, the contrapositive local criterion gives a pair of points of R whose difference is an isolated torsion point, of some order $m_i \equiv 1 \pmod{d_0}$. For $m = \prod_i m_i$, multiplication identifies each pair, so $g_m(R)$ has at least r locally reducible singularities. The image remains rational because the map on R is birational. Letting $r \rightarrow \infty$, the singularity contributions to arithmetic genus grow; by adjunction the self-intersections tend to infinity. This proves the first theorem. The key point is not merely that multiplication gives many images, but that non-quasi-torsion supplies isolated coincidences in every chosen region.

Why the lifted curves are dense

Choose a line bundle M of fiber degree d_0 . For $n \equiv 1 \pmod{d_0}$, tensoring by $M^{-(n-1)/d_0}$ identifies $J^n(X)$ back with $J^1(X)$ and gives a self-rational map G_n . In a standard model it is simply

$$G_n(z, t) = (nz, t). \quad (3)$$

For a local branch $C = (f(t), t)$ of a non-quasi-torsion curve, density of torsion differences produces points t_j and $n_j \equiv 1 \pmod{d_0}$ with

$$(n_j - 1)f(t_j) = a_j\tau(t_j) + b_j, \quad a_j, b_j \in \mathbb{Z}. \quad (4)$$

Thus the curves $G_{n_j^p}(C)$ meet over t_j for all $p > 0$. Their lifted tangent directions cannot eventually all coincide. Indeed, the deck-transformation formula reduces such coincidence to

$$\partial_t f(t_j) = \frac{a_j}{n_j - 1} \partial_t \tau(t_j). \quad (5)$$

In the isotrivial case $\partial_t \tau = 0$; in the non-isotrivial case the holomorphic ratios in (5) take values in $\mathbb{R}$. Constancy then forces $f = a\tau + b$, exactly the forbidden quasi-torsion form. Hence over every base open one obtains infinitely many distinct tangent directions in a projective-line fiber, forcing the Zariski density of the lifted curves.

Scope

The argument concerns elliptic complex projective K3 surfaces and allows the elliptic pencil to change during the existence step. It does not prove the stronger open statement that every polarization contains infinitely many rational curves in its multiples. Its output is instead a sequence of singular rational curves with unbounded classes, together with density of their first-jet lifts. As a secondary consequence, Corollary 7.3 combines Theorem 4.1 with Chen–Lewis to show that rational curves are dense in the usual topology on every projective elliptic K3 surface.