

Base change for semiorthogonal decompositions

Alexander Kuznetsov

arXiv:0711.1734

<http://arxiv.org/abs/0711.1734v2>

How a categorical piece changes base

Let $f: X \rightarrow S$ be a morphism of quasiprojective varieties and let $\phi: T \rightarrow S$ be a base change, with $X_T = X \times_S T$. Write f and ϕ also for the two projections from X_T . A subcategory of $D^b(X)$ is S -linear when it is stable under tensoring with $f^*D^{\text{perf}}(S)$. The base change is *faithful for f* when its Cartesian square is exact, equivalently when the canonical base-change map

$$\phi^* f_* \xrightarrow{\sim} f_* \phi^* \quad (1)$$

is an isomorphism. Flat f or flat ϕ is enough for faithfulness.

For a semiorthogonal decomposition $D^b(X) = \langle \mathcal{A}_1, \dots, \mathcal{A}_m \rangle$, *strong* means that each $\mathcal{A}_k$ is admissible in the tail category $\langle \mathcal{A}_k, \dots, \mathcal{A}_m \rangle$ (Definition 2.6). The problem is to construct meaningful components on a possibly singular X_T . Merely generating inside $D^b(X_T)$ from pulled-back bounded objects can recover only the perfect complexes, so bounded categories cannot be handled directly.

Main theorem (Kuznetsov, Theorem 5.6). *Assume that $D^b(X) = \langle \mathcal{A}_1, \dots, \mathcal{A}_m \rangle$ is an S -linear strong semiorthogonal decomposition, every projection functor has finite cohomological amplitude, and ϕ is faithful for f . Then the subcategories $\mathcal{A}_{iT} \subset D^b(X_T)$ constructed below form a T -linear semiorthogonal decomposition*

$$D^b(X_T) = \langle \mathcal{A}_{1T}, \dots, \mathcal{A}_{mT} \rangle. \quad (2)$$

Its projection functors have the same cohomological amplitudes as the original projections. Moreover, ϕ_ and ϕ^* are compatible with the induced decompositions in the precise unbounded and bounded-above categories stated in the paper.*

Ingredient I: begin with compact objects

Set $\mathcal{A}_i^{\text{perf}} = \mathcal{A}_i \cap D^{\text{perf}}(X)$ and define

$$\mathcal{A}_{iT}^p = \text{thick} \left\langle \phi^* F \otimes f^* G : F \in \mathcal{A}_i^{\text{perf}}, G \in D^{\text{perf}}(T) \right\rangle. \quad (3)$$

Faithfulness (1), perfectness, and S -linearity identify the relative RHom between generators with the pullback of the original one, proving semiorthogonality. Generation is subtler: Lemma 5.2 resolves a perfect object on X_T by external tensor products and shows that a far enough stupid truncation contains it as a direct summand. Decomposing the X -factor then proves $D^{\text{perf}}(X_T) = \langle \mathcal{A}_{1T}^p, \dots, \mathcal{A}_{mT}^p \rangle$ (Proposition 5.1).

Ingredient II: enlarge before intersecting back

Let $\widehat{\mathcal{A}}_{iT}$ be the smallest triangulated subcategory of $D_{\text{qc}}(X_T)$ containing $\mathcal{A}_{iT}^p$ and closed under arbitrary direct sums. Compactness of perfect complexes preserves semiorthogonality, while their generation of D_{qc} supplies fullness. Intersecting in stages gives

$$\mathcal{A}_{iT}^- = \widehat{\mathcal{A}}_{iT} \cap D^-(X_T), \quad \mathcal{A}_{iT} = \mathcal{A}_{iT}^- \cap D^b(X_T). \quad (4)$$

The unbounded detour is essential, not cosmetic: arbitrary sums and homotopy colimits allow perfect approximations to converge to coherent complexes that need not themselves be perfect.

Ingredient III: boundedness is detected after ample twists

Formula (4) is useful only if the projection of a bounded complex remains bounded. Lemma 5.4 supplies the detector. For a quasiprojective $\psi: Y \rightarrow X$, a line bundle L ample over X , and $F \in D_{\text{qc}}(Y)$, membership in $D^{[p,q]}$ is equivalent to the same uniform cohomological range for homotopy colimits of $\psi_*(F \otimes L^{k_j})$ along every sequence of maps $L^{k_1} \rightarrow L^{k_2} \rightarrow \dots$ with $k_j \rightarrow \infty$.

Apply this to $H \in D^{[p,q]}(X_T)$ and its i th bounded-above projection $\alpha_{iT}^-(H)$. Taking $L = f^*M$ for M ample over S , compatibility of pushforward and projection gives

$$\phi_*(\alpha_{iT}^-(H) \otimes f^*M^k) \simeq \hat{\alpha}_i(\phi_*(H \otimes f^*M^k)). \quad (5)$$

For $k \gg 0$, Lemma 2.20 writes the argument on the right as $\text{hocolim } G_j$ with every $G_j \in D^{[p,q]}(X)$. Since $\hat{\alpha}_i$ commutes with homotopy colimits and the original α_i has amplitude (a_i, b_i) , (5) lies in $D^{[p+a_i, q+b_i]}(X)$. Lemma 5.4 pulls this bound back to X_T . This is the hardest step: it simultaneously proves boundedness in (4) and preserves the exact cohomological amplitude, completing Theorem 5.6.

The flat-base-change model

When $T \rightarrow S$ is flat, faithfulness is automatic. For an exceptional collection, (3) says to pull each exceptional object to X_T and allow tensoring by perfect complexes from T ; the relative RHom calculation immediately transports the vanishing order. This is the clean geometric picture behind (2). The analogy stops when X_T is singular: pulled-back perfect objects generally generate only $D^{\text{perf}}(X_T)$, and the closure/intersection procedure (4) is what reaches all of D^b .

Consequences and scope

Corollary 5.7 shows that the base-changed category attached to one admissible component depends only on that component, not on a surrounding decomposition. Section 6 proves independence from a chosen geometric realization under its kernel-functor hypotheses. Finally, Theorem 7.1 applies the product decomposition to the diagonal object and represents each finite-amplitude projection functor by a kernel on $X \times X$.

The theorem assumes quasiprojectivity, strongness, S -linearity, finite cohomological amplitude, and faithful base change; it is not an arbitrary base-change assertion. Smooth projective X makes strongness automatic, and smooth X supplies the relevant amplitude finiteness noted in the introduction, but the stated theorem deliberately allows singular varieties when its explicit hypotheses are verified.