

24 rational curves on K3 surfaces

Slawomir Rams, Matthias Schütt

arXiv:1907.04182

<http://arxiv.org/abs/1907.04182v3>

Counting low-degree curves on a polarized K3 surface

Let (X, H) be a smooth K3 surface over an algebraically closed field, with H very ample and $H^2 = 2h$. Put $\deg C = H \cdot C$, let r_d count irreducible rational curves of degree exactly d , and set

$$S_d = r_1 + \cdots + r_d = \#\{C \subset X : C \text{ rational and } H \cdot C \leq d\}.$$

The point is that, once the polarization is sufficiently large relative to d , this unweighted count has a sharp universal bound.

Main theorem (Theorem 1.1). *Fix $d \geq 1$.*

- (i) *If $\text{char } k \neq 2, 3$ and $h > 42d^2$, then $S_d \leq 24$.*
- (ii) *For $h \gg 0$, if $S_d > 21$, then all rational curves of degree at most d are components of fibres of one genus-one fibration on X .*
- (iii) *If $d \geq 3$ and $h \geq d(2d+1) - 1$, then, over every characteristic other than 2, there are degree- $2h$ polarized K3 surfaces with $r_d = 24$.*

Part (ii) explains the number 24: for an elliptic K3 fibration the total Euler contribution of the singular fibres is $e(X) = 24$. Part (iii) shows that the bound is not only asymptotic but attained throughout a large range of polarizations.

The intersection-graph trichotomy

Let Γ be the graph of all curves counted by S_d , with edge multiplicities $C \cdot C'$, and let $M = \mathbf{Z}\Gamma$ carry the intersection form. The Hodge index theorem leaves three possibilities.

- If M is negative definite, its (-2) -curves form ADE root diagrams; the Picard rank bounds $\#\Gamma \leq 21$.
- If M is negative semidefinite with kernel, an isotropic vertex or affine Dynkin cycle produces a genus-one fibration. Every curve in Γ is a rational fibre component. Outside characteristics 2, 3, the Euler formula bounds their number by 24.
- The remaining case is hyperbolic. This is the only place where the numerical threshold $42d^2$ is needed.

Intrinsic polarization and exclusion of the hyperbolic case

For a hyperbolic subgraph Γ_0 , solve inside its rational span for an intrinsic class H_0 satisfying $H_0 \cdot C = H \cdot C$ on the chosen curve basis. Orthogonally projecting the actual polarization into that span gives the decisive inequality

$$2h = H^2 \leq H_0^2.$$

If G is the Gram matrix and $\mathbf{d} = (H \cdot C)$, then $H_0^2 = \mathbf{d}^T G^{-1} \mathbf{d}$; since every coordinate is at most d , the positive entries of G^{-1} give explicit upper bounds for H_0^2 .

Under $h > 42d^2$, the same determinant estimates force every curve in a hypothetical hyperbolic Γ to be a smooth (-2) -curve and rule out small Kodaira cycles. Any non-ADE graph nevertheless contains an affine Dynkin diagram, hence supports a Kodaira fibre D ; necessarily $\deg D > 6d$. In the non-quasi-elliptic case relevant to part (i), if $\#\Gamma > 24$, rank and Euler-number arguments force at least three multisections of $|D|$. Successively replacing D by smaller supported fibre configurations reduces the possibilities to I_1^* , I_2^* , or IV^* . Classified extremal alternatives have at most 24 components or belong to the characteristic-2, 3 exceptions. The remaining attachments of three sections create a forbidden small cycle, violate a multisection-degree bound, or contradict an intrinsic-polarization estimate (for example, for the relevant I_2^* configuration). Thus a hyperbolic graph with more than 24 vertices cannot occur, completing the trichotomy.

Why the bound is sharp

The construction begins with K3 surfaces whose generic Picard lattice has Gram matrix

$$\begin{pmatrix} 0 & d \\ d & c \end{pmatrix}, \quad c \in 2\mathbf{Z}.$$

After reflections, the isotropic generator becomes the fibre class E of a genus-one fibration. A general member has exactly 24 nodal singular fibres; each fibre is an irreducible rational curve. If B is the reflected second generator, then $E \cdot B = d$ and $NE + B$ is very ample for $N > 2d$, so every nodal fibre has degree d . Varying c and N realizes every $h \geq d(2d + 1) - 1$ and gives $r_d = 24$.

This model also exposes the proof's central theme: a large collection of low-degree rational curves is forced to behave like fibre components, and the K3 Euler number then caps the collection. The technically hardest step is making that heuristic effective in the hyperbolic lattice, where the constant 42 comes from explicit Gram-matrix bounds and a finite Kodaira-configuration analysis.

Exceptional characteristics and omissions

Quasi-elliptic fibrations in characteristics 2 and 3 break the clean Euler count. The paper proves separate bounds: after excluding cuspidal arithmetic-genus-one curves, at most 40 curves occur when $p = 2$ and $h > 46.25d^2$, and at most 30 when $p = 3$ and $h > 43d^2$; additional hypotheses recover the bound 24. This note omits the detailed classification of those exceptional configurations and the explicit Fermat, Hesse, and $12I_2$ models used to realize smooth equality cases.