

The Dynamical Manin-Mumford Conjecture and the Dynamical Bogomolov Conjecture for split rational maps

Dragos Ghioca, Khoa D. Nguyen, Hexi Ye

arXiv:1511.06081

<http://arxiv.org/abs/1511.06081v2>

Small points force an invariant curve

Let $f_i \in \mathbf{C}(z)$ have degrees $d_i > 1$, and let $C \subset \mathbf{P}^1 \times \mathbf{P}^1$ be irreducible and dominant over both factors. A degree- d rational map is called *exceptional* here if it is linearly conjugate to $z^{\pm d}$, to $\pm T_d$, or to a Lattès map.

Main theorem (Ghioca–Nguyen–Ye, Theorem 1.1). *Assume that at least one of f_1, f_2 is nonexceptional. Then C contains infinitely many points (x_1, x_2) with each x_i preperiodic for f_i if and only if there are $\ell_1, \ell_2 \geq 1$ such that*

$$d_1^{\ell_1} = d_2^{\ell_2}, \quad C \text{ is preperiodic under } (f_1^{\ell_1}, f_2^{\ell_2}). \quad (1)$$

If f_1, f_2, C are defined over $\overline{\mathbf{Q}}$, condition (1) is also equivalent to the existence of infinitely many $P_n = (x_{1,n}, x_{2,n}) \in C(\overline{\mathbf{Q}})$ satisfying

$$\widehat{h}_{f_1}(x_{1,n}) + \widehat{h}_{f_2}(x_{2,n}) \longrightarrow 0. \quad (2)$$

Thus the conclusion synchronizes unequal-degree systems by suitable iterates. The exceptional exclusion is essential without an equal-degree hypothesis: algebraic-group covers create extra invariant relations.

Ingredient I: one small sequence gives two equal measures

Work first over $\overline{\mathbf{Q}}$ and replace C by its normalization. For the projections $\pi_i : C \rightarrow \mathbf{P}^1$, pull back the canonical semipositive adelic metric attached to f_i on $\mathcal{O}(1)$. The induced height is exactly

$$\widehat{h}_{\mathcal{L}_i}(P) = \widehat{h}_{f_i}(\pi_i(P)). \quad (3)$$

Dominance supplies infinitely many points above preperiodic values, so Zhang’s height inequality gives $\widehat{h}_{\mathcal{L}_i}(C) = 0$. Thus the points in (2), together with nonnegativity, are small for both $\mathcal{L}_i$. Yuan–Chambert-Loir equidistribution gives the same weak limit for their Galois orbits. At the complex place this says

$$\widetilde{\mu}_1 := \frac{\pi_1^* \mu_{f_1}}{\deg \pi_1} = \frac{\pi_2^* \mu_{f_2}}{\deg \pi_2} =: \widetilde{\mu}_2, \quad (4)$$

where μ_{f_i} is the maximal-entropy measure. This is Proposition 4.1: arithmetic smallness has become a local analytic rigidity statement on C .

Ingredient II: convert (4) into a local conjugacy

Choose an unramified point (x_0, y_0) above a repelling periodic point of f_1 and write the nearby branch of C as $y = h(x)$. Coordinatewise iterates and an image of C make x_0, y_0 fixed while preserving (4). Put locally $\tilde{g} = h^{-1} \circ f_2 \circ h$, using a chosen branch of h^{-1} ; if h has local degree j , the multiplier of $\tilde{g}$ is a j th root of the multiplier of f_2 . Measure scaling first forces y_0 to be repelling: otherwise a sufficiently large power of $\tilde{g}$ would map a small disc inside its f_1 -image but multiply its measure by a larger degree, contradicting that $x_0 \in \text{supp } \mu_{f_1}$.

Linearize both germs. There are pairs $n_j, m_j \rightarrow \infty$ for which their multiplier ratio tends to 1, so

$$f_1^{n_j} \circ \tilde{g}^{-m_j} \quad (5)$$

forms a nontrivial normal family of local symmetries of the Julia set. Levin's theorem says that an infinite such family occurs exactly in the postcritically finite parabolic-orbifold case, hence in the exceptional case. Nonexceptionality therefore makes two germs in (5) coincide. Cancellation gives, for some $n, m \geq 1$,

$$f_1^n = h^{-1} \circ f_2^m \circ h. \quad (6)$$

This finite-collision step is the hardest point of the proof.

Ingredient III: globalize and match degrees

Equation (6) says that C and $(f_1^n, f_2^m)(C)$ share an analytic arc; irreducibility makes them the same algebraic curve. Applying the common measure to a small injectivity disc in (6) then yields $d_1^n = d_2^m$. After one pass obtains equal-degree iterates, a second pass from the original curve gives its preperiodicity, not merely periodicity of a later image. Conversely, if equal-degree iterates make C preperiodic, the fibres of $C \rightarrow \mathbf{P}^1$ show that every preperiodic first coordinate has a preperiodic second coordinate; hence C contains infinitely many such points (Lemma 6.1).

For data over an arbitrary characteristic-zero subfield of $\mathbf{C}$, the authors spread the maps, curve, and preperiodic points over a finite-type $\mathbf{Q}$ -variety. Yuan–Zhang specialization keeps the specialized set infinite on a dense open; (4) holds there and continuity of the potentials returns it to the original complex fibre. This is the bridge from the number-field equidistribution argument to the complex statement.

Toy model and scope

For $f_1 = f_2 = f$ and the diagonal $C = \{x = y\}$, one has $\tilde{\mu}_1 = \tilde{\mu}_2$, equation (6) is tautological, and every (x, x) with x preperiodic lies on C . The model displays the conclusion but removes the proof's main difficulty: extracting a global algebraic relation from equality of two measures on an arbitrary correspondence.

The theorem concerns curves dominant over both factors; horizontal and vertical cases are separate and elementary. When both maps are exceptional, the paper instead proves the equal-degree tangent-space formulation of dynamical Manin–Mumford (Theorem 1.3). Its later classifications of unicritical-polynomial relations and invariant curves are important further applications, but are not needed for the mechanism explained here.