Equidistribution of periodic points of some automorphisms on K3 surface

Chong Gyu Lee

arXiv:1102.4860v1 (2011), 10 pages
http://arxiv.org/abs/1102.4860v1

Several-map polarization

Let W be a projective variety over a number field and let $\mathcal{M} = \{\phi_1, \dots, \phi_t\}$ be endomorphisms. The system is *polarizable* by an ample line bundle L if

$$\bigotimes_{i=1}^{t} \phi_i^* L \simeq L^{\otimes q}, \qquad q > t.$$

Since words of length N number t^N while polarization grows as q^N, the inequality $q > t$ lets Kawaguchi's limit put one semipositive adelic metric on L.

Main theorem (Theorem 1.2). *For a polarizable system $(W, \mathcal{M}, L)$ and a generic sequence of closed points x_m that is small for its canonical height, the probability measures on the Galois orbits Γ_{x_m} converge weakly at every place v to*

$$\mu_{\mathcal{M},v} = \frac{c_1(\overline{L})_v^{\dim W}}{\deg_L W}.$$

For a surface automorphism with $\sigma^* L \otimes (\sigma^{-1})^* L \simeq L^{\otimes q}$, $q > 2$, Theorem 1.3 and Corollary 4.3 also claim density and equidistribution of periodic points. The density proof has the gap described below, so these remain claimed applications rather than established consequences.

For the $(1,1)$–$(2,2)$ K3 surface $S \subset \mathbb{P}^2 \times \mathbb{P}^2$, the projections π_i give involutions ι_i. With $L_i = \pi_i^* \mathcal{O}(1)$, $\sigma = \iota_2 \iota_1$, and $L = L_1 \otimes L_2$, the paper computes

$$\sigma^* L \otimes (\sigma^{-1})^* L \simeq L^{\otimes 14},$$

for a concrete 18-parameter family.

Proof mechanism

1. Canonical metric and height. Kawaguchi supplies an admissible semipositive metric whose height satisfies

$$\sum_{i=1}^{t} \widehat{h}_L(\phi_i(x)) = q \widehat{h}_L(x), \qquad \widehat{h}_L(x) \geq 0.$$

Periodic points have height zero. At $q = t$, however, word averaging need not decrease the height of a common fixed point.

2. Equidistribution of any generic small sequence. Yuan's theorem applies to $\overline{L}$: smallness controls height, genericity prevents concentration on a proper subvariety, and Galois-orbit measures converge to $c_1(\overline{L})_v^{\dim W} / \deg_L W$.

3. Polarization survives iteration. Lemma 4.1 inductively produces $q_m > 2$ with

$$(\sigma^m)^* L \otimes (\sigma^{-m})^* L \simeq L^{\otimes q_m}.$$

via $q_{m+1} = q q_m - q_{m-1}$. Thus one may pass to an iterate fixing a hypothetical curve component.

4. The intended density step, and its gap. The intended toy contradiction fixes a curve component C of the periodic-point closure by some σ^m and intersects with C: $2(L \cdot C) = q_m (L \cdot C)$, impossible for $q_m > 2$ and ample L. But the paper neither excludes a finite zero-dimensional closure nor first proves infinitely many periodic points. Density is therefore unproved by this argument. The subsequent use of Zhang's successive minima to obtain a generic small periodic sequence, and hence periodic equidistribution, remains conditional on a valid density input.

Scope

The fully supported result is Theorem 1.2, resting on the semipositive metric and Yuan's theorem. Theorem 1.3 is printed for a projective variety although its proof and Lemma 4.1 use surfaces, where the zero-dimensional case still remains. The examples concern only K3 automorphisms polarizable together with their inverses, not arbitrary ones.

Codex mathematical paper digest, 2026-10-27. Source paper should be consulted for full statements and proofs.