Polynomials with many rational preperiodic points

John R. Doyle, Trevor Hyde

arXiv:2201.11707

http://arxiv.org/abs/2201.11707v2

Dynamical compression and two extremal questions

For a polynomial $f \in K[x]$, let $\mathrm{PrePer}(f, K)$ be the points with finite forward orbit. The paper says that a degree-r polynomial g exhibits *dynamical compression* if it is linearly conjugate to a polynomial f satisfying

$$f([m]) \subseteq [n] \qquad \text{for some } m \geq n > r+1,$$

where $[m] = \{1, \ldots, m\}$. For the interval-compression mechanism itself, if

$$f([m]) \subseteq [n] \qquad (m > n),$$

then every point of $[m]$ is rational preperiodic. Doyle and Hyde call this condition into service to produce many rational preperiodic points and, less visibly, many common complex preperiodic points for the translates $f + i$.

Main theorem (Doyle–Hyde, Theorem 1.3)**.** *For every integer $d \geq 2$ there is $f_d \in \mathbf{Q}[x]$ with $2 \leq \deg(f_d) \leq d$ such that*

(i) *f_d has at least $d + \lfloor \log_2 d \rfloor$ rational preperiodic points;*
(ii) *for $0 \leq i < j \leq \lfloor \log_2 d \rfloor$, the sets $\mathrm{PrePer}(f_d + i, \mathbf{C})$ and $\mathrm{PrePer}(f_d + j, \mathbf{C})$ are distinct and have finite intersection;*
(iii)

$$\left| \bigcap_{i=0}^{\lfloor \log_2 d \rfloor} \mathrm{PrePer}(f_d + i, \mathbf{C}) \right| \geq \deg(f_d)(d-1) + 1.$$

Consequently, for infinitely many degrees d, two degree-d polynomials have at least $d^2 + d\lfloor \log_2 d \rfloor - 2d + 1$ common complex preperiodic points.

The first bound improves the $d+1$ points supplied by unconstrained interpolation by a logarithmic term. The theorem gives lower bounds; the corresponding uniform upper bounds remain conjectural.

Ingredient 1: an evaluation lattice with exact discriminant

For $e \geq 0$, let $\Lambda_{d,e} \subset \mathbf{R}^{d+e+1}$ be generated by

$$u_i = \left(\tbinom{0}{i}, \tbinom{1}{i}, \ldots, \tbinom{d+e}{i} \right), \qquad 0 \leq i \leq d.$$

Thus its lattice points are evaluation vectors of degree-at-most-d integer-valued polynomials on $0, \ldots, d+e$. Theorem 2.1 computes

$$\delta(\Lambda_{d,e}) = \prod_{i=0}^{d} \prod_{j=1}^{e} \frac{d+i+j+1}{i+j}.$$

The proof identifies the Gram determinant with nonintersecting rhombic paths by Lindström–Gessel–Viennot, hence with plane partitions in a $(d+1) \times (d+1) \times e$ box; MacMahon's formula gives the product. This exact expression, rather than its leading-order size alone, provides the lower-order saving needed when e is logarithmic in d.

Ingredient 2: Minkowski finds a nonlinear short vector

Adjoin e mutually orthogonal vectors of a very large common length in the orthogonal complement of $\Lambda_{d,e}$. For $e = \lfloor \log_2(d/2) \rfloor$, Corollary 2.2 and Minkowski's product bound for successive minima imply that the third minimum of the enlarged lattice is less than $d/2$ when $d \geq 33$. Three independent short vectors cannot use the long directions; at least one therefore comes from an integer-valued polynomial g of degree at least two, with

$$|g(r)| < d/2 \qquad (0 \leq r \leq d + e).$$

After translating the input and output, one obtains Theorem 2.5:

$$f_d\big([d + \lfloor \log_2 d \rfloor]\big) \subseteq [d].$$

The cases $2 \leq d \leq 32$ are supplied by interpolation and the paper's explicit auxiliary family. This is the existence step behind the rational lower bound.

Ingredient 3: compression survives every translate

Suppose f has degree $r \geq 2$ and $f([m]) \subseteq [n]$. For $0 \leq i \leq m - n$,

$$(f + i)\big(f^{-1}([n])\big) \subseteq [n + i] \subseteq [m] \subseteq f^{-1}([n]).$$

Hence $f^{-1}([n])$ is a finite forward-invariant set for every $f + i$. Counting ramification via f' gives

$$|f^{-1}([n])| \geq rn - r + 1.$$

Proposition 3.2 also proves that the preperiodic sets for distinct positive integer translates are unequal; the Baker–DeMarco theorem then makes each pairwise intersection finite. Applying this with $m = d + \lfloor \log_2 d \rfloor$ and $n = d$ proves the simultaneous-intersection part of the main theorem. A degree lower bound for interval-compressing polynomials due to Cohen–Shpilka–Tal supplies infinitely many actual degrees for the final quadratic-plus-logarithmic estimate.

A quadratic model and the hard point

The polynomial

$$f(x) = \frac{x^2 - 9x + 22}{2}$$

satisfies $f([8]) \subseteq [7]$. Thus its eight integer inputs are rational preperiodic, while $f^{-1}([7])$ gives at least $2 \cdot 7 - 2 + 1 = 13$ common complex preperiodic points for f and $f + 1$. This small example contains the whole bridge from an evaluation vector to both dynamical conclusions.

The difficult quantitative point is forcing a genuinely nonlinear short lattice vector while taking $e \sim \log_2 d$: the exact discriminant estimate is what makes the third successive minimum small enough. The logarithmic construction is nonexplicit. The paper separately gives explicit degree-d polynomials with $d + 6$ rational preperiodic points and records stronger low-degree computations; those refinements, and the detailed closed formulas for the explicit family, are omitted here.