Stability of some vector bundles on Hilbert schemes of points on K3 surfaces

Fabian Reede, Ziyu Zhang

arXiv:2103.10328; doi:10.1007/s00209-021-02920-6

https://arxiv.org/abs/2103.10328

Turning a universal family around

Let X be a projective K3 surface over $\mathbf{C}$ and $X^{[k]}$ its Hilbert scheme of length-k subschemes. A universal bundle on $X \times X^{[k]}$ is usually read as a family on X, parametrized by $X^{[k]}$. Reede and Zhang reverse the roles: for $x \in X$, restrict to the "wrong-way" fibre E_x on $X^{[k]}$, and ask whether all these bundles are stable for one polarization and form a complete moduli component.

The paper treats exactly two lattice configurations:

$$\begin{aligned}
&\text{(I)} \quad \mathrm{NS}(X) = \mathbf{Z}h, \quad h^2 = 4k, \quad k \geq 1, \qquad v = (k+1, -h, 1);\\
&\text{(II)} \quad \mathrm{NS}(X) = \mathbf{Z}e \oplus \mathbf{Z}f, \quad \begin{pmatrix} e^2 & ef \\ ef & f^2 \end{pmatrix} = \begin{pmatrix} -2k & 2k+1 \\ 2k+1 & 0 \end{pmatrix}, \quad k \geq 2,\\
&\qquad h = e + (2k-1)f, \qquad v = (2k-1, h, 2k).
\end{aligned}$$

Main theorem (Theorem 0.1)**.** *In either case there is a fine moduli space M of stable vector bundles of Mukai vector v on X, an isomorphism $M \simeq X^{[k]}$, and a universal bundle $\mathcal{E}$. Some ample divisor H on M makes every wrong-way fibre E_x simultaneously μ_H-stable. The resulting classifying map*

$$X \longrightarrow \mathcal{M}_H(M), \qquad x \longmapsto [E_x],$$

identifies X with a smooth connected component of the moduli space of μ_H-stable sheaves on M.

Thus a K3 surface is recovered inside a moduli space on its higher-dimensional hyperkähler Hilbert scheme. When $k \geq 2$, the bundles are not tautological: their ranks $k+1$ and $2k-1$ are not divisible by k (Remark 3.17).

Ingredient 1: explicit Fourier–Mukai families

In case (I), $\mathcal{O}_X(1)$ is k-very ample. For $Z \in X^{[k]}$ the evaluation sequence gives

$$0 \longrightarrow E_Z \longrightarrow H^0(I_Z(1)) \otimes \mathcal{O}_X \longrightarrow I_Z(1) \longrightarrow 0.$$

The bundle E_Z has Mukai vector $(k+1, -h, 1)$ and is the shifted spherical twist $T_{\mathcal{O}_X}(I_Z(1))[1]$. Since the twist is an autoequivalence, the classifying map $X^{[k]} \to M_h(v)$ separates points; equal dimensions and properness give the isomorphism (Theorem 2.3).

In case (II), the autoequivalence $\Theta = (- \otimes \mathcal{O}_X(f)) \circ T_{\mathcal{O}_X}^{-1} \circ (- \otimes \mathcal{O}_X(e))$ sends I_Z to a bundle fitting into

$$0 \longrightarrow \mathcal{O}_X(f)^{\oplus(2k-2)} \longrightarrow E_Z \longrightarrow I_Z(e+f) \longrightarrow 0.$$

The lattice inequalities and the elliptic fibration defined by f control every torsion-free quotient, proving μ_h-stability (Theorem 3.7). Again the equivalence separates the E_Z, yielding $X^{[k]} \simeq M_h(v)$.

Ingredient 2: stability in the wrong direction

For (I), each wrong-way fibre has an exact sequence

$$0 \longrightarrow E_x \longrightarrow K \longrightarrow I_{S_x} \longrightarrow 0, \qquad c_1(E_x) = c_1(K) = -h_k + \delta,$$

where S_x is the incidence locus, h_k is the natural nef class, and 2δ is the Hilbert–Chow exceptional divisor. Pulling to the ordered configuration space X^k turns K into the kernel of an evaluation map. It has no global sections, so an $\mathfrak{S}_k$-invariant destabilizing subsheaf cannot have slope zero; the remaining integral first-Chern classes leave no integer between the slopes of E_x and K (Propositions 2.6–2.7).

For (II), use the ample class $\widehat{h} = 2ke + (2k-1)f$ on X, for which $e\widehat{h} = -1$. After passage to X^k, the common subbundle in $0 \to \mathcal{H} \to E_x \to I_{S_x} \to 0$ is an equivariant extension of $\bigoplus_i q_i^* \mathcal{O}_X(e)$, where $q_i \colon X^k \to X$ is the ith projection, by a trivial bundle and has no invariant global section. A putative destabilizer has either negative degree, already too small, or nonnegative degree; the latter would force precisely such an invariant trivial summand. This proves stability for the nef class $\widehat{h}_k$ (Proposition 3.14).

Nef stability is not enough for the moduli problem. The subtle uniform step is to perturb to one ample class that works for every x. In case (I), all E_x embed in the fixed bundle K, so the first Chern classes of possible near-destabilizers form a finite set. Case (II) needs an extra bridge. From the exact sequence for E_x, a candidate F is an extension of $F_2 \subset I_{S_x}$ by $F_1 \subset \mathcal{H}$. Replace it by $F' = F_1 \oplus F_2$, which has the same first Chern class and lies in $E'_x = \mathcal{H} \oplus I_{S_x} \subset \mathcal{H} \oplus \mathcal{O}_{X^{[k]}}$; the last ambient sheaf is fixed, so finiteness follows here as well. Intersecting the semistability chambers then gives a full-dimensional convex region whose closure contains the nef class. An ample interior point makes every E_x stable simultaneously (Theorems 2.8 and 3.15).

Ingredient 3: the component calculation

Let $\Phi \colon D^b(X) \to D^b(X^{[k]})$ be the universal-ideal functor. Addington's result makes it a $\mathbf{P}^{k-1}$-functor, so

$$\operatorname{Ext}^*(\Phi A, \Phi B) \simeq \operatorname{Ext}^*(A, B) \otimes H^*(\mathbf{P}^{k-1}, \mathbf{C}).$$

In case (I), $E_x = \Phi(I_x(1))$; in case (II) an autoequivalent transform of $\mathcal{O}_x$ plays the same role. Consequently $E_x \not\simeq E_y$ for $x \neq y$ and

$$\operatorname{Ext}^1(E_x, E_x) \simeq T_x X.$$

The classifying map is injective on closed points and its target has tangent dimension two along the image, hence the image is a smooth connected component (Theorems 2.10 and 3.16).

The result is intentionally not a general theorem for all K3 lattices or all universal bundles. Its hardest point is uniform stability: individual nef stability does not by itself produce a single ample polarization for the whole family. The two explicit constructions supply the fixed ambient sheaves and lattice gaps that make that passage possible.