Gap for geometric canonical height functions

Yugang Zhang

arXiv:2307.08561

http://arxiv.org/abs/2307.08561v2

A discrete gap for a geometric height

Let B be a smooth complex projective variety of dimension $b \geq 1$, $K = \mathbf{C}(B)$, and $f\colon \mathbf{P}^k_K \to \mathbf{P}^k_K$ an endomorphism of degree $d \geq 2$. After restricting to a nonempty open $\Lambda \subset B$, it is a family f_t of endomorphisms of $\mathbf{P}^k_{\mathbf{C}}$. Its canonical height $\widehat{h}_f\colon \mathbf{P}^k(\overline{K}) \to \mathbf{R}_{\geq 0}$ satisfies $\widehat{h}_f(f(x)) = d\widehat{h}_f(x)$. A geometric point of height zero is called *stable*; preperiodic points are stable, but the converse need not hold.

Main theorem (Zhang, Theorem 1.1)**.** *There is an $\varepsilon > 0$, depending on f and K, such that for every $x \in \mathbf{P}^k(K)$,*

$$\widehat{h}_f(x) < \varepsilon \quad \Longrightarrow \quad \widehat{h}_f(x) = 0.$$

If the stable K-points are Zariski dense, then f is birationally isotrivial: the maps f_t on general fibres are pairwise birationally conjugate.

The paper also notes that both statements extend to polarized endomorphisms of smooth projective varieties; the gap statement needs only normality.

Ingredient 1: turn height into bounded orbit-cycles

Choose an ample M on B and an ample model line bundle L on $\mathcal{X} = \mathbf{P}^k_{\mathbf{C}} \times B$. Normalized graph models for the iterates of f give projections $\pi_n\colon \mathcal{X}_n \to B$ and nef line bundles L_n. If $\sigma_x \subset \mathcal{X}$ and $\sigma_{x,n} \subset \mathcal{X}_n$ are the Zariski closures of x in the corresponding models, the height is the intersection limit

$$\widehat{h}_f(x) = \frac{1}{[k(x) : K]} \lim_n [\sigma_{x,n}] \cdot c_1(L_n) \cdot c_1(\pi_n^* M)^{b-1}.$$

The vertical error between consecutive models is bounded by a geometric series. Consequently $\widehat{h}_f$ differs from the naive model height by a constant independent of x (Proposition 2.1).

Proposition 2.3 then identifies height zero with uniform boundedness of all orbit closures $f^n(\sigma_x)$. A mixed-intersection inequality upgrades the bound involving $c_1(L)c_1(\pi^* M)^{b-1}$ to a bound on the full degree $[f^n(\sigma_x)] \cdot c_1(L)^b$. This is what makes the relevant orbit closures live in a fixed projective piece of the cycle space.

Ingredient 2: Noetherianity makes the gap finite

For every $r \geq 0$, use the normalized graph of (id, f^r) to define a closed locus S_r of bounded b-cycles whose cycle and r-th image both retain the required degree bound. The intersection $S = \bigcap_r S_r$ parameterizes cycles with stable horizontal part (after discarding the allowed vertical contribution over $B \setminus \Lambda$). Because the ambient cycle space is Noetherian, this infinite intersection already equals $S_0 \cap \cdots \cap S_N$ for some finite N.

If c_u is the uniform model-height error, then

$$[f^n(\sigma_x)] \cdot c_1(L)c_1(\pi^* M)^{b-1} < c_u + d^n \widehat{h}_f(x).$$

Thus $\widehat{h}_f(x) < c_u/d^N$ puts the first N orbit-cycles in the bounded locus; finite stabilization puts every iterate there, and Proposition 2.3 forces $\widehat{h}_f(x) = 0$. The essential mechanism is a useful toy model: an apparently asymptotic orbit condition becomes a finite test once its parameter space is Noetherian.

Ingredient 3: dense stable sections trivialize the family

Let Z parameterize stable irreducible horizontal cycles birational to B. Under the density hypothesis, choose an irreducible component of maximal dimension and continue to denote it by Z. Let $\Gamma(Z)$ be its universal graph with projections $\pi_1\colon \Gamma(Z) \to \mathcal{X}$ and $\pi_2\colon \Gamma(Z) \to Z$. Assume stable K-points are Zariski dense. The decisive step is that π_1 is birational.

Indeed, take a repelling periodic point p in a regular fibre and replace f by an iterate so that p is fixed. The implicit-function theorem continues p to a local fixed section σ_0. Uniform expansion near σ_0 says

$$\operatorname{dist}(f_t(q), \sigma_0(t)) > \mu \operatorname{dist}(q, \sigma_0(t)), \qquad \mu > 1.$$

A stable section through p has orbit-cycles in a compact bounded family; their subsequential limit cannot acquire a vertical component over the regular locus. Expansion therefore forces that section to equal σ_0. Repelling periodic points are Zariski dense in each regular fibre, so a general point lies on a unique stable cycle and π_1 has generic degree one.

On a dense open $Y \subset \mathcal{X}_\Lambda$ where π_1 is an isomorphism, the regular maps $\Psi_t = \pi_2 \circ \pi_1^{-1}|_{Y_t}$ induce birational maps from $\mathcal{X}_t$ to Z. Hence f_t and $f_{t'}$ are conjugate by $\Psi_t^{-1} \circ \Psi_{t'}$, which is the asserted birational isotriviality.

The projective-plane Northcott consequence

If $f\colon \mathbf{P}^2_K \to \mathbf{P}^2_K$ is not isotrivial, Corollary 1.3 produces an $\varepsilon > 0$ and a proper preperiodically isotrivial subvariety E such that

$$\{x \in (\mathbf{P}^2_K \setminus E)(K) : \widehat{h}_f(x) < \varepsilon\} \quad \text{is finite.}$$

Here every curve component of E eventually becomes periodic with an isotrivial return map, and every zero-dimensional component eventually enters such a curve. In particular, only finitely many preperiodic K-points lie outside E. The proof uses the theorem above together with the projective-plane result that birational isotriviality of an endomorphism is equivalent to isotriviality, then analyzes the zero- and one-dimensional stable-cycle loci.

Hard point, meaning, and scope

The algebraic heart is not the functional equation for height, but the uniform passage from height zero to one bounded projective cycle space; without the mixed-intersection estimate, Noetherian stabilization cannot be applied uniformly. The analytic heart is the uniqueness of a bounded stable section through a repelling point. The result supplies a genuine gap, not merely a description of the zero set. It does not assert that height-zero points are always preperiodic, nor that birational isotriviality is ordinary isotriviality in every dimension.