

The Dynamical Mordell–Lang problem

Jason P. Bell, Dragos Ghioca, Thomas J. Tucker

arXiv:1401.6659

<http://arxiv.org/abs/1401.6659v1>

A topological almost Mordell–Lang theorem

Let X be a Noetherian topological space, so descending chains of closed subsets stabilize, and let $\Phi: X \rightarrow X$ be continuous. For a closed set Y and $x \in X$, write

$$S_Y(x) = \{n \in \mathbb{N} : \Phi^n(x) \in Y\}, \quad \delta(S) = \limsup_{|I| \rightarrow \infty} \frac{|S \cap I|}{|I|},$$

where I ranges over intervals in $\mathbb{N}$. The second quantity is upper Banach density.

Main theorem (Bell–Ghioca–Tucker, Theorem 1.4 and Corollary 1.5). *The set $S_Y(x)$ is a union of finitely many arithmetic progressions and a set of Banach density zero.*

Consequently, let X be a quasi-projective variety over any field K , $\Phi: X \dashrightarrow X$ a rational map, $Y \subseteq X$ a K -subvariety, and $x \in X(K)$. If every point of the forward orbit of x lies in the domain of definition of Φ , then

$$\{n \in \mathbb{N} : \Phi^n(x) \in Y(K)\}$$

has the same form.

Conjecture 1.1, stated for endomorphisms over $\mathbb{C}$, predicts no exceptional remainder. Extending that prediction to rational maps is a further formulation. The theorem is weaker, but it needs neither characteristic zero nor an unramified map: all arithmetic geometry has been compressed into the Noetherian Zariski topology.

Ingredient I: density forces a repeated gap

The combinatorial engine is Lemma 2.1. If $S \subseteq \mathbb{N}$ has $\delta(S) > 0$, divide long intervals on which S is dense into blocks of a fixed length $N > 1/\delta(S)$. A positive proportion of the blocks contain two points of S . Pigeonholing their differences produces an integer $1 \leq k < N$ and a positive-density set $Q \subseteq S$ such that

$$n \in Q \implies n + k \in S. \tag{1}$$

The positive density of Q , not merely its infinitude, is what lets the argument descend to a smaller closed set without losing its hypothesis.

Ingredient II: Noetherian descent creates a progression

Proposition 3.1 says that a positive-density hitting set already contains an infinite arithmetic progression. If not, choose by Noetherianity a minimal closed W for which

$$T_W = \{n : \Phi^n(x) \in W\}$$

has positive density but no progression. Apply (1) to T_W . For $n \in Q$, both $\Phi^n(x)$ and $\Phi^{n+k}(x)$ lie in W , hence

$$\Phi^n(x) \in Z := W \cap \Phi^{-k}(W). \tag{2}$$

If $\Phi^{-k}(W) \supseteq W$, then T_W is closed under adding k and already contains a progression. Otherwise $Z \subsetneq W$; its hitting set contains the positive-density set Q , so minimality forces a progression in $T_Z \subseteq T_W$. Either case is the desired contradiction.

Ingredient III: one progression controls every residue

To upgrade Proposition 3.1 to the theorem, suppose a counterexample exists and choose a minimal Noetherian closed space V supporting one. Fix its counterexample data: a continuous map $g: V \rightarrow V$, a closed $W \subset V$, and $y \in V$, with

$$T = \{n : g^n(y) \in W\}$$

not of the required form. This bad set has positive density, so Proposition 3.1 supplies $a, b \in \mathbb{N}$ with $\{an+b : n \geq 0\} \subseteq T$. Let C_i be the closure of the tail $\{g^{an+b}(y) : n \geq i\}$. The descending chain C_i stabilizes; call a stable member V_0 . Then $g^{-a}(V_0) \supseteq V_0$. For $0 \leq j < a$, put $V_j = g^{-j}(V_0)$. These are proper closed subsets of V , and $h = g^a$ restricts to a self-map of every V_j . With $y_j = g^{a+b-j}(y)$, minimality of V decomposes each explicitly defined set

$$T_j = \{n \geq m : h^n(y_j) \in W \cap V_j\}$$

into finitely many progressions plus a zero-density set. Recombining the a sets T_j , one for each residue class, reconstructs T up to finitely many terms, giving the forbidden decomposition.

Ingredient IV: rational maps become continuous

A rational map is only continuous on its open domain U . Theorem 3.2 passes to the invariant subspace

$$Z = \bigcap_{n \geq 0} \Phi^{-n}(U). \tag{3}$$

It is Noetherian with the inherited topology, contains the chosen orbit, and $\Phi|_Z: Z \rightarrow Z$ is continuous. Applying Theorem 1.4 to $Y \cap Z$ proves the algebraic corollary. This small reduction is why the result covers rational, not only regular, self-maps.

Toy model, scope, and bottleneck

On a finite space every orbit is eventually periodic, so every hitting set is already a finite union of progressions and a finite set. The proof replaces finiteness of points by Noetherian finiteness of closed-set descent; the stabilized tail closures play the role of the eventual cycle.

The zero-density remainder can still be infinite, and the paper gives positive-characteristic examples showing that very sparse nonperiodic return sets are natural. It therefore does not prove the full dynamical Mordell–Lang conjecture. Theorem 1.7 also proves the same finite-progressions-plus-zero-density conclusion for a coherent backward orbit $\{x_{-n}\}$ satisfying $f(x_{-n-1}) = x_{-n}$; its proof parallels the forward minimal-counterexample argument. The hardest step is the upgrade from the density lemma’s one progression to a finite global decomposition using stabilized tail closures and residue classes.