

Vector bundles on Fano threefolds and K3 surfaces

Arnaud Beauville

arXiv:1906.03594

<http://arxiv.org/abs/1906.03594v2>

Restriction to an anticanonical K3 surface

Let X be a Fano threefold over $\mathbf{C}$ and let $S \in |-K_X|$ be smooth, hence a K3 surface. For a simple vector bundle E on X , put $E_S = E|_S$. A component $\mathcal{M}_S$ of the moduli space of simple bundles on S carries Mukai's holomorphic symplectic form. The paper asks when the restriction map from a component $\mathcal{M}_X$ of bundles on X cuts out a Lagrangian subvariety.

Main theorem (Tyurin's theorem, as used and proved in Section 1). *Assume*

$$H^2(X, \mathcal{E}nd(E)) = 0 \quad (1)$$

for every $E \in \mathcal{M}_X$. Then $\mathcal{M}_X$ is smooth, every E_S is simple, and

$$\text{res} : \mathcal{M}_X \longrightarrow \mathcal{M}_S$$

is locally an isomorphism onto a Lagrangian subvariety of $\mathcal{M}_S$.

Beauville's contribution is to make (1) verifiable through curves and to construct concrete families to which the theorem applies, including five-dimensional Lagrangians in a moduli space birational to O'Grady's ten-dimensional hyperkähler manifold OG₁₀.

Ingredient I: the half-dimensional isotropic tangent space

Tensoring the anticanonical sequence by $\mathcal{E}nd(E)$ gives

$$0 \rightarrow \mathcal{E}nd(E) \otimes K_X \rightarrow \mathcal{E}nd(E) \rightarrow \mathcal{E}nd(E_S) \rightarrow 0. \quad (2)$$

The vanishing (1) and its Serre-dual group $H^1(X, \mathcal{E}nd(E) \otimes K_X)$ yield simplicity of E_S and an exact tangent-space sequence

$$0 \rightarrow H^1(X, \mathcal{E}nd(E)) \rightarrow H^1(S, \mathcal{E}nd(E_S)) \rightarrow H^2(X, \mathcal{E}nd(E) \otimes K_X) \rightarrow 0. \quad (3)$$

Serre duality identifies the dimensions of the outer terms, so the image has half the dimension of the middle space. Mukai's form is cup product followed by trace to $H^2(S, \mathcal{O}_S) \simeq \mathbf{C}$. For two classes coming from X , the product factors through $H^2(X, \mathcal{E}nd(E))$ and then $H^2(X, \mathcal{O}_X) = 0$. Thus the half-dimensional image is isotropic, hence Lagrangian. This linear-algebra picture is the proof's toy model; the real work is manufacturing (1) in geometric families.

Ingredient II: Serre's construction supplies the vanishing

Let $C \subset X$ be a smooth (or lci) curve and let L be ample with $(K_X \otimes L)|_C \simeq K_C$. Serre's construction gives a unique rank-two bundle

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow \mathcal{I}_C \otimes L \rightarrow 0, \quad E|_C \simeq N_{C/X}. \quad (4)$$

Proposition 1 assumes

$$H^1(C, N_{C/X}) = 0, \quad H^0(X, K_X \otimes L) \twoheadrightarrow H^0(C, K_C), \quad (5)$$

and concludes (1). Indeed, tensoring (4) by E^* gives $0 \rightarrow E^* \rightarrow \mathcal{E}nd(E) \rightarrow \mathcal{I}_C E \rightarrow 0$. Serre duality and the second condition in (5) kill $H^2(E^*)$; the first condition, together with Kodaira vanishing and ampleness of $-K_X$, kills $H^2(\mathcal{I}_C E)$. These two vanishings give $H^2(\mathcal{E}nd(E)) = 0$.

Proposition 2 adds the surjectivity $H^0(X, L) \twoheadrightarrow H^0(C, L|_C)$. Then the Serre bundles form an open subset of $\mathcal{M}_X$: deformation theory of the pair (E, s) reduces smoothness of the forgetful map to $H^1(X, E) = 0$, which is precisely that restriction condition. Hence moving curves produce an open family in a moduli component, not merely isolated examples.

Stability and numerical size

For the examples, $\text{Pic}(X) = \mathbf{Z}[\mathcal{O}_X(1)]$, the curve C is projectively normal, and L is $\mathcal{O}_X(1)$ or $\mathcal{O}_X(2)$. Lemma 1 proves that E and E_S are stable; in the second case one also assumes that C is not contained in a hyperplane. If $Z = C \cap S$ and $L = \mathcal{O}_X(j)$, the restricted Serre sequence

$$0 \rightarrow \mathcal{O}_S \rightarrow E_S \rightarrow \mathcal{I}_Z(j) \rightarrow 0 \quad (6)$$

makes the same stability mechanism visible on the K3 surface. For rank r and discriminant $\Delta_E = 2rc_2(E) - (r-1)c_1(E)^2$, Riemann–Roch and (1) give

$$\dim \mathcal{M}_X = \frac{1}{2}(-K_X \cdot \Delta_E) + 1 - r^2. \quad (7)$$

This formula predicts the half-dimension required by the Lagrangian conclusion and provides a numerical check on every family below.

Ingredient III: four families that realize the mechanism

The paper specializes to Picard-rank-one Fano threefolds and sufficiently general anticanonical sections with $\text{Pic}(S) = \mathbf{Z}[\mathcal{O}_S(1)]$, so the constructed bundles and their restrictions are stable.

Conics on a cubic threefold (Proposition 3). For $X \subset \mathbf{P}^4$ cubic and $S = X \cap Q$, conics have $H^1(N_{C/X}) = 0$. The resulting moduli spaces are

$$\mathcal{M}_X \simeq F(X), \quad \mathcal{M}_S \simeq S^{[2]}, \quad (8)$$

where $F(X)$ is the Fano surface of lines. If ℓ is the line residual to a conic in its plane section, restriction is the closed Lagrangian embedding $\ell \mapsto \ell \cap Q$.

Conics on an intersection of two quadrics (Proposition 4). For $X \subset \mathbf{P}^5$ of degree 4, the quadrics containing S form a net $\Pi \simeq \mathbf{P}^2$ with smooth discriminant sextic Δ . The double cover $\pi : \widehat{S} \rightarrow \Pi$ branched over Δ is a K3 surface. The quadrics containing X form a line $\ell \subset \Pi$, and $\Gamma = \pi^{-1}(\ell)$ has genus 2. The same normal-bundle vanishing identifies

$$\mathcal{M}_X \simeq \Gamma, \quad \mathcal{M}_S \simeq \widehat{S}, \quad (9)$$

with restriction equal to the natural embedding $\Gamma \hookrightarrow \widehat{S}$. Here every curve in a K3 surface is already Lagrangian, illustrating that Tyurin's theorem is most informative when $\mathcal{M}_S$ has higher dimension.

Normal elliptic curves on index-two Fanos (Proposition 6). For $X_d \subset \mathbf{P}^{d+1}$, $3 \leq d \leq 5$, a normal elliptic curve of degree $d+2$ produces a rank-two Ulrich bundle. Here

$$\dim \mathcal{M}_X = 5, \quad \dim \mathcal{M}_S = 10, \quad (10)$$

$\mathcal{M}_S$ is birational to OG_{10} , and restriction is injective and locally Lagrangian. Ulrich resolutions also prove the injectivity: an isomorphism after restriction lifts to a nonzero map between stable bundles on X , hence to an isomorphism.

Canonical curves on index-one Fanos (Proposition 8). For $X_{2g-2} \subset \mathbf{P}^{g+1}$, a canonical curve of genus $g+2$ with $H^1(N_{C/X}) = 0$ gives another local Lagrangian map to a moduli space birational to OG_{10} (with X general when $g=3$). For $g \geq 4$, the curve is the zero locus of a section of a globally generated twist of a stable bundle; the vanishing follows from the bundle construction. The quartic case $g=3$ is checked directly by the normal sequence of a complete intersection of three quadrics.

Proof roadmap and hardest point

The common chain, using both the normal-bundle vanishing and the restriction surjectivity in (5), is

$$\text{curve } C \xrightarrow{\text{Serre}} E \xrightarrow{\text{both conditions in (5)}} H^2(\mathcal{E}nd(E)) = 0 \xrightarrow{\text{Tyurin}} \text{Lagrangian restriction.} \quad (11)$$

The difficult step varies with the family: one must prove unobstructedness $H^1(N_C) = 0$ without losing stability or the needed restriction surjectivities. The examples use smooth Hilbert schemes, global generation of normal bundles, explicit normal sequences, and in genus 6 a nested-Hilbert-scheme argument. Once this curve-theoretic gate is passed, (2)–(3) make the symplectic conclusion short and formal.

Scope, transversality, and a useful warning

Tyurin's conclusion is local: it does not say that restriction is globally injective. Proposition 7 exhibits the failure for an index-one complete intersection of three quadrics. There $\mathcal{M}_X$ is an étale double cover of a discriminant curve, and restriction is its normalization map onto a singular curve in a K3 double cover, identifying pairs of points above the intersections of two discriminants.

For genus 6, the paper goes further. A general (2,3)-complete-intersection $X \subset \mathbf{P}^5$ contains canonical curves obtained from quintic del Pezzo surfaces, and a nested-Hilbert argument proves $H^1(N_C) = 0$ for all such canonical curves (Lemma 2). The hypersurface of Pfaffian cubics containing X maps birationally to $\mathcal{M}_X$ (Proposition 9). Meanwhile $\mathcal{M}_S$ has a rational Lagrangian fibration h to the space of cubics containing S , and $h \circ \text{res}$ is generically finite (Proposition 10): the constructed Lagrangian is weakly transverse to a general fiber. The degree-5 conic point and the detailed projective calculations are omitted; no global Lagrangian fibration by the moving $\mathcal{M}_X$ is claimed.