

The canonical heights for Jordan blocks of small eigenvalues, preperiodic points, and the arithmetic degrees

Kaoru Sano

arXiv:1712.07533

<http://arxiv.org/abs/1712.07533v1>

The spectral-gap question

Let X be a smooth projective variety over $\overline{\mathbf{Q}}$, let $f : X \rightarrow X$ be a surjective endomorphism, and fix an ample divisor H . Write $V_H \subset \text{Pic}(X)_{\mathbf{Q}}$ for the span of $(f^n)^*H$, $n \geq 0$, and $\overline{V}_H \subset N^1(X)_{\mathbf{Q}}$ for its image modulo numerical equivalence. For $P \in X(\overline{\mathbf{Q}})$, the arithmetic degree is

$$\alpha_f(P) = \lim_{n \rightarrow \infty} \max\{h_H(f^n(P)), 1\}^{1/n}.$$

It is always 1 on a preperiodic orbit. The problem is to recognize when slow height growth forces the orbit itself to be finite.

Main theorem (Theorem 1.1). *Assume that no eigenvalue of $f^* : \overline{V}_H \rightarrow \overline{V}_H$ has complex absolute value 1, and put*

$$\mu_H(f) = \min\{|\lambda| : \lambda \text{ is an eigenvalue of } f^*|_{\overline{V}_H}, |\lambda| \geq 1\}.$$

Then, for every $P \in X(\overline{\mathbf{Q}})$, the following are equivalent:

$$P \text{ is preperiodic}, \quad \alpha_f(P) < \mu_H(f), \quad \alpha_f(P) = 1.$$

Thus the numerical spectrum has a genuine gap above the finite orbits. Unlike the polarized case, the cyclic divisor space may contain nontrivial Jordan blocks and eigenvalues of modulus less than one; controlling those contracting directions is the new issue.

Canonical heights in the contracting direction

Choose a Jordan chain of divisors $D_0, \dots, D_l$ with $f^*D_j \sim D_{j-1} + \lambda D_j$ and $D_{-1} = 0$. For $|\lambda| > 1$, the usual forward normalization removes the bounded errors in the Weil-height relations. It diverges when $0 < |\lambda| < 1$. Sano instead works on

$$\varprojlim_f X(\overline{\mathbf{Q}}) = \{(P_m)_{m \in \mathbf{Z}} : f(P_m) = P_{m+1}\},$$

where surjectivity supplies backward histories and the right shift is a bijection.

Let Λ be the Jordan block with eigenvalue λ , let $h = (h_{D_0}, \dots, h_{D_l})^t$, and let $E = h \circ R - \Lambda^{-1}h$ be the bounded functoriality error. When $|\lambda| < 1$, the correction

$$\widehat{h} = h + \sum_{n \geq 0} \Lambda^{n+1}(E \circ R^n)$$

converges uniformly because $\|\Lambda^n\| \ll n^l |\lambda|^{n-l}$. It is the unique bounded perturbation satisfying $\widehat{h} \circ R = \Lambda^{-1}\widehat{h}$. Translating back to the left shift shows that every ordinary height $h_{D_j}(f^n(P))$ in a contracting Jordan block is bounded along the forward orbit (Theorem 1.2 and Proposition 3.3).

How the gap forces preperiodicity

1. **Separate the numerical Jordan blocks.** Lemma 4.2 gives an integral polynomial annihilating $\text{Pic}(X)$, so V_H is finite-dimensional. Over $\mathbf{C}$ it splits into Jordan blocks. Blocks visible in $\overline{V}_H$ have modulus either > 1 or < 1 by the hypothesis; blocks killed by $V_H \rightarrow \overline{V}_H$ consist of algebraically trivial divisors.

2. **An expanding coordinate detects exponential growth.** Suppose a canonical-height coordinate in a block with $|\lambda| > 1$ is nonzero, and choose the first nonzero coordinate in its Jordan chain. Its value at $f^n(P)$ is exactly λ^n times its initial value. Comparison with the ample height gives $\alpha_f(P) \geq |\lambda| \geq \mu_H(f)$. Hence $\alpha_f(P) < \mu_H(f)$ makes every expanding canonical-height coordinate vanish, so the associated ordinary heights are bounded.
3. **All remaining contributions are harmless.** The contracting-block heights are bounded by the inverse-limit construction. Heights of the algebraically trivial blocks are $o(h_H)$. Since H is a linear combination of the Jordan-chain divisors, along the orbit one gets

$$h_H(f^n(P)) - C \leq o(h_H(f^n(P))).$$

Therefore $h_H(f^n(P))$ is bounded. The orbit is defined over one finite extension of the field of definition of X , f , and P ; Northcott's theorem then makes it finite.

Here the $o(h_H)$ estimate is genuinely sufficient. If the ample heights along a subsequence escaped to infinity, dividing the displayed inequality by h_H would give $1 \leq o(1)$, a contradiction. This is why the blocks that disappear under numerical equivalence need not have bounded heights of their own.

The reverse implications are immediate: a finite orbit has arithmetic degree 1, while the spectral hypothesis ensures $1 < \mu_H(f)$.

Model, application, and scope

For $f(z) = z^d$ on $\mathbf{P}^1$, with an integer $d \geq 2$, V_H has the single eigenvalue d . Roots of unity together with 0 and ∞ are preperiodic and have arithmetic degree 1; every other algebraic point has arithmetic degree d . The theorem replaces this one expanding coordinate by an arbitrary numerical Jordan decomposition, and its inverse-limit height handles directions that contract rather than expand.

For the main application (Proposition 5.4), let δ_f be the first dynamical degree and let $\deg f = [\overline{\mathbf{Q}}(X) : f^*\overline{\mathbf{Q}}(X)]$. Assume that the numerical cyclic space has dimension at most two and $\delta_f > (\deg f)^2$. In dimension one the map is polarized on this space and the conclusion is standard. In dimension two, if λ is the second eigenvalue, the integral push-pull maps give

$$|\delta_f \lambda| = |\det(f^*|_{\overline{V}_H})| \leq (\deg f)^2,$$

so $|\lambda| < 1$ and $\mu_H(f) = \delta_f$. Thus every wandering point has arithmetic degree δ_f , proving the Kawaguchi–Silverman prediction in this setting.

Example 5.7 shows the contracting phenomenon concretely on $E \times E$: after composing a suitable automorphism with multiplication by 2, the two numerical eigenvalues are $28 + 16\sqrt{3} > 1$ and $28 - 16\sqrt{3} < 1$. Theorem 1.1 says that preperiodicity is exactly the strict inequality $\alpha_g(P) < 28 + 16\sqrt{3}$. By contrast, an eigenvalue of modulus one is a real obstruction: translation factors can create wandering points of arithmetic degree 1. The hardest point is therefore not Northcott, but constructing a canonical normalization for the small-eigenvalue Jordan blocks so that every non-expanding height contribution is controlled.