A case of the Dynamical Andre-Oort Conjecture

Dragos Ghioca, Holly Krieger, Khoa Nguyen

arXiv:1404.6011

http://arxiv.org/abs/1404.6011v1

A dynamical unlikely-intersection problem

Fix $d \geq 2$ and write

$$f_t(z) = z^d + t, \qquad \mathcal{P}_d = \{t \in \mathbf{C} : 0 \text{ is preperiodic for } f_t\}.$$

The finite critical point of f_t is 0, so f_t is postcritically finite (PCF) exactly when $t \in \mathcal{P}_d$. The dth multibrot set

$$\mathcal{M}_d = \{t : \{f_t^n(0)\}_{n\geq 0} \text{ is bounded}\}$$

is the connectedness locus of this family. The question is an unlikely-intersection problem: when can the graph of a polynomial h meet the special set $\mathcal{P}_d \times \mathcal{P}_d$ infinitely often?

Main theorem (Theorem 1.1)**.** *Let $h \in \mathbf{C}[z]$ be nonconstant. If there are infinitely many $t \in \mathbf{C}$ for which both f_t and $f_{h(t)}$ are PCF, then*

$$h(z) = \zeta z \qquad \textit{with} \qquad \zeta^{d-1} = 1.$$

Thus the only polynomial graphs with infinitely many simultaneous PCF parameters are the visible rotational symmetries of the multibrot family.

From arithmetic coincidence to set invariance

1. Upgrade infinitely many parameters to all parameters. Every PCF parameter is algebraic. Since h takes infinitely many algebraic arguments to algebraic values, interpolation first puts its coefficients in $\overline{\mathbf{Q}}$. The imported unlikely-intersection theorem quoted as Theorem 2.1 then applies to the two families $z^d + t$ and $z^d + h(t)$: the infinite coincidence implies

$$t \in \mathcal{P}_d \iff h(t) \in \mathcal{P}_d \qquad \text{for every } t \in \mathbf{C}.$$

This is the arithmetic input. Its proof in the cited earlier work uses equidistribution of small points together with variation of canonical heights; the present paper does not reprove it.

2. Recover the whole boundary from PCF points. The boundary $\partial\mathcal{M}_d$ is characterized by the property that every neighborhood contains infinitely many distinct PCF parameters. The equivalence above, continuity of h, and the open mapping theorem therefore give

$$h^{-1}(\partial\mathcal{M}_d) = \partial\mathcal{M}_d.$$

Because $\mathbf{C} \setminus \mathcal{M}_d$ is connected and h has full degree near infinity, the exterior is totally invariant as well; hence $h^{-1}(\mathcal{M}_d) = \mathcal{M}_d$.

If h is linear, it is an affine symmetry of $\mathcal{M}_d$. Proposition 2.2 computes all such symmetries from the Laurent expansion at infinity of the parameter Böttcher map $\Phi : \mathbf{C} \setminus \mathcal{M}_d \to \mathbf{C} \setminus \overline{\mathbf{D}}$: they are precisely $z \mapsto \zeta z$ with $\zeta^{d-1} = 1$.

Suppose instead that $\deg h \geq 2$. The Julia set J_h is the minimal closed set with at least three points that is totally invariant under h, so $J_h \subset \partial\mathcal{M}_d$. The connected exterior of $\mathcal{M}_d$ has no bounded complementary component, giving $K_h \subset \mathcal{M}_d$. Conversely, total invariance and boundedness of $\mathcal{M}_d$ keep every orbit starting there bounded, so $\mathcal{M}_d \subset K_h$. Thus $K_h = \mathcal{M}_d$. The remaining task is the paper's independent complex-dynamical obstruction:

$$\mathcal{M}_d \text{ is not the filled Julia set of any polynomial.} \qquad \text{(Theorem 1.2)}$$

Why a multibrot set cannot be a filled Julia set

The paper gives two proofs; the shorter algebraic one runs through Theorem 4.4. If polynomials P, g have degree at least two and either $P(S^1) = J_g$ or $P(\mathbf{D})$ is a Fatou component of g, then J_g is a circle.

3. Turn a relation on the circle into an invariant algebraic curve. For a nonconstant rational map q, embed its image of S^1 in $\mathbf{P}^1 \times \mathbf{P}^1$ by

$$C_q = \{(q(x), \overline{q}(y)) : xy = 1\}.$$

If $r \circ q(u) = q(w)$ for infinitely many $(u, w) \in (S^1)^2$, then the two irreducible curves C_q and $C_{r\circ q}$ meet infinitely often. They are equal, so C_q is invariant under $(r, \overline{r})$.

When q is a nonconstant *polynomial* and r is a non-exceptional polynomial of degree at least two, Proposition 4.8 applies the Medvedev–Scanlon classification of invariant curves to forbid this. Indeed, that classification would express C_q as the image of a graph $y = \psi(x)$ (or its transpose) under polynomial semiconjugacies. Such a presentation has (∞, ∞) as its unique point over first coordinate ∞, whereas the definition of C_q gives $(\infty, \overline{q}(0))$. Hence r must be

exceptional: linearly conjugate to a powering map or to a signed Chebyshev polynomial. Its Julia set is then a circle or a line segment.

4. Feed a Fatou component into that rigidity. In the Julia-set case of Theorem 4.4 the lemma is used with $q = P$, so the polynomial hypothesis is satisfied. If $P(\mathbf{D})$ is a bounded Fatou component, Sullivan's no-wandering-domain theorem makes an eventual image periodic. After replacing g by an iterate, the boundaries of $g \circ P(\mathbf{D})$ and $g^2 \circ P(\mathbf{D})$ coincide in an infinite set. The preceding invariant-curve argument makes g exceptional; here Proposition 4.8 is used with the polynomial $q = g \circ P$. The Chebyshev case is impossible because its only Fatou component is unbounded, so J_g is a circle.

Now the main hyperbolic component of $\mathcal{M}_d$ is

$$\mathcal{H}_d = \{c : f_c \text{ has an attracting fixed point}\} = \{\alpha - \alpha^d : |\alpha| < d^{-1/(d-1)}\}.$$

If $\mathcal{M}_d = K_g$, then $\mathcal{H}_d$ is the polynomial image of a disk and is a Fatou component of g. Theorem 4.4 would make $\partial\mathcal{M}_d = J_g$ a circle, a contradiction. This proves Theorem 1.2 and closes the nonlinear branch of Theorem 1.1.

What the other proof adds

The paper's first proof of Theorem 1.2 makes the incompatibility visible in external-angle combinatorics. If one assumes $K_h = \mathcal{M}_d$ and writes $D = \deg h$, the Böttcher coordinates can be normalized so that h is monic and the parameter coordinate Φ conjugates h on the exterior of $\mathcal{M}_d$ to $z \mapsto z^D$. Proposition 3.5 proves that, for every $m \geq 2$, the parameter rays with angles

$$\frac{1}{d^m - 1} \quad \text{and} \quad \frac{d}{d^m - 1}$$

land together on the boundary of the main hyperbolic component. Exterior conjugacy would therefore make the rays of angles $D^k/(d^m - 1)$ and $dD^k/(d^m - 1)$ land together for every $k \geq 0$.

Choose $m = m(k)$ by

$$d^m - 1 \leq D^k(d^2 - 1) < d^{m+1} - 1.$$

For large k, Bang's theorem supplies a primitive prime divisor $p \mid d^m - 1$ with $p \nmid D$. At a common landing point, the corresponding dynamic rays lie in one cycle under multiplication by d; if its exact period is $n \mid m$, this gives a congruence

$$D^k(1 - d^{rn+1}) \equiv 0 \pmod{d^m - 1} \qquad (0 \leq r < m/n).$$

Primitivity of p forces $rn + 1 = m$, hence $n = 1$. The landing point must therefore lie on the main hyperbolic component. Its wake geometry then forces D^k to be one of the two endpoint numerators $(d^m - 1)/(d^2 - 1)$ or $d(d^m - 1)/(d^2 - 1)$. This is impossible because the primitive divisor p divides $d^m - 1$ but divides neither D nor $d^2 - 1$. Thus the ray proof reaches the same obstruction by combining a precise landing-pair theorem with primitive divisors.

For $d = 2$ the conclusion is especially rigid: $\zeta = 1$, so the graph must be the diagonal. This toy case removes the visible rotations but not the two hard inputs—arithmetic equidistribution in Theorem 2.1 and invariant-curve rigidity in Theorem 4.4. The result treats polynomial graphs in the unicritical parameter plane; it does not prove the full dynamical André–Oort conjecture for arbitrary curves or arbitrary rational maps.