

Geometry of moduli spaces of rational curves in linear sections of Grassmannian $\mathrm{Gr}(2, 5)$

Kiryong Chung, Jaehyun Hong, Sanghyeon Lee

arXiv:1608.02275

<http://arxiv.org/abs/1608.02275v2>

Low-degree curves on the degree-five Fano sections

Let $G = \mathrm{Gr}(2, 5) \subset \mathbf{P}^9$ be the Pluecker Grassmannian and, over $\mathbf{C}$, let

$$Y_5^m = G \cap H_1 \cap \cdots \cap H_{6-m}, \quad 2 \leq m \leq 6, \quad (1)$$

for general hyperplanes. Thus Y_5^m is a smooth Fano m -fold of degree 5 and coindex 2. Write $R_d(Y_5^m)$ for the moduli space of smooth rational curves of Pluecker degree d , and $H_d(Y_5^m)$ for its Hilbert closure with Hilbert polynomial $dt + 1$.

Main theorem (Chung–Hong–Lee, Theorems 1.1 and 1.2). *For $d \leq 3$ and $2 \leq m \leq 6$, the spaces $R_d(Y_5^m)$ are rational. Moreover the Hilbert schemes $H_2(Y_5^m)$ of conics are smooth and irreducible for $m = 4, 5$; their dimensions are respectively 7 and 10.*

For $m = 2$ the source gives an explicit componentwise description: there are ten lines; the conic locus is five copies of $\mathbf{P}^1 \setminus \{0, 1, \infty\}$; and the cubic locus is one $\mathbf{P}^2$ minus four lines together with four copies of $\mathbf{P}^2 \setminus \mathbf{P}^1$.

Ingredient I: split the universal bundle

A degree- d map $f : \mathbf{P}^1 \rightarrow G$ pulls back the dual universal bundle as

$$f^*\mathcal{U}^\vee \simeq \mathcal{O}_{\mathbf{P}^1}(d_1) \oplus \mathcal{O}_{\mathbf{P}^1}(d_2), \quad d_1 + d_2 = d. \quad (2)$$

For $d = 1, 2, 3$, the possible splittings turn the family of lines in $\mathbf{P}^4$ parameterized by the curve into a small rational normal scroll. Consequently a line in G has a *vertex* $p \in \mathbf{P}^4$; a conic has an *envelope* $\mathbf{P}^3 \subset \mathbf{P}^4$ containing all its lines; and a general twisted cubic has an *axis*, a line meeting every member of the family. These define

$$\eta_1 : R_1(G) \rightarrow \mathbf{P}^4, \quad \eta_2 : R_2(G) \dashrightarrow \mathrm{Gr}(4, 5) = \mathbf{P}^4, \quad \eta_3 : R_3(G) \dashrightarrow \mathrm{Gr}(2, 5). \quad (3)$$

The general fibers are low-dimensional Grassmannian parameter spaces inside the corresponding Schubert cycles.

Ingredient II: intersect the fibers, not the moduli space

For Y_5^3 , known descriptions give

$$H_1(Y_5^3) \simeq \mathbf{P}^2, \quad H_2(Y_5^3) \simeq \mathbf{P}^4, \quad H_3(Y_5^3) \simeq \mathrm{Gr}(2, 5). \quad (4)$$

An incidence correspondence with linear subspaces containing the curve then proves $R_d(Y_5^m)$ irreducible for $m \geq 3$. For $m \geq 4$, imposing the hyperplanes in (1) cuts the vector spaces in the fibers of (3). The resulting birational models are

m	R_1	R_2	R_3
6	$\mathrm{Gr}(2, 4)/\mathbf{P}^4$	$\mathrm{Gr}(3, 6)/\mathbf{P}^4$	$\mathrm{Gr}(4, 7)/\mathrm{Gr}(2, 5)$
5	$\mathbf{P}^2/\mathbf{P}^4$	$\mathrm{Gr}(3, 5)/\mathbf{P}^4$	$\mathrm{Gr}(4, 6)/\mathrm{Gr}(2, 5)$
4	$\mathbf{P}^4$	$\mathbf{P}^3/\mathbf{P}^4$	$\mathbf{P}^4/\mathrm{Gr}(2, 5)$,

(5)

where “fiber/base” denotes a generically Grassmannian bundle. Since both the bases and fibers are rational, (5), together with (4) and the del Pezzo surface calculation, proves the first theorem.

Ingredient III: control the conic boundary

Rationality only sees the open locus. To understand its Hilbert compactification, let $\mathcal{U}$ be the universal rank-4 bundle on $\mathrm{Gr}(4, 5)$ and, for $m = 5$ or 4 , put

$$\mathcal{K} = \ker \left(\bigwedge^2 \mathcal{U} \longrightarrow \mathcal{O}^{\oplus(6-m)} \right), \quad S(Y) = \mathrm{Gr}(3, \mathcal{K}). \quad (6)$$

Over $\mathrm{Gr}(4, 5)$, the relative Hilbert scheme of conics in $\mathrm{Gr}(2, \mathcal{U})$ maps to $H_2(G)$ and gives a common resolution: one morphism blows up the two flag loci corresponding to the two types of planes in G , while the other blows up the locus of conics lying in a $\sigma_{2,2}$ -plane.

Intersecting this diagram with $S(Y)$ is subtle. Lemma 5.6 identifies tangent spaces and proves that the intersection with the two flag loci is scheme-theoretic, so its blow-up is the proper transform. The restricted normal bundle in the exceptional direction is $\mathcal{O}(-1)$; the Fujiki–Nakano criterion therefore contracts the other ruling smoothly. Let $\tilde{S}(Y)$ denote this proper transform, equivalently the common blow-up of $S(Y)$ along the intersected flag loci. Thus

$$S(Y) \longleftarrow \tilde{S}(Y) \longrightarrow H_2(Y) \quad (7)$$

is one blow-up followed by one blow-down. Irreducibility comes from the ambient diagram, and the local-complete-intersection argument used for lines shows the Hilbert scheme is reduced. Proposition 6.7 repeats the construction for the second hyperplane.

Toy model, hardest point, and scope

For the threefold Y_5^3 , the map in (6) has rank 3 and Y_5^3 contains no planes. Hence no boundary modification is needed: $H_2(Y_5^3) \simeq S(Y) \simeq \mathrm{Gr}(4, 5) = \mathbf{P}^4$. This is the clean model for (7): the envelope already determines the conic. In dimensions four and five, planes create competing degenerations, and the hardest point is proving that their scheme-theoretic centers and normal directions permit a smooth flip-like replacement.

Generality of the hyperplane sections and the ground field $\mathbf{C}$ are part of the setup. The rationality theorem concerns the open spaces of smooth curves, while the smoothness assertion concerns only conic Hilbert schemes for $m = 4, 5$. The paper also records compactifications for lines and twisted cubics in G from earlier work, but does not claim analogous new smoothness theorems for every degree and dimension.