

Bifurcations, intersections, and heights

Laura DeMarco

arXiv:1412.2708

<http://arxiv.org/abs/1412.2708v3>

Three faces of a marked orbit

Let V be an irreducible quasi-projective complex curve, $k = \mathbb{C}(V)$, and $f: V \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ an algebraic family of degree $d \geq 2$. A marked point $a \in \mathbb{P}^1(k)$ is *stable* when the maps $t \mapsto f_t^n(a(t))$ form a normal family on V ; it is *preperiodic* when two of these maps agree identically. Isotriviality allows a finite branched base change and a moving Möbius conjugacy that make both the map and marked point constant.

Main theorem (DeMarco, Theorems 1.1, 1.2, and 1.4). *If (f, a) is stable, then it is isotrivial or preperiodic. In particular, if f is non-isotrivial and V is a curve, then for every $a \in \mathbb{P}^1(k)$ the following are equivalent:*

$$(f, a) \text{ is stable,} \quad \widehat{h}_f(a) = 0, \quad (f, a) \text{ is preperiodic.}$$

Moreover, only finitely many $a \in \mathbb{P}^1(k)$ are stable. More quantitatively, if f is not isotrivial over k , there is $b > 0$ for which $\{a : \widehat{h}_f(a) < b\}$ is finite.

This identifies an analytic condition in the parameter variable with an algebraic orbit relation and an arithmetic height condition. The paper also applies it to simultaneous preperiodicity: for a family of dimension $N > 0$ in moduli and any $r \leq N$ marked points, their common preperiodic locus is Zariski dense (Theorem 1.6).

Ingredient I: small height forces a constant family

For $g_n(t) = f_t^n(a(t))$, the function-field canonical height is

$$\widehat{h}_f(a) = \lim_{n \rightarrow \infty} \frac{\deg g_n}{d^n}, \quad (1)$$

for $a \in \mathbb{P}^1(k)$ (the paper includes the factor $1/\deg p_a$ after a finite cover $p_a: W \rightarrow V$). To prove finiteness below a positive height threshold, suppose instead that $\widehat{h}_f(a_m) \rightarrow 0$. The height axioms produce increasingly long initial orbit segments whose maps to $\mathbb{P}^1$ have uniformly bounded degree.

After marking three forward-invariant points and conjugating them to $0, 1, \infty$, each iterate outside its finite inverse image is a map to $\mathbb{P}^1 \setminus \{0, 1, \infty\}$. Lemma 2.4, via Riemann–Hurwitz, gives only finitely many such nonconstant maps from a fixed punctured curve. A long orbit segment therefore contains $2d + 2$ consecutive constant, distinct iterates. Two degree- d rational maps agreeing on $2d + 1$ points coincide (Lemma 2.5), forcing the conjugated family to be constant. Lemma 2.1 removes the residual finite monodromy. This contradicts non-isotriviality and proves Theorem 1.2.

Ingredient II: control at a puncture

Near a missing parameter $t = 0$, choose a homogeneous lift $F_t: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ and a lift A_t of $a(t)$. Put

$$a_n = \text{ord}_{t=0} F_t^n(A_t), \quad F_n(t) = t^{-a_n} F_t^n(A_t), \quad G_n(t) = d^{-n} \log \|F_n(t)\|. \quad (2)$$

If $q = \text{ord}_0 \text{Res}(F_t)$, the resultant identity gives, for small $t \neq 0$,

$$\alpha |t|^q \leq \frac{\|F_t(z)\|}{\|z\|^d} \leq \beta. \quad (3)$$

Thus $0 \leq a_{n+1} - da_n \leq q$. This bounded defect makes G_n converge locally uniformly on the punctured disk to a continuous G with $G(t) = o(\log |t|)$ (Proposition 3.1).

If the marked orbit is stable, G is harmonic off the puncture; the sublogarithmic estimate removes the singularity. Rescaling A_t by a holomorphic factor normalizes $G \equiv 0$ (Corollary 3.4), after which Lemma 3.5 bounds every F_n above and away from zero on compact annuli. This is the local estimate that survives degeneration of f_t at the boundary of V .

Ingredient III: normality cannot hide degree growth

Reduce Theorem 1.1 to a curve by slicing. Assume that (f, a) is stable and non-isotrivial but that $\deg g_n$ is unbounded. A normal subsequence g_{n_j} converges on V . The argument principle then forces, near some puncture, unbounded numbers of both zeros and poles of g_{n_j} (Lemma 4.1).

On an annulus $r_0 \leq |t| \leq r_1$, factor the two coordinates of F_{n_j} so that P_{n_j} and Q_{n_j} contain $\kappa(n_j) \rightarrow \infty$ zeros inside $|t| < r_0$. The normalization from Ingredient II gives uniform two-sided bounds for F_{n_j} . Hence the remaining factors are at most $(r_1 - r_0)^{-\kappa(n_j)}$ on the outer circle but one is at least $(2r_0)^{-\kappa(n_j)}$ on the inner circle. Choosing $2r_0 < r_1 - r_0$ contradicts the maximum principle. Therefore the degrees are bounded, and Proposition 1.8 makes a preperiodic.

From instability to dense intersections

The same analytic bridge yields Theorem 1.6. Proposition 5.1 says that every open set meeting the bifurcation locus contains infinitely many parameters where the marked point lands on a repelling cycle. Indeed, choose three holomorphically moving repelling cycles outside the initial orbit; Montel's theorem turns failure of normality into an iterate hitting one of them. For $r \leq N$ marked points, impose these preperiodic relations successively. Persistence of each repelling cycle makes the new relation a codimension-one condition in moduli, so after r steps a nonempty locus of moduli dimension at least $N - r$ remains. Starting outside an arbitrary proper closed subset proves Zariski density. For an elliptic family, quotienting by $P \sim -P$ turns multiplication by 2 into a Lattès map and torsion into preperiodicity, giving the paper's torsion-specialization application.

A scale model and the hard point

For $Q_t(z) = z^2 + t$ over $\mathbb{C}(t)$, every $a \neq \infty$ satisfies $\deg Q^n(a) = 2^{n-1} \deg Q(a)$, while ∞ is the sole rational preperiodic point. Every finite constant $a \in \mathbb{C}$ has height $1/2$, so the optimal threshold in Theorem 1.2 is $b = 1/2$. This elementary model displays the degree-growth quantity in (1), but it removes the paper's hardest issue: in a general family, zeros and poles can be absorbed at degenerate parameters. Estimates (2)–(3) show that their loss is too controlled to coexist with normality.

The stability theorem itself permits higher-dimensional parameter spaces through slicing, but the height equivalence and finiteness statement above are formulated for a curve base. The argument is specific to rational maps on $\mathbb{P}^1$; the final unlikely- intersection conjecture is only proved in two of its three directions.