The tropicalization of the moduli space of curves

Dan Abramovich, Lucia Caporaso, Sam Payne

arXiv:1212.0373

http://arxiv.org/abs/1212.0373v2

Problem and setup

Work over an algebraically closed field k with the trivial valuation, and in the stable range $2g-2+n>0$. A stable pointed curve has a weighted dual graph G: vertices record irreducible components and their genera, edges record nodes, and legs record marked points. An extended tropical curve is such a graph with an edge length in $\mathbf{R}_{>0}\cup\{\infty\}$.

For a fixed stable graph, the cone

$$\sigma_G=(\mathbf{R}_{\geq 0})^{E(G)},\qquad \overline{\sigma}_G=(\mathbf{R}_{\geq 0}\cup\{\infty\})^{E(G)}$$

records ordinary and extended edge lengths. Gluing these cones along faces corresponding to graph contractions, and quotienting by graph automorphisms, gives $M_{g,n}^{\mathrm{trop}}$ and its compactification $\overline{M}_{g,n}^{\mathrm{trop}}$.

The algebraic boundary has the parallel stratification $\overline{\mathcal{M}}_{g,n}=\coprod_G \mathcal{M}_G$: the stratum $\mathcal{M}_G$ has codimension $|E(G)|$, and one stratum specializes to another exactly when the associated graphs are related by contraction. On the tropical side $\sigma_G^\circ/\operatorname{Aut}(G)$ has dimension $|E(G)|$. Thus dimensions and codimensions are exchanged, as expected for a boundary complex.

A point of the Berkovich analytification of the coarse moduli space $\overline{M}_{g,n}$ is represented, after extension, by a stable curve over a valuation ring. If a node of the special fiber has local equation $xy=f_e$, its naive tropicalization assigns the corresponding edge the length $\mathrm{val}(f_e)$.

Main theorem (Theorems 1.2.1 and 1.2.2)**.** *There are natural isomorphisms of generalized cone complexes with integral structure*

$$\Phi_{g,n}:\Sigma(\overline{\mathcal{M}}_{g,n})\xrightarrow{\sim} M_{g,n}^{\mathrm{trop}},\qquad \overline{\Phi}_{g,n}:\overline{\Sigma}(\overline{\mathcal{M}}_{g,n})\xrightarrow{\sim}\overline{M}_{g,n}^{\mathrm{trop}}.$$

If p is the canonical retraction of $\overline{M}_{g,n}^{\mathrm{an}}$ onto the skeleton, then

$$\mathrm{Trop}=\overline{\Phi}_{g,n}\circ p.$$

Hence tropicalization is continuous, proper, and surjective. It also commutes with the algebraic and tropical forgetful, clutching, and gluing maps.

The skeleton is constructed from the stack $\overline{\mathcal{M}}_{g,n}$ but lies inside the analytification of its coarse moduli space; this distinction matters because the coarse space need not itself be toroidal.

Proof mechanism

1. Extend skeletons from schemes to toroidal stacks. For a toroidal Deligne–Mumford stack $\mathcal{X}$ with coarse moduli space X, choose an étale atlas $V\to\mathcal{X}$ without self-intersections and set $V_2=V\times_{\mathcal{X}}V$. Thuillier's retractions on V and V_2 descend through the topologically right-exact groupoid $V_2^i\rightrightarrows V^i\to X^i$. Proposition 6.1.4 thereby produces a canonical deformation retraction

$$p_{\mathcal{X}}:X^i\longrightarrow\Sigma(\mathcal{X}),\qquad \Sigma(\mathcal{X})=\operatorname{colim}\big(\Sigma(V_2)\rightrightarrows\Sigma(V)\big).$$

Functoriality for sub-toroidal morphisms is Proposition 6.1.8.

2. Read each cone and its quotient from a boundary stratum. At a geometric point w of a stratum, effective boundary divisors form a sharp characteristic monoid S_w; its dual is a rational polyhedral cone σ_w. The étale fundamental group of the stratum permutes the local boundary branches, with finite image H_w. Proposition 6.2.6 gives the decisive decomposition

$$\Sigma(\mathcal{X})=\coprod_W \sigma_W^\circ/H_W,$$

and its extended analogue. This accounts for self-intersections and monodromy rather than pretending that every stratum contributes an unquotiented cone. Concretely, a small toric chart trivializes the boundary branches and contributes one cone. Two points of its relative interior descend to the same point of the skeleton exactly when a monodromy operator relates them. Lower-dimensional faces are supplied by strata whose closures contain the original stratum. This upgrades the displayed disjoint union to a genuine colimit of cone complexes.

3. Specialize to stable curves. The boundary stratum $\mathcal{M}_G$ consists of curves with dual graph G. A versal deformation has one normal-crossings divisor $D_e=(f_e)$ for every node, so its local skeleton is $\overline{\sigma}_G$. The cover that

labels the components, nodes, and branches has Galois group $\mathrm{Aut}(G)$; therefore Proposition 7.2.1 identifies the monodromy group with the image of $\mathrm{Aut}(G)$ acting on $E(G)$. Thus

$$\overline{\Sigma}(\overline{\mathcal{M}}_{g,n}) = \coprod_G \overline{\sigma}_G^\circ / \mathrm{Aut}(G),$$

exactly the same stratumwise description as extended tropical moduli.

4. Check that the gluing data also agree. In the algebraic compactification, the closure relations among strata are precisely contractions $G \to G'$. On both sides these induce the same face maps $\overline{\sigma}_{G'} \hookrightarrow \overline{\sigma}_G$. Consequently the skeleton and tropical moduli space are colimits of the same diagram of extended cones, proving the isomorphism with its integral structure, not merely a set-theoretic bijection. The graph automorphisms appear as the self-maps of a cone in this diagram, while every proper face inclusion is induced by contracting the complementary set of edges. No extra identifications remain.

5. Identify the analytic retraction with naive tropicalization. Near a node q_e, a family is étale-locally $xy = f_e$. The boundary divisors (f_e) form a basis of the characteristic monoid, and Thuillier's monomial valuation sends this basis element to $\mathrm{val}(f_e)$. The retraction therefore records exactly the same metric graph as the naive rule. Properness and surjectivity follow from the retraction and the cone-complex isomorphism.

Tautological maps and a local model

Forgetting a marking may destabilize a rational component. If it lies between two nodes with equations $xy = f_1$ and $xy = f_2$, stabilization replaces them by one node with parameter $f_1 f_2$; hence the new edge length is

$$\mathrm{val}(f_1 f_2) = \mathrm{val}(f_1) + \mathrm{val}(f_2),$$

which is exactly tropical stabilization: remove the two-valent vertex and merge its adjacent edges. If forgetting the marking instead creates a one-pointed rational tail, stabilization merely deletes that tail and restricts the old length function. Clutching or gluing marked points creates a node already present on the generic fiber, with local equation $xy = 0$; its tropical edge has length $\mathrm{val}(0) = \infty$. All other nodes keep their original smoothing parameters. These local calculations drive the three commutative diagrams of Theorem 1.2.2.

The one-node family $xy = t$ is the basic toy model. A unit t gives length zero and contracts the edge; $0 < \mathrm{val}(t) < \infty$ gives a finite tropical edge; and $t = 0$ gives an infinite edge. The full proof globalizes this dictionary while retaining graph automorphisms and stack monodromy.

Allowing the value ∞ is therefore not cosmetic. It compactifies each cone in precisely the direction in which a node persists on the generic fiber, and it makes clutching and gluing land inside the same extended tropical moduli space rather than on an ad hoc boundary.

Difficulty and scope

The hard point is not assigning a length to one node, but showing that all local cones descend canonically through an étale groupoid and glue with exactly the automorphism quotients and contraction maps of tropical moduli. The paper develops this for arbitrary toroidal Deligne–Mumford stacks before applying it to curves. Its result is a statement about the Berkovich skeleton over a trivially valued algebraically closed base and the moduli of stable curves; it does not identify every tropical moduli problem with an algebraic skeleton.